

## **Annex VI report**

# **PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING**

**Substance Name:** Abamectin

**EC Number:**

**CAS Number:** 71751-41-2

**Submitted by:** Bureau REACH, RIVM, The Netherlands, [bureau-reach@rivm.nl](mailto:bureau-reach@rivm.nl)

**Version:** Draft 2.1 June 2009



# CONTENTS

1	IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES .....	6
1.1	Name and other identifiers of the substance .....	6
1.2	Composition of the substance .....	6
1.3	Physico-chemical properties .....	9
2	MANUFACTURE AND USES .....	11
3	CLASSIFICATION AND LABELLING .....	11
3.1	Classification in Annex I of Directive 67/548/EEC.....	11
3.2	Self classification(s) .....	11
4	ENVIRONMENTAL FATE PROPERTIES .....	13
4.1	Degradation .....	13
4.1.1	Stability .....	13
4.1.2	Biodegradation .....	14
4.1.3	Summary and discussion of persistence .....	16
4.2	Environmental distribution .....	17
4.2.1	Adsorption/desorption .....	17
4.2.2	Volatilisation .....	17
4.2.3	Distribution modelling .....	17
4.3	Bioaccumulation.....	18
4.3.1	Aquatic bioaccumulation.....	18
4.3.2	Terrestrial bioaccumulation.....	18
4.3.3	Summary and discussion of bioaccumulation .....	18
4.4	Secondary poisoning.....	18
5	HUMAN HEALTH HAZARD ASSESSMENT.....	19
5.1	Toxicokinetics (absorption, metabolism, distribution and elimination) .....	19
5.1.1	Absorption.....	19
5.1.2	Distribution.....	20
5.1.3	Metabolism.....	20
5.1.4	Excretion .....	20
5.1.5	Summary and conclusion .....	20
5.2	Acute toxicity .....	21
5.2.1	Acute oral toxicity.....	21
5.2.2	Acute inhalation toxicity .....	21
5.2.3	Acute dermal toxicity .....	23
5.2.4	Acute toxicity: other routes .....	23
5.2.5	Summary and discussion of acute toxicity .....	23
5.3	Irritation .....	24
5.3.1	Skin Irritation .....	24
5.3.2	Eye irritation.....	25
5.3.3	Respiratory tract .....	25
5.3.4	Summary and discussion of irritation.....	25

5.4	Corrosivity.....	25
5.5	Sensitisation.....	25
5.5.1	Skin .....	25
5.5.2	Respiratory system .....	26
5.5.3	Summary and discussion of sensitisation .....	26
5.6	Repeated dose toxicity.....	26
5.6.1	Repeated dose toxicity: oral .....	26
5.6.2	Repeated dose toxicity: inhalation.....	28
5.6.3	Repeated dose toxicity: dermal .....	28
5.6.4	Other relevant information .....	29
5.6.5	Summary and discussion of repeated dose toxicity:.....	29
5.7	Mutagenicity.....	30
5.7.1	In vitro data .....	30
5.7.2	In vivo data.....	30
5.7.3	Human data .....	30
5.7.4	Other relevant information .....	30
5.7.5	Summary and discussion of mutagenicity .....	31
5.7.6	Conclusion.....	31
5.8	Carcinogenicity.....	31
5.8.1	Carcinogenicity: oral .....	31
5.8.2	Carcinogenicity: inhalation .....	31
5.8.3	Carcinogenicity: dermal .....	31
5.8.4	Carcinogenicity: human data .....	31
5.8.5	Other relevant information .....	31
5.8.6	Summary and discussion of carcinogenicity .....	32
5.9	Toxicity for reproduction.....	32
5.9.1	Effects on fertility.....	32
5.9.2	Developmental toxicity .....	33
5.9.3	Human data .....	36
5.9.4	Other relevant information .....	36
5.9.5	Summary and discussion of reproductive toxicity.....	36
5.10	Other effects .....	38
5.10.1	P-glycoprotein expression and increased susceptibility to abamectin.....	38
5.11	Derivation of DNEL(s) or other quantitative or qualitative measure for dose response.....	42
6	HUMAN HEALTH HAZARD ASSESSMENT OF PHYSICO-CHEMICAL PROPERTIES .....	43
6.1	Explosivity.....	43
6.2	Flammability.....	43
6.3	Oxidising potential .....	43
7	ENVIRONMENTAL HAZARD ASSESSMENT .....	44
7.1	Aquatic compartment (including sediment).....	44
7.1.1	Toxicity test results .....	44
7.1.2	Calculation of Predicted No Effect Concentration (PNEC) .....	46
7.2	Terrestrial compartment.....	46
7.2.1	Toxicity test results .....	46
7.2.2	Calculation of Predicted No Effect Concentration (PNEC_soil).....	46

7.3 Atmospheric compartment.....	47
7.4 Microbiological activity in sewage treatment systems .....	47
7.4.1 Toxicity to aquatic micro-organisms.....	47
7.4.2 PNEC for sewage treatment plant .....	47
7.5 Calculation of Predicted No Effect Concentration for secondary poisoning (PNEC_oral) .....	47
7.6 Conclusion on the environmental classification and labelling.....	47

## TABLES

Table 1.3-1: Summary of physico- chemical properties .....	9
Table 4.1-1 Hydrolysis of abamectin.....	13
Table 4.1-2 Photolysis of avermectin B <sub>1a</sub> in water .....	13
Table 4.1-3 Photolysis of avermectin B <sub>1a</sub> in soil .....	14
Table 4.1-4 Ready biodegradability of abamectin .....	14
Table 4.1-5 Degradation of abamectin in <i>aerobic</i> water/sediment systems.....	15
Table 4.1-6 Degradation of abamectin in <i>anaerobic</i> water/sediment systems.....	15
Table 4.1-7 Overview of DT <sub>50</sub> -values from aerobic laboratory degradation studies with avermectin B <sub>1a</sub> .....	16
Table 4.2-1 Adsorption of avermectin B <sub>1a</sub> onto soils.....	17
Table 4.3-1 Measurements of aquatic bioconcentration .....	18
Table 5.1-1 summarises toxicokinetics of abamectin in rats and humans. ....	20
Table 5.2-1 Key acute toxicity (LD50/LC50) studies reported for abamectin .....	23
Table 5.2-2 Summary of acute neurotoxicity study .....	24
Table 5.3-1 Skin Irritation .....	24
Table 5.3-2 Eye irritation.....	25
Table 5.5-1 Sensitisation .....	25
Table 5.6-1 Summary of neurotoxicity studies.....	26
Table 5.6-2 Summary of repeated dose toxicity studies .....	27
Table 5.6-3 Summary of repeated dose inhalation toxicity study.....	28
Table 5.7-1 Genotoxicity studies: In vitro .....	30
Table 5.7-2 Genotoxicity studies: In vivo.....	30
Table 5.8-1 Carcinogenicity study.....	31
Table 5.9-1 Fertility study .....	33
Table 5.9-2 Developmental effects in the rat study. ....	34
Table 5.9-3 Developmental effects in the rabbit study. ....	35
Table 5.9-4 Summary of teratogenicity studies .....	35
Table 7.1-1 Acute and chronic toxicity of abamectin and avermectin B <sub>1a</sub> to fish .....	44
Table 7.1-2 Acute and chronic toxicity of abamectin and avermectin B <sub>1a</sub> to invertebrates .....	44
Table 7.1-3 Chronic toxicity to sediment organisms .....	45
Table 7.1-4 Acute and chronic toxicity to saltwater organisms.....	46

## **PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING**

**Substance Name: Abamectin**

**EC Number:**

CAS number: 71751-41-2

Registration number (s): CIPAC 495 (collaborative international pesticides analytical council code number)

Purity:           Min. 90 % w/w abamectin (sum of avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>)  
                      Min. 83 % w/w avermectin B<sub>1a</sub>  
                      Max. 8 % w/w avermectin B<sub>1b</sub>

Impurities:       Based on the available environmental and (eco)toxicological information, there are no relevant impurities

Remark:

The present proposal for harmonized Classification and Labelling applies to the technical active substance abamectin as proposed for inclusion in Annex I of Council Directive 91/414/EEC and Annex I of Directive 98/8/EC.

Confidential information on the content and identity of isomers, impurities and additives is available in Volume 4, Annex C of the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands).

### **Proposed classification based on Directive 67/548/EEC:**

Phys/Chem hazards:

-

Health hazards:

Repr. Cat.3; R63

T+; R26/28

T ; R48/23/25

Environment:

N; R50/53

### **Proposed classification based on Regulation EC 1272/2008:**

Phys/Chem hazards:

-

Health hazards:

Repr. 2                H361d

Acute Tox. 2        H300

Acute Tox. 1        H330

STOT-RE 1         H372

Environment:

Aquatic Acute 1     H400

Aquatic Chronic 1   H410

**Proposed labelling:**

Directive 67/548/EEC:

Symbol            : T+, N

Risk phrases     : R26/28-R48/23/25-R63-R50/53

Safety phrases   : S28-S36/37-S45-S60-S61

Regulation EC 1272/2008:

Signal word: Danger

Symbol: GHS06, GHS09

Hazard statement codes: H300, H330, H361d, H372, H400, H410

As precautionary statements are not included in Annex VI of Regulation EC 1272/2008, no proposal is made.

**Proposed specific concentration limits (if any):**

M-factor for 67/548 EEC and EC 1272/2008:

The M-factor is 10,000. This value is based on two LC50 values of 0.020 µg/l and 0.022 µg/l obtained for the marine crustacean *Mysidopsis bahia* in a 96-h flow-through study.

## JUSTIFICATION

### 1 IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES

#### 1.1 Name and other identifiers of the substance

Chemical Name: Abamectin

EC Name:

CAS Number: 71751-41-2

IUPAC Name: none

Abamectin is the ISO common name for a mixture of 80 % avermectin B1a and 20 % avermectin B1b. The use of the word “mixture” in the ISO description is not in line with REACH and CLP terminology. Following the terminology of REACH and CLP Abamectine is a substance containing 80 % avermectin B1a and 20 % avermectin B1b. The material considered in this report is not an exact 80/20 ratio of these two constituents and therefore strictly speaking this ISO name can not be used. The registrant is currently trying to get the ratio amended to better describe the considered material. When information is available confirming the change in ISO name this proposal will be adapted accordingly.

According to the REACH guidance document on substance identification the substance is a mono-constituent substance with avermectin B1a (CAS Number 65195-55-3) as its main constituent (purity > 83%) and with avermectin B1b as an impurity. However, part 1.1.1.4 of Annex VI of EC 1272/2008 (CLP) states that whenever possible plant protection products and biocides are designated by their ISO names. As abamectin is used as both a plant protection product and as a biocide in this proposal preference is given to the use of the ISO name abamectin as the International Chemical Identifier for inclusion in Annex VI of EC 1272/2008.

#### 1.2 Composition of the substance

The present proposal for harmonized Classification and Labelling applies to the technical active substance abamectin as proposed for inclusion in Annex I of Council Directive 91/414/EEC and Annex I of Directive 98/8/EC.

Confidential information on the content and identity of isomers, impurities and additives is available in Volume 4, Annex C of the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands).

Purity and impurities



Purity: Min. 90 % w/w abamectin (sum of avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>)  
 Min. 83 % w/w avermectin B<sub>1a</sub>  
 Max. 8 % w/w avermectin B<sub>1b</sub>

Main constituent

Chemical Name: avermectin B<sub>1a</sub>

EC Number: 265-610-3


CAS Number: 65195-55-3

IUPAC Name: (10*E*,14*E*,16*E*,22*Z*)-(1*R*,4*S*,5'*S*,6*S*,6'*R*,8*R*,12*S*,13*S*,20*R*,21*R*,24*S*)-6'-[(*S*)-*sec*-butyl]-21,24-dihydroxy-5',11,13,22-tetramethyl-2-oxo-3,7,19-trioxatetracyclo[15.6.1.1<sup>4,8</sup>.0<sup>20,24</sup>]pentacosa-10,14,16,22-tetraene-6-spiro-2'-(5',6'-dihydro-2'*H*-pyran)-12-yl 2,6-dideoxy-4-*O*-(2,6-dideoxy-3-*O*-methyl- $\alpha$ -L-arabino-hexopyranosyl)-3-*O*-methyl- $\alpha$ -L-arabino-hexopyranoside

Molecular Formula:

C<sub>48</sub>H<sub>72</sub>O<sub>14</sub>

Structural Formula:



Molecular Weight: 873.1

Typical concentration (% w/w):  $\geq$  83%

Concentration range (% w/w): confidential

Impurity

Chemical Name:

avermectin B<sub>1b</sub>:

EC Number: 265-611-9

CAS Number: 65195-56-4


IUPAC Name: (10*E*,14*E*,16*E*,22*Z*)-(1*R*,4*S*,5'*S*,6*S*,6'*R*,8*R*,12*S*,13*S*,20*R*,21*R*,24*S*)-21,24-dihydroxy-6'-isopropyl-5',11,13,22-tetramethyl-2-oxo-3,7,19-trioxatetracyclo[15.6.1.1<sup>4,8</sup>.0<sup>20,24</sup>]pentacosa-10,14,16,22-tetraene-6-spiro-2'

(5',6'-dihydro-2'*H*-pyran)-12-yl 2,6-dideoxy-4-*O*-(2,6-dideoxy-3-*O*-methyl- $\alpha$ -L-arabino-hexopyranosyl)-3-*O*-methyl- $\alpha$ -L-arabino-hexopyranoside

Molecular Formula:

$C_{47}H_{70}O_{14}$

Structural Formula:



Molecular Weight: 859.1

Typical concentration (% w/w):  $\leq 8\%$

Concentration range (% w/w): confidential

Other impurities:

The natural fermentation process for the production of abamectin produces several impurities, which are structurally similar to avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>. Because of their low concentration level and their expected similar (eco)toxicity to avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>, these impurities are considered not (eco)toxicologically relevant in the material (see DAR October 2005 + addendum February 2008, RMS The Netherlands and the CAR; July 2008; RMS The Netherlands).

Test material:

The active substance abamectin, produced in a natural fermentation process, contains both avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>. All studies, unless otherwise stated, were carried out using abamectin which varied in purity between 88.3 and 96.7% (sum of avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub>). Where information on the ratio between avermectin B<sub>1a</sub> and avermectin B<sub>1b</sub> was available for batches used in the toxicological studies, these were above 80% for avermectin B<sub>1a</sub> and below 20% for avermectin B<sub>1b</sub>. The variation in purity and ratio is not expected to substantially affect the toxicity and the classification and labelling. Studies not carried out using abamectin were mostly conducted with the major component avermectin B<sub>1a</sub>.

### 1.3 Physico-chemical properties

**Table 1.3-1: Summary of physico- chemical properties**

REACH ref Annex, §	Property	IUCLID section	Value
VII, 7.1	Physical state at 20°C and 101.3 KPa	3.1	Powder at 25 °C (96.7% w/w)
VII, 7.2	Melting/freezing point	3.2	161.8 °C – 169.4 °C (96.7% w/w) with thermal decomposition during melting (at 162 °C)
VII, 7.3	Boiling point	3.3	Not determined, due to thermal decomposition during melting of abamectin
VII, 7.4	Relative density	3.4 density	1.18 at 22 °C (96.7% w/w)
VII, 7.5	Vapour pressure	3.6	< 3.7 x 10 <sup>-6</sup> Pa at 25 °C (96.7% w/w) using the gas saturation method
VII, 7.6	Surface tension	3.10	52.4 mN/m at 90% of the saturation concentration at 20 °C (purity 96.7% w/w)
VII, 7.7	Water solubility	3.8	Water solubility at 25 °C (purity not stated) using the shake flask method pH 7.57: 1.21 mg/L (in water)
VII, 7.8	Partition coefficient n-octanol/water (log value)	3.7 partition coefficient	log Kow = 4.4 at pH 7.2 at 20 °C (water), (purity 96.7% w/w) using shake flask method
VII, 7.9	Flash point	3.11	No data
VII, 7.10	Flammability	3.13	Abamectin is considered to be not highly flammable
VII, 7.11	Explosive properties	3.14	No explosive properties
VII, 7.12	Self-ignition temperature		No self-ignition was observed before the melting point
VII, 7.13	Oxidising properties	3.15	No oxidising properties
VII, 7.14	Granulometry	3.5	No data
XI, 7.15	Stability in organic solvents and identity of relevant degradation products	3.17	No data
XI, 7.16	Dissociation constant	3.21	No dissociation in the pH-range from 1 to 12
XI, 7.17	Viscosity	3.22	Not data, Abamectin is a solid, not a liquid
	Auto flammability	3.12	No data
	Reactivity towards container material	3.18	Abamectin is packed in a conical bucket inside another conical bucket (inside the buckets are two polyethylene bags which contain the material).

			The registrant indicates that there is no record of any reaction to the container material
	Thermal stability	3.19	Thermal decomposition during melting (above 162 °C, 96.7% w/w) Identification of breakdown products was not performed. The registrant indicates that combustion products are likely to be oxides of carbon and water. The registrant indicates that dangerous products are unlikely to be formed.

The above data are obtained from the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands) and the Competent Authority Report (CAR; July 2008; RMS The Netherlands) on the inclusion of abamectin in Annex I to Directive 98/8/EC concerning the placing of biocidal products on the market.

## 2 MANUFACTURE AND USES

Abamectin is used as an insecticide and acaricide for the control of motile stages of mites, leaf mines, suckers, Colorado beetles, etc. on ornamentals, cotton, citrus fruit, pome fruit, nut crops, vegetables, potatoes and other crops. Also used for the control of fire ants.

The assessment of the biocidal activity of the active substance demonstrates that it has a sufficient level of efficacy against some of the target organisms (pharao ants and cockroaches)

## 3 CLASSIFICATION AND LABELLING

### 3.1 Classification in Annex I of Directive 67/548/EEC

*Current classification:* None

### 3.2 Self classification(s)

The registrant has proposed the following classification and labelling of the active substance abamectin.

Proposal of the registrant according to Directive 67/548/EEC for abamectin:

Hazard symbol:	T+	Very toxic
	N	Dangerous for the environment
Risk phrases	R26	Very toxic by inhalation
	R28	Very toxic if swallowed
	R50/53	Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment
Safety phrases	S28	After contact with skin, wash immediately with plenty of ... (to be specified by the manufacturer)
	S36/37	Wear suitable protective clothing and gloves
	S45	In case of accident or if you feel unwell seek medical advice immediately (show the label where possible)
	S60	This material and its container must be disposed of as hazardous waste
	S61	Avoid release to the environment. Refer to special instructions/safety data sheet

Proposal of the registrant according to Regulation EC 1272/2008: No proposal

## 4 ENVIRONMENTAL FATE PROPERTIES

The environmental fate properties assessment for abamectin is based on the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands) and the Competent Authority Report (CAR; July 2008; RMS The Netherlands) on the inclusion of abamectin in Annex I to Directive 98/8/EC concerning the placing biocidal products on the market.

All tables in the present assessment are copied from the DAR or CAR. The tables are renumbered in accordance with the paragraph numbers in chapter 4.

### 4.1 Degradation

#### 4.1.1 Stability

##### Hydrolysis

Both <sup>14</sup>C- and <sup>3</sup>H-avermectin B<sub>1a</sub> are hydrolytically stable at environmentally relevant pH (4 - 7) and temperature (25 °C). Under basic conditions (pH 9), DT<sub>50,hydrolysis</sub> of avermectin B<sub>1a</sub> was 213, 9.9 and 4.9 days at 25, 50 and 60 °C, respectively, and the calculated DT<sub>50,hydrolysis</sub> at 20 °C is 380 days.

Table 4.1-1 Hydrolysis of abamectin

Guideline/ Test method	Substance	pH	Temperature [°C]	Initial test substance concentration, C <sub>0</sub> [mg/L]	Reaction rate constant, K <sub>h</sub> [1/d] <sup>a</sup>	Half-life, DT <sub>50,hydro</sub> [d]	Reference
OECD 111; EPA N 161- 1; BBA 55, I and II	<sup>14</sup> C-avermectin B <sub>1a</sub>	4	50	0.11 (with 20% acetonitrile)	3.25 x 10 <sup>-3</sup>	no hydrolysis	Ellgehausen, 2001
		5	50	0.11 (with 20% acetonitrile)		no hydrolysis	
		7	50	0.11 (with 20% acetonitrile)		no hydrolysis	
		9	25	0.11 (with 20% acetonitrile)		213	
		9	50	0.11 (with 20% acetonitrile)		9.9	
		9	60	0.11 (with 20% acetonitrile)		4.9	

a: calculated as ln2/DT<sub>50</sub>

##### Photolysis in water

The data on aqueous photodegradation of avermectin B<sub>1a</sub> are summarised in Table 4.1-2.

Table 4.1-2 Photolysis of avermectin B<sub>1a</sub> in water

Guideline/ Test method	Substance	Initial test substance concentration, C <sub>0</sub> [µg/L]	Total recovery of test substance [% of added radioactivity]	Photolysis rate constant, K <sub>p</sub> <sup>c</sup> [1/d] <sup>a</sup>	Reaction quantum yield [Φ <sup>°E</sup> ]	Half-life, DT <sub>50,photo</sub> [d]	Reference
EPA 161-2	<sup>14</sup> C-avermectin B <sub>1a</sub>	100 (in 1% acetonitrile)	91.8	0.35		2	Adam, 2001b
EPA 188	<sup>3</sup> H-avermectin B <sub>1a</sub>	3 (in 1% acetonitrile)	93.3	0.53	0.0287-0.0347	1.3	Halley et al., 1991

a: calculated as ln2/DT<sub>50</sub>

The DT<sub>50,photolysis</sub> for avermectin B<sub>1a</sub> of 2 days obtained in the first study (Adam, 2001b) is equivalent to 1.5 sunlight days at 30 - 50 °N.

In a second study by Halley et al., (1991), the DT<sub>50,photolysis</sub> of 1.3 days from this study was obtained after irradiation with natural sunlight at Three Bridges, NJ, USA from September 26 to October 2,

1990. Samples received about 8 hours sunlight each day. Quantum yield was determined to be 0.0347, 0.0316 and 0.0287 at 40 °N in summer, fall and winter, respectively.

### Photolysis in soil

The data on photodegradation of avermectin B<sub>1a</sub> in soil are summarised in Table 4.1-3.

Table 4.1-3 Photolysis of avermectin B<sub>1a</sub> in soil

Guideline/ Test method	Substance	Initial tests substance application rate, C <sub>0</sub> [kg/ha]	Total recovery of test substance [% of added radioactivity]	Photolysis rate constant, K <sub>p</sub> <sup>c</sup> [1/d] <sup>a</sup>	Reaction quantum yield [Φ°E]	Half-life, DT <sub>50,photo</sub> [d]	Reference
EPA 161-3	<sup>14</sup> C-avermectin B <sub>1a</sub>	0.09	100	0.05		13	Phaff, 2001

a: calculated by RMS as  $\ln 2/DT_{50}$

The DT<sub>50,photolysis</sub> of 13 days is equivalent to 22 days at 30 – 50 °N. Mineralisation and bound residues were 7.6 and 25.9 %, respectively, after 28 days.

### Photo-oxidative degradation in air

The atmospheric half-life time of abamectin is estimated according to Atkinson as < 1 hour (Stamm, 1998).

## **4.1.2 Biodegradation**

### **4.1.2.1 Biodegradation estimation**

No data available.

### **4.1.2.2 Screening test**

#### *Readily biodegradability*

Abamectin at concentration (100 mg/L) in the study of Dietschy (1999) was far above the solubility in water of 1.21 mg/L but within the concentration range recommended by the test guideline. In the absence of other data, abamectin is considered as *not readily biodegradable*.

Table 4.1-4 Ready biodegradability of abamectin

Guideline/ Test method	Test type	Test parameter	Inoculum			Additional substrate	Test substance concentration	Degradation		Remarks	Reference
			Type	Concen- tration	Adaptation			Incubation period [d]	Degree [%]		
OECD 301F	manometric respirometry	oxygen demand	STP	26 mg/L		synthetic	100 mg/L	28	3		Dietschy (1999)

### **4.1.2.3 Simulation tests**

#### Biodegradation in water/sediment systems



*Aerobic water/sediment system*

In the study of Buckel (2002), a sandy loam system (River Rhine) and a silty clay loam system (pond) were treated with  $^{14}\text{C}$ -avermectin B<sub>1a</sub> and incubated under aerobic or anaerobic conditions at 20 °C in the dark. The results of the aerobic incubation are summarised in Table 4.1-5.

Table 4.1-5 Degradation of abamectin in *aerobic* water/sediment systems

Guideline/ Test method	Substance	System name	Sediment type	Condition	T [°C]	pH <sub>water</sub>	OM [%]	Duration [d]	DT <sub>50</sub> water <sup>1</sup> [d]	DT <sub>50</sub> sediment [d]	DT <sub>50</sub> system [d]	Reference
OECD draft 2000; BBA IV, 5-1	$^{14}\text{C}$ - avermectin B <sub>1a</sub>	River Rhine	sandy loam	aerobic	20	7.9- 8.4	2.5	100	1.8	87	87	Buckel, 2002
OECD draft 2000; BBA IV, 5-1	$^{14}\text{C}$ - avermectin B <sub>1a</sub>	Rothenfluh pond	silty clay loam	aerobic	20	7.7- 8.4	7.7	100	2.9	111	91	Buckel, 2002

1: DT<sub>50,water</sub> determined by sorption, value represents dissipation

The decline of concentrations in the water phase was mainly determined by a rapid initial sorption, and the DT<sub>50,water</sub> thus represents dissipation rather than degradation.

The maximum level of avermectin B<sub>1a</sub> found in sediment was 78.1% of added radioactivity (pond) and 82.8% of added radioactivity (river) after 14 days. At the end of the study after 100 days, levels of avermectin B<sub>1a</sub> had declined to 44.3 and 45.3% of added radioactivity for river and pond, respectively.

Bound residues increased to 20.4% of added radioactivity (river) and 23.2% of added radioactivity (pond) at the end of the study after 100 days, mineralisation was low with a maximum of 3.0 and 3.2% of added radioactivity after 100 days in the river and pond system, respectively.

*Anaerobic water/sediment system*

In the *anaerobic* systems, dissipation from the water phase was fast (DT<sub>50,water</sub> of 5.6-7.2 days, see tabel 4.1-6), but degradation in the total system was much slower with < 50% degradation at the end of the study after 100 days (see Table 4.1-6). DT<sub>50,sediment</sub>-values could not be estimated because there were too few data points with decline.

Table 4.1-6 Degradation of abamectin in *anaerobic* water/sediment systems

Guideline/ Test method	Substance	System name	Sediment type	Condition	T [°C]	pH <sub>water</sub>	OM [%]	Duration [d]	DT <sub>50</sub> water <sup>1</sup> [d]	DT <sub>50</sub> sediment [d]	DT <sub>50</sub> system <sup>2</sup> [d]	Reference
OECD draft 2000; BBA IV, 5-1	$^{14}\text{C}$ - avermectin B <sub>1a</sub>	River Rhine	sandy loam	anaerobic	20	8.4- 9.2	2.5	100	7.2		230	Buckel, 2002
OECD draft 2000; BBA IV, 5-1	$^{14}\text{C}$ - avermectin B <sub>1a</sub>	Rothenfluh pond	silty clay loam	anaerobic	20	7.8- 9.8	7.7	100	5.6		312	Buckel, 2002

1: DT<sub>50,water</sub> determined by sorption, value represents dissipation

2: extrapolated value

Biodegradation in soil

*Aerobic biodegradation*

The rate of degradation of avermectin B<sub>1a</sub> under aerobic conditions was assessed in four laboratory experiments in eight different soil types. The experiments are summarised in Table 4.1-7. Results from the study of Ku and Jacob (1983a) indicate that there is no clear relation between dose and degradation rate. This study was performed at ambient temperature, which is supposed to be 20 °C. Moisture content may influence the degradation rate: at 30 °C, the DT<sub>50</sub> under dry conditions (pF 4) is 24.4 days, which is higher than the value found at field capacity (16.6 days).

Where multiple experiments were performed with a single soil type under the same conditions (temperature, moisture), the geometric mean of the DT<sub>50</sub>-values is calculated.

Table 4.1-7 Overview of DT<sub>50</sub>-values from aerobic laboratory degradation studies with avermectin B<sub>1a</sub>.

Guideline/ Test method	Label	Soil type	Dose [mg/kg]	T [°C]	OM [%]	pH	pF	DT <sub>50</sub> [d]	DT <sub>50</sub> , 20 °C [d]	Reference
BBA IV, 4-1; draft OECD	<sup>14</sup> C	loam	0.22	20	3.2	7.3	2	18.8	18.8	Nicollier, 2001
BBA IV, 4-1; draft OECD	<sup>14</sup> C	silt loam	0.1	20	4	7.2	2.5	23.3	23.3	Adam, 2001a
	<sup>14</sup> C	silt loam	0.1	10	4	7.2	2.5	50.6 <sup>1</sup>		
	<sup>14</sup> C	silt loam	0.1	30	4	7.2	2.5	16.6		
	<sup>14</sup> C	silt loam	0.1	30	4	7.2	4	24.4		
BBA IV, 4-1; draft OECD	<sup>14</sup> C	loamy sand	0.125	20	2.4	7.41	2.5	23.6	23.6	Phaff, 2003
	<sup>14</sup> C	sandy clay loam	0.125	20	4.3	5.81	2.5	11.2	11.2	
	<sup>14</sup> C	silty clay loam	0.125	20	2.4	7.92	3.5	49.6	49.6	
not specified	<sup>3</sup> H	sandy loam	0.1	ambient	1.1	6.8	2.5	26.9	geometric mean for sandy loam: 28.3	Ku & Jacob, 1983a
	<sup>3</sup> H	sandy loam	1	ambient	1.1	6.8	2.5	22.3		
	<sup>3</sup> H	sandy loam	50	ambient	1.1	6.8	2.5	42.6		
	<sup>14</sup> C	sandy loam	1	ambient	1.1	6.8	2.5	15.1		
	<sup>14</sup> C	sandy loam	1	ambient	1.1	6.8	2.5	47.0		
	<sup>3</sup> H	sand	1	ambient	0.6	8	2.5	65.7		
	<sup>3</sup> H	clay	0.1	ambient	1.3	6.8	2.5	34.9	geometric for clay: 39.6	
	<sup>3</sup> H	clay	1	ambient	1.3	6.8	2.5	44.9		

1: actual temperature 8.6 °C, value at 10 °C estimated using the Arrhenius equation

The overall geometric mean DT<sub>50</sub> of avermectin B<sub>1a</sub> at 20 °C is 28.4 days (range 11.2 – 65.7 days; n = 8; r<sup>2</sup> 0.9471 - 0.9970).

The highest formation of bound residues was 39.1% of added radioactivity after incubation for 91 days at 20 °C and further increased to 44.1% of added radioactivity after 196 days (Phaff, 2003). Highest mineralisation accounted for 12.4% of added radioactivity after 91 days (Phaff, 2003) and reached 27.6% of added radioactivity in another study at the end of a 365-days incubation period (Nicollier, 2001).

Based on degradation rates at 8.6, 20 and 30 °C (Adam, 2001a), the DT<sub>50</sub> of avermectin B<sub>1a</sub> at 10 °C is estimated as 50.6 days.

#### 4.1.3 Summary and discussion of persistence

##### Biodegradation in water

Avermectin was found to be not readily biodegradable in a ready biodegradability study.

In natural aerobic water/sediments systems, the dissipation of avermectin from the water phase was dominated by sorption with a DT<sub>50,water</sub> of 2.4 days. The average DT<sub>50,system</sub> was 89 days whereas the DT<sub>50,sediment</sub> was 99 days.

In natural anaerobic water/sediment systems, dissipation of abamectin was fast with  $DT_{50,water}$  of 6.4 days. In contrast,  $DT_{50,sediment}$  could not be determined due to limited degradation. The  $DT_{50,system}$  was on average 271 days.

### Biodegradation in soil

The geometric mean  $DT_{50}$  of avermectin  $B_{1a}$  in soil at 20 °C is 28.4 days. The highest formation of bound residues was 39.1% of added radioactivity after incubation for 91 days at 20 °C. Highest mineralisation accounted for 12.4% of added radioactivity after 91 days.

## 4.2 Environmental distribution

### 4.2.1 Adsorption/desorption

Batch equilibrium experiments have been performed with avermectin  $B_{1a}$  in eight different soils. One of the soils was a sand with 0.1% OM (0.06% OC), which is considered not relevant for risk assessment. Accepted  $K_{OC}$ -values are summarised in Table 4.2-1. The average  $K_{OC}$  is 5638 L/kg (range 1495 – 7893; n = 7). Sorption of avermectin  $B_{1a}$  is related to OC-content, linear regression of  $K_F$  versus % OC gives a regression coefficient  $r^2$  of 0.919. Abamectin can be considered as immobile in soil.

Table 4.2-1 Adsorption of avermectin  $B_{1a}$  onto soils

Guideline / Test method	Adsorbed a.s. [%]	$K_a^1$ [L/kg]	$K_{aOC}^2$ [L/kg]	$K_d^3$ [L/kg]	$K_{dOC}^4$ [L/kg]	$K_a / K_d^5$	Degradation products		Remarks	Reference
							Name	[%] of a.s.		
<b><sup>14</sup>C-avermectin <math>B_{1a}</math></b>										
OECD 106		87.2	5701				test substance was stable during mass balance experiment	loamy sand loamy sand sandy loam loam silt loam	Morgenroth, 2001	
OECD 106		77.3	7893							
OECD 106		76.8	6004							
OECD 106		178	6875							
OECD 106		334	6682							
<b><sup>3</sup>H-avermectin <math>B_{1a}</math></b> not spec.		18.2 134	1495 4814				test substance was stable during mass balance experiment	silt loam clay loam	Gruber & Wislocki, 1988	
		average	5638							
1: $K_a$ = Adsorption coefficient						4: $K_{dOC}$ = Desorption coefficient based on organic carbon content				
2: $K_{aOC}$ = Adsorption coefficient based on organic carbon content						5: $K_a / K_d$ = Adsorption / Desorption distribution coefficient				
3: $K_d$ = Desorption coefficient										

### 4.2.2 Volatilisation

Abamectin has a vapour pressure of  $< 3.7 \times 10^{-6}$  Pa at 25 °C.

### 4.2.3 Distribution modelling

Not relevant for this dossier

### 4.3 Bioaccumulation

#### 4.3.1 Aquatic bioaccumulation

##### 4.3.1.1 Bioaccumulation estimation

Abamectin has a log Kow of 4.4 at pH 7.2 at 20 °C.

##### 4.3.1.2 Measured bioaccumulation data

Results of a bioconcentration study with abamectin are summarised in the table below. The value of 69 L/kg ww was calculated from the Total Radioactive Residues (TRR) in whole fish (plateau) and the average TRR in water. Based on the fitted uptake and elimination rate constants, the BCF is 52 L/kg ww.

Table 4.3-1 Measurements of aquatic bioconcentration

Guideline / Test method	Exposure	Log P <sub>ow</sub>	Initial concentr. [µg/L]	Steady-state BCF [L/kg ww]	Uptake rate constant [mL/g.d]	Depuration rate constant [1/d]	Depuration time (DT <sub>50</sub> ) [d]	Metabolites	Remarks	Reference
ASTM 1978	flow-through	4.4	0.1	69 <sup>1</sup> 52 <sup>2</sup>	11	0.21	3.3	N.D. <sup>3</sup>	whole fish; based on TRR	Forbis & Franklin, 1983

1: based on plateau TRR in whole fish and average TRR in water

2: estimated from uptake and elimination rate constants

3: ND, Not determined

#### 4.3.2 Terrestrial bioaccumulation

#### 4.3.3 Summary and discussion of bioaccumulation

Abamectin has a log Kow of 4.4. However, a BCF of 52 L/kg ww (based on the total radioactive residue) and 69 L/kg (based on whole fish) was obtained in a bioaccumulation study. Based on the results of the bioaccumulation study, abamectin does not significantly bioaccumulate.

### 4.4 Secondary poisoning

Not relevant for this dossier.

## 5 HUMAN HEALTH HAZARD ASSESSMENT

The human health hazard assessment for abamectin is based on the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands), the EFSA conclusion (EFSA Scientific Report (2008) 147, 1-106) and the Competent Authority Report (CAR; January 2009; RMS The Netherlands) on the inclusion of abamectin in Annex I to Directive 98/8/EC concerning the placing biocidal products on the market.

It should be noted that in the present human health hazard assessment most data on studies in CF-1 mice are excluded. Many studies with abamectin were performed with the CF-1 mouse, which is very sensitive to the observed developmental effects. Based on a recent extensive overview of the literature, it was however concluded that the CF-1 mouse is not relevant for human risk assessment (see 5.10.1) because some CF-1 mice lack the p-glycoprotein which has a function in restricting the brain penetration of avermectins including abamectin. Absence of p-glycoprotein is not known to occur in humans. The results of studies with the CF-1 mouse are therefore also not relevant for classification and labelling.

All tables in the present human health hazard assessment are copied from the DAR or CAR. The tables are renumbered in accordance with the paragraph numbers in chapter 5.

### 5.1 Toxicokinetics (absorption, metabolism, distribution and elimination)

#### 5.1.1 Absorption

##### 5.1.1.1 Oral Absorption

Avermectin B1a, administered in sesame oil or in polyethylene glycol, is absorbed from the gastrointestinal tract of the rat and is distributed throughout all major tissues and organs sampled. Maximum concentrations in blood are achieved within 4-8 h after administration.

The comparison of urinary excretion after oral or i.v. administration indicate almost complete oral absorption, with a calculated bioavailability of 0.86.

##### 5.1.1.2 Inhalation Absorption

For the inhalation route no data are available. Absorption through inhalation is assumed to be 100%.

##### 5.1.1.3 Dermal Absorption

The extent of dermal penetration of avermectin B1a is minimal in the rhesus monkey, amounting to less than 1% of the applied dose. The low dermal absorption of <1% was confirmed by a recent *in vitro* dermal absorption study with human skin (see revised addendum to pesticides Draft Assessment Report, February 2008).

### 5.1.2 Distribution

Avermectin B1a and/or metabolites do not accumulate in liver, kidneys, muscle or fat on repeated administration of a low dose. Seven days after the last of 14 daily consecutive doses less than 1% of the total administered dose was present in tissues and organs. The highest residue levels were found in fat, with more than 10 times higher levels compared to other tissue residue levels. A comparative distribution and clearance study with avermectin B1b following single oral doses showed that the toxicokinetic profile was essentially the same as that of avermectin B1a.

### 5.1.3 Metabolism

The metabolite pattern in urine, faeces and bile is complex, and 11 metabolites were isolated. In faeces, avermectin B1a accounted for 24 to 45% of the dose, and the metabolite 3''-O-desmethyl-avermectin B1a [=3''DM] accounted for 19-27%. These major faecal components were not present in urine. In fat and muscle, avermectin B1a was the major component (92% and 72%, respectively), and metabolite [3''DM] accounted for 1.7% and 19% in the fat and muscle, respectively. The major reactions involved in the biotransformation of avermectin B1a in the rat are demethylation, hydroxylation, cleavage of the oleandrosyl ring and oxidation reactions.

In rat the 8,9-Z isomer of abamectin B1a is not formed.

### 5.1.4 Excretion

Avermectin B1a and/or metabolites is rapidly eliminated from the body, almost exclusively in the faeces (more than 92% of the dose within 7 days, urinary excretion accounting for 0.9-1.6% of the dose in males and 0.5-1.0% in females of low and high dose groups). Initially, the rate of excretion was slower in females as compared to males. The excretion via expired air accounted for only 0.01% of the dose within 48h after administration. Tissue half-lives were mostly within the range of  $1.2 \pm 0.3$  days, with the tissue half-lives of avermectin B1a being lower in males (12 to 17 h) compared to females (13 to 33 h). So, with the exception of dose-dependence for tissue residue levels and excretion by urine, the toxicokinetic profile is not influenced by sex, dose level or treatment regime.

### 5.1.5 Summary and conclusion

Table 5.1-1 summarises toxicokinetics of abamectin in rats and humans.

Absorption	Oral: Complete oral absorption with a calculated bioavailability of 0.86. For risk assessment of abamectin a value of 100% is assumed. Inhalation: No data are available. Absorption is assumed to be 100%. Dermal: less than 1% absorption.
Distribution	Widely distributed
Metabolism	Extensive metabolism (demethylation, hydroxylation, cleavage of oleandrosyl ring and oxidation reactions).
Excretion	Rapidly eliminated from body, almost exclusively in faeces.

## CONCLUSION

Abamectin is almost completely absorbed in the gastrointestinal tract of the rat (calculated oral bioavailability is 0.86) and distributed throughout tissues and organs. It is rapidly eliminated from the body, almost exclusively in the faeces, and does not accumulate in tissues/organs after repeated exposure. The major reactions involved in the biotransformation of abamectin in the rat are demethylation, hydroxylation, cleavage of the oleandrosyl ring and oxidation reactions.

Dermal penetration is very low, less than 1% is absorbed through the skin of monkeys. For the inhalation route 100% absorption is assumed.

## 5.2 Acute toxicity

### 5.2.1 Acute oral toxicity

#### *Animal data*

Abamectin is very toxic to the rat by oral administration in sesame oil (LD50 8.7-12.8 mg/kg bw). However, a subsequent study with an aqueous vehicle showed that abamectin was significantly less toxic orally with this vehicle. In the toxicokinetic studies performed with sesame oil or polyethylene glycol there are no indications for this observed difference in toxicity. Characteristic signs of abamectin toxicity after oral administration are tremors and ataxia. As abamectin is lipophilic the sesame oil is considered to be a more suitable vehicle than water, and classification of abamectin for acute toxicity will be based on the LD50 values from the studies in which sesame oil was used as vehicle.

In an acute neurotoxicity study in rats abamectin, administered by gavage induced clinical signs of neurotoxicity, i.e. reduced splay reflex, tiptoe gait and splayed gait. The NOAEL was 0.5 mg/kg bw, based on reduced splay reflex at 1.5 mg/kg bw. At 6 mg/kg bw, reduced splay reflex, tip toe and splayed gait and a transient reduction in motor activity was observed.

#### *Human data*

Available human data from suicide attempts show that typical clinical signs of abamectin toxicity in animal studies, like tremors and convulsions, do not occur in humans. No signs of poisoning were reported in a few cases after ingestion of low doses (up to 40 mg/kg bw). In other cases (estimated exposure 4.2 - 67 mg/kg bw), nausea, vomiting and diarrhoea or short lasting CNS depressions like dizziness, drowsiness and weakness were observed. Severe poisoning after suicidal ingestion of high amounts of an abamectin formulation (equivalent to 38.5 - 227.3 mg/kg bw abamectin) resulted in a comatose state within 3 hours after ingestion, shock, respiratory failure and even death as a result of multiple organ failure. The dose of abamectin ingested orally by a patient with lethal outcome in suicidal intention was 88.1 mg/kg. The maximum tolerated dose via the same route by another patient was 227.3 mg/kg.

### 5.2.2 Acute inhalation toxicity

Two acute inhalation toxicity studies with rats are available and included in this report.

#### **Characteristics**

Reference/notifier	: Ruddock, W. (2001a)	Exposure	: 4 h (nose only)
Type of study	: Acute inhalation toxicity study	Dose	: 0.21, 1.78 and 5.25 mg/L ( MMAD 4.2, 3.7 and 2.7 resp., GSD 3.8, 4.9 and 3.4 resp.)
Year of execution	: 2001	Vehicle	: -
Test substance	: Abamectin (purity 89.3% and 96.7%)	GLP statement	: yes
Route	: inhalation	Guideline	: In accordance with OECD 403
Species	: Rat (CrI:Han Wist)	Acceptability	: acceptable
Group size	: 5/sex/dose	LC50 rats	: < 0.21 mg/L

#### **Study design**

The study is in accordance with OECD 403.

**Results**

**Mortality:** There was 100% mortality in all dose groups. Death or moribund sacrifice occurred during exposure or within 2 h for the 5.25 mg/L animals, within 5 h for the 1.78 mg/L animals and by the day following exposure for the 0.21 mg/L animals.

**Symptoms of toxicity:** tremors, rigid tail and prostrate body position were observed in all animals, whereas signs observed in some animals included ataxia, cyanosis, subdued behaviour, piloerection, noisy respiration, coloured tears, staining of the eye and fur, squinting, wet fur, vocalisation on handling and tail flicking.

**Body weight:** Body weight analysis was not appropriate due to the early termination of the rats.

**Pathology:** dark, darkened or red areas in the lungs were observed in all dose groups, inflated lungs and firmness along the length of the tail was noted for all 5.25 mg/L animals.

**Acceptability**

The study is considered acceptable.

**Conclusions**

The acute 4-hour inhalatory LC<sub>50</sub> in rats is <0.21 mg/L.

**Characteristics**

Reference/notifier	: Noakes, J.P. (2003)	Exposure	: 4 h (nose only)
Type of study	: Acute inhalation toxicity study	Dose	: m + f: 0.051 mg/L (MMAD 2.11, 2.29; GSD 1.69, 1.83); f: 0.034 mg/L (MMAD 2.80, 2.57; GSD 1.73, 1.70)
Year of execution	: 2003	Vehicle	: -
Test substance	: Abamectin (purity 88.3%)	GLP statement	: yes
Route	: inhalatoir	Guideline	: In accordance with OECD 403
Species	: Rat (Alp:AP,SD)	Acceptability	: acceptable
Group size	: 5/sex (0.05 mg/L) and 5f (0.03 mg/L)	LC50 rats	: >0.051 mg/L (m); >0.034 mg/L and <0.051 mg/L (f)

**Study design**

The study is in accordance with OECD 403, with the following deviation: there are only 2 concentrations tested, and exposure to 0.03 mg/L was performed with 5 females only, pathology was not performed.

**Results**

**Mortality:** In the 0.05 mg/L group, one female was found dead and 2 females were killed on day 2 due to the severity of the clinical signs on day 2.

**Symptoms of toxicity:** reduced splay reflex, prostrate and tip toe gait, shaking, comatose, increased response to touch, reduced stability, decreased visual placing response, abnormal respiratory noise, increased breathing depth were observed in animals of the 0.051 mg/L group. In the 0.034 mg/L group, abnormal respiratory noise was observed staining of the oral and nasal cavities and eye discharge, wet fur, hunched posture, piloerection and chromodacryorrhea. Full recovery was apparent by day 4 for surviving females and for males by day 15.

**Body weight:** normal

**Pathology:** not performed



## Acceptability

The study is considered acceptable.

## Conclusions

The acute 4-hour inhalatory LC<sub>50</sub> in rats is >0.051 mg/L for males and between 0.051mg/L and 0.034 mg/L for females.

### 5.2.3 Acute dermal toxicity

Topical application of abamectin resulted in the rabbit in a 24hr LD50 value >2000 mg/kg and in the rat in a 24hr LD50 value >330 mg/kg (highest dose tested). The low order of toxicity by topical application indicates a low order of percutaneous penetration. This is supported by data in rhesus monkeys which demonstrate that < 1% of the applied dose is absorbed through the skin in to the systemic circulation.

Characteristic signs of abamectin toxicity, tremors and ataxia, occur in rats at 330 mg/kg bw about 3 days after administration and in rabbits at 2120 mg/kg bw within 6 days after administration.

### 5.2.4 Acute toxicity: other routes

No data

### 5.2.5 Summary and discussion of acute toxicity

Table 5.2-1 Key acute toxicity (LD50/LC50) studies reported for abamectin

Route	Method Guideline	Species Strain Sex no/group	Dose levels duration of exposure	Value LD50/LC50	Reference
Oral	Not fully in compliance with OECD 401 (but rated acceptable in 91/414/EEC DAR)	Rat/ CRCD 10 m/f	6.67, 10, 15, 22.5, 33.75 mg/kg bw in sesame oil single dose	M: 8.7 F: 12.8	Robertson 1981f
Oral	In compliance with OECD 401	Rat/- 5 m/f	20, 50, 100, 275, 500 mg/kg bw in 0.5% methylcellulose in water single dose	M: 232 F: 214	Glaza 2001
Dermal	Not fully in compliance with OECD 402 (but rated acceptable in 91/414/EEC DAR)	Rat/CD(SD)BR 5 m/f	330 mg/kg bw; 24h exposure	> 330	Gordon 1985a
Dermal	Not fully in compliance with OECD 402 (but rated acceptable in 91/414/EEC DAR)	Rabbit 5 m/f	2120 mg/kg bw; 24h exposure	> 2000	Gordon 1984a
Inhalation	In compliance with OECD 403	Rat/Crl:Han Wist	0.21, 1.78 and 5.25 mg/L 4 h (nose only)	<0.21	Ruddock, W. (2001)
Inhalation	Not fully in compliance with OECD 403 (but rated acceptable in 91/414/EEC DAR)	Rat/Alp:ApSD) 5 m/f	0.051 mg/L (m/f) 0.034 mg/L (f) 4 h (nose only)	M: >0.051 F: between 0.034 and 0.051	Noakes 2003

Table 5.2-2 Summary of acute neurotoxicity study

Species	Study type or duration; Dose levels	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Effect at LOAEL	Reference
Rat	Oral study of acute neurotoxicity, OECD 424. 0, 0.5, 1.5, 6 mg/kg bw by gavage	0.5	1.5	reduced splay reflex	Brammer, A. 2006a

### Classification proposals according to Directive 67/548/EEC

Based on the acute oral LD50 values (8.7-12.8 mg/kg bw) observed in the rat, abamectin needs to be classified as R28 “very toxic if swallowed”. Based on the acute inhalation LC50 value (<0.21 mg/l) abamectin needs to be classified as R26 “very toxic by inhalation”. Based on the available data the compound needs not to be classified when exposed via skin.

The limited number of human cases seem to indicate a somewhat lower acute oral toxicity of abamectin towards humans (lethality at 88 mg/kg bw/day) compared to rats. However, most patients were intensively treated. Also, it is unclear whether the vehicle has affected the human toxicity. The rat data are considered relevant for humans in a quantitative way.

### Classification proposals according to Regulation EC 1272/2008

According to CLP abamectin should be classified in acute hazard category 2 for oral exposure and in acute hazard category 1 for inhalation exposure, and labeled with signal word ‘Danger’ and hazard statements: H300 and H330.

Clinical signs of mild neurotoxicity were observed in an acute oral (gavage) neurotoxicity study at 1.5 and 6 mg/kg bw. It is noted that this is not much lower than the LD50s of 8.7-12.8 mg/kg bw (on which basis it is proposed to classify abamectin in acute hazard category 2 for oral exposure). In humans ingestion of doses up to 40 mg/kg bw induced no signs of poisoning.

Mild signs of neurotoxicity were observed in animals while in human poisoning cases with relative low doses no neurotoxic effects were reported. Since it is already proposed to classify abamectin for acute toxicity on the basis of the LD50 studies, no additional classification of abamectin for Specific Target Organ Toxicity-Single Exposure (STOT-SE) is necessary.

## 5.3 Irritation

### 5.3.1 Skin Irritation

In a study with rabbits, abamectin did not cause any irritation of the skin (table 5.3-1).

Table 5.3-1 Skin Irritation

Species	Method	Average score 24, 48, 72 h		Reversibility yes/no	Result	Reference
		Erythema	Oedema			
Rabbits; New Zealand White	Not fully in compliance with OECD 404 (but rated acceptable in 91/414/EEC DAR	0, 0, 0	0, 0, 0	-	negative	Robertsen (1981b)

### 5.3.2 Eye irritation

In a study with rabbits, abamectin did not cause any irritation of the eyes (table 5.3-2).

Table 5.3-2 Eye irritation

Species	Method	Average Score 24, 48, 72 h				Result	Reversibility yes/no	Reference
		Cornea	Iris	Conjunctiva Redness	Chemosis			
Rabbits; New Zealand White	in accordance with OECD 405	0	0	0	0	Not irritating	-	Glaza (2000)

### 5.3.3 Respiratory tract

In an acute inhalation toxicity study (Ruddock, 2001) dark, darkened or red areas in the lungs and inflated lungs were observed in all dose groups (0.21-5.25 mg/L).

### 5.3.4 Summary and discussion of irritation

#### Classification proposals according to Directive 67/548/EEC

Abamectin is considered not irritating to skin or eyes according to the criteria of Annex VI of Directive 67/548/EC.

The description of the effects on the lung in the acute toxicity study is limited. Since in repeated dose studies histological examination revealed no signs of respiratory irritation and since abamectin is not irritating to the eyes and the skin no classification for respiratory irritation is required.

#### Classification proposals according to Regulation EC 1272/2008

Abamectine also needs not to be classified for skin and eye irritation according to the criteria in the new EU C&L Regulation based on GHS. For reasons described above abamectin needs not to be classified for respiratory irritation.

### 5.4 Corrosivity

Based on the data from the skin irritation study it can be concluded that abamectin is not corrosive.

### 5.5 Sensitisation

#### 5.5.1 Skin

Abamectin showed no skin sensitizing properties in a Guinea Pig Maximization test (table 5.5-1).

Table 5.5-1 Sensitisation

Species	Method	Number of animals sensitised/total number of animals	Result	Reference
Guinea Pig	GPMT in accordance with OECD 406	0/19	negative	Ruddock (2001b)

### 5.5.2 Respiratory system

No data.

### 5.5.3 Summary and discussion of sensitisation

Abamectin needs not to be classified for skin or respiratory sensitization according to Directive 67/548/EC.

Abamectine needs not to be classified for skin or respiratory sensitization according to Regulation EC 1272/2008.

## 5.6 Repeated dose toxicity

No repeated exposure toxicity data in humans are available.

### 5.6.1 Repeated dose toxicity: oral

In the rat an 8-week and a 90-day dietary study were performed. In the dog 12, 18 and 53-week toxicity studies have been performed by dietary, gavage and dietary administration respectively. The studies were performed using abamectin except the 18 week toxicity study in dogs which used avermectin B1a. The 8-week study in the rat and the 12 week study in the dog were range finding studies, with determination of very few parameters, not in accordance with OECD guidelines.

In a 90-day study of (neuro-)toxicity in rats abamectin, administered daily by gavage at 4 mg/kg bw/day induced low incidences of clinical signs from week 2 onwards. In these animals a marked increase in clinical signs (shaking, tiptoe gait, reduced righting reflex, reduced stability, reduced splay reflex, hunched posture, “pinched-in” sides, subdued behaviour, irregular breathing, decreased activity, stains around the mouth or nose, upward spinal curvature) and body weight loss occurred in week 7 of treatment. The animals were killed for humane reasons. Pathological examinations revealed macroscopic and histological changes in the stomach. The NOAEL in this study was 1.6 mg/kg bw/day. The LOAEL was 4.0 mg/kg bw/day.

In the 18 week oral toxicity study with dogs, a very steep dose-response relationship for avermectin B1a in the dog was observed, since the oral NOAEL by gavage is 0.25 mg/kg bw/day and death, clinical signs (ataxia, tremors, mydriasis and ptyalism), reduced weight gain and histopathological changes in the liver occurred at 0.5 mg/kg bw/day. At the highest dose these effects were observed after the first dose. At the lower dose levels, the effects were observed after several exposures.

In the 53-week oral toxicity study with abamectin in dogs, death occurred at the high dose level of 1.0 mg/kg bw/day, and pupil reactivity was decreased or absent at the dose level of 0.5 mg/kg bw/day. Based on this effect on pupil reactivity, the NOAEL in this study is 0.25 mg/kg bw/day. The results of both these studies show that a similar steep dose response exists for abamectin .

The overall NOAEL in the short-term toxicity studies is 0.25 mg/kg bw/day for both abamectin and avermectin B1a, observed in an 18-week and a 1-year study in the dog.

Table 5.6-1 Summary of neurotoxicity studies

Species	Study type or duration; Dose levels	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Effect at LOAEL	Reference
Rat	90-day oral study of (neuro-)toxicity; OECD 408, OECD 424. 0, 0.4, 1.6, 4 mg/kg bw/day	1.6	4	clinical signs (shaking, tiptoe gait, reduced righting reflex, reduced stability, reduced splay reflex, hunched posture,	Brammer, A. 2006b

by gavage	"pinched-in" sides, subdued behaviour, irregular breathing, decreased activity, stains around the mouth or nose, upward spinal curvature) and body weight loss and macroscopic and histological changes in the stomach
-----------	--

### Chronic toxicity

Long-term toxicity and carcinogenicity studies were performed in the rat and the mouse. There was no evidence of carcinogenicity in either the rat or the mouse at any of the dose levels employed. The long-term dietary administration of abamectin did not reveal any primary target organ toxicity. Although clinical signs of neurotoxicity were evident in rats and to a lesser extent in mice, no histopathologic correlate was evident. The overall NOAEL determined in long-term toxicity studies was 1.5 mg/kg bw/day found in the rat carcinogenicity and toxicity study.

Table 5.6-2 Summary of repeated dose toxicity studies

Test substance	Duration, route Dose levels	Species	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Critical effects	Reference/ Registrant
Abamectin (vehicle acetone)	4 or 8 weeks, oral 0, 5, 10, 15, 20/25, 40 (and 60) ppm (mean weekly achieved dose 0, 0.3-0.7, 1.0- 1.4, 1.6-2.2, 1.7-2.7 and 4.1-5.8 mg/kg bw/day)	rat	-	-	Range-finding study (only bw, food consumption and clin. signs)	Gordon, L.R. (1984b)
Abamectin (acetone)	12 weeks, oral 0, 0.25, 0.5, 1.0 and 4.0/2.0 mg/kg (0, 6, 13, 25, 100/50 ppm)	dog	(0.5)	(1.0)	Range-finding study (only bw, food consumption and pupil response)	Gordon, L.R. (1984c)
Avermectin B1a (vehicle sesame oil)	18 weeks, oral (gavage) 0, 0.25, 0.5, 2.0 and 8.0 mg/kg bw/day	dog	0.25	0.5	Mortality, clinical signs of toxicity (ataxia, tremors, mydriasis, ptyalism), reduced weight gain, histopathologic changes in the liver	Robertson, R.T. & Allen, H.L. (1976)
Abamectin (vehicle acetone)	53 weeks, oral (diet) 0, 0.25, 0.5 and 1.0 mg/kg bw/day	dog	0.25	0.5	Absent or decreased pupil reflex (death at 1.0 mg/kg bw/day)	Gordon, L.R. (1984d)
Abamectin (vehicle acetone)	105 weeks, oral (diet) 0, 0, 0.75, 1.5 and 2.0 mg/kg bw/day	Rat	1.5	2.0	Increased mortality in males, clinical signs (tremors, unthrifty appearance)	Gordon, L.R. (1985b)
Abamectin (vehicle acetone)	94 weeks, oral (diet) 0, 0, 2.0, 4.0 and 8.0 mg/kg bw/day	CD-1 mice	4.0	8.0	Increased mortality in males, reduced body weight gain in males and females, extramedullary haematopoiesis in spleen of males.	Gordon, L.R. (1985c)

### 5.6.2 Repeated dose toxicity: inhalation

In a preliminary study in rats (2/sex/dose) daily inhalation exposure for 5 consecutive days induced dose-dependent increases in clinical signs after exposure at all doses (1.03-24.7 µg/L). The severity of the clinical signs was such that at 9.59 and 24.7 µg/L (part of) the animals were humanely killed during the treatment period. Pathological examination revealed no relevant macroscopic or microscopic effects.

A repeated dose inhalation study was described in the addendum to the DAR (February 2008). Daily inhalation exposure (nose only) of rats for 6 h/day, 5 days/week over a 30 day period (total of 21 exposures) induced clinical signs and reduced motor activity at 2.69 µg/L. The NOAEC was 0.577 µg/L.

Pathological examination revealed no relevant macroscopic or microscopic effects.

Table 5.6-3 summarises the repeated dose inhalation toxicity study.

Table 5.6-3 Summary of repeated dose inhalation toxicity study.

Test substance	Duration, route Dose levels	Species	NOAEC (µg/L)	LOAEC (µg/L)	Critical effects	Reference/ Registrant
Abamectin	6h/day, for 5 days.  Inhalation (nose only) 0, 0.103, 3.71, 9.59 and 24.7 µg/L.	Rat			Dose-dependent increase in number and severity of clinical signs (splay reflex, hunched posture reduced foot splay reflex, tremors, decreased activity, piloerection, shaking, reduced stability, pale skin, tail erection, reduced breathing rate, decreased visual placing response), and body weight loss and reduced food consumption	Pinto, P.J. (2006a)
Abamectin	6h/day, 5 days/week over a 30-day period. Inhalation (nose only) 0, 0.111, 0.577 and 2.69 µg/L.	rat	0.577	2.69	Clinical signs (lying prostrate, shaking and gasping, with a swollen head. Ungroomed appearance, with stains around the mouth, hunched posture and piloerection, abnormal respiratory noise) and decreased motor activity.	Pinto, P.J. (2006b)

### 5.6.3 Repeated dose toxicity: dermal

No repeated dose studies were available.

Acute dermal toxicity studies with rat and rabbit has shown that abamectin has a low order of toxicity. A dermal penetration study with monkeys has shown that less than 1% of abamectin is absorbed through the skin. Based on these findings, percutaneous exposure will not be a significant route of exposure.

#### 5.6.4 Other relevant information

None.

#### 5.6.5 Summary and discussion of repeated dose toxicity:

##### Effect:

With respect to oral exposure, based on the severity of clinical signs of neurotoxicity and mydriasis and the dose levels at which death occurs, the dog is more sensitive than the rat to abamectin. Repeated dose dietary administration of abamectin reveals that the nervous system is a primary target organ for toxicity. A steep dose response curve exists for this effect. Although clinical signs of neurotoxicity occur in all species evaluated, no histopathologic correlates are evident in central or peripheral nerves. In addition, histopathologic changes in the liver of dogs and extramedullary haematopoiesis in the spleen of mice were observed.

With respect to inhalation toxicity the data in the rat study indicate that the nervous system is a primary target organ for toxicity.

##### Dose:

The overall NOAEL in oral repeat dose toxicity studies is 0.25 mg/kg bw/day for both abamectin and avermectin B1a, observed in an 18-week and a 1-year study in the dog. This is about 30 times lower than the acute oral LD50.

The overall NOAEC in inhalation repeat dose toxicity studies is 0.577 µg/L for both abamectin, observed in a 30 days study in the dog. The LOAEC was 2.69 µg/L. This is about 20 times lower than the acute LC50.

##### Relevant repeated exposure effect levels for classification

Clear signs of oral neurotoxicity were observed in a 90-day study in rats at a dose of 4 mg/kg bw/day. In an 18-week oral (gavage) study in dogs severe signs of toxicity, including mortality, were observed at 0.5 mg/kg bw/day. In a 2-year dietary study in rats severe signs of toxicity, including mortality, were observed at 2.0 mg/kg bw/day. Clear signs of neurotoxicity were also observed in a 30-day inhalation study (6h/day, 5 days/week) in rats at a NOAEC of 0.00269 mg/L.

##### Classification proposal according to Directive 67/548/EEC

In view of the effects and effect levels for oral and inhalation (neuro-)toxicity in repeated exposure studies abamectin should be classified with R48/23/25: Toxic: danger of serious damage to health by prolonged exposure through inhalation and if swallowed. There was clear neurotoxicity at 0.00269 mg/L which is below the guidance value for R48/23 in a 30 day inhalation study of 0.075 mg/L. In the oral dog study, neurotoxicity was observed at dose levels below the guidance value for R48/25 of 5 mg/kg bw/day in a 13-week study.

##### Classification proposal according to Regulation EC 1272/2008

In oral repeated dosing studies in animals (rats and dogs) abamectin appears to be (neuro-)toxic at doses < 10 mg/kg bw/day. In a 30-day repeated exposure inhalation study in rats abamectin is neurotoxic at concentrations of 0.00269 mg/L and above (range-finding study). This is below the guidance value for STOT-RE Cat 1 in a 30 day inhalation study of 0.06 mg/L. In the oral dog study,

neurotoxicity was observed at dose levels below the guidance value of 10 mg/kg bw/day in a 13-week study. According to CLP criteria abamectin should be classified with STOT/RE cat. 1.

## 5.7 Mutagenicity

### 5.7.1 In vitro data

Table 5.7-1 summarises in vitro genotoxicity studies. Abamectin does not induce gene mutations in either bacteria or mammalian cells with or without metabolic activation. There is no evidence of clastogenicity in an in vitro test system.

Table 5.7-1 Genotoxicity studies: In vitro

Test system Method Guideline	organism/ strain(s)	concentrations tested (give range)	Result		Remark give information on cytotoxicity and other	Reference
			+	-		
Point mutation OED 471	S. typhimurium (5 strains)	3, 10, 30, 100, 300, and 1000 µg/plate	neg	neg		Gordon L.R. (1986a)
Point mutation OED 471	S. typhimurium (5 strains) & E.coli (1 strain)	312.5, 625, 1250, 2500, 5000 µg/plate	neg	neg		Deperade, E (2001)
Chromosome aberrations (in vitro) OECD 473	Chinese hamster ovary cells (CHO-WBL)	-S9: 0.0100, 0.0150, 0.0200, 0.0250, 0.0300 and 0.0350 mM +S9: 0.0050, 0.0100, 0.0150, 0.0200 and 0.0250 mM Solvent: DMSO	neg	neg		Gordon L.R. (1986b)
Mammalian point Mutations OECD 476	Chinese hamster lung cells (V79)	-S9 <sup>1</sup> : 0.003, 0.004, 0.005 and 0.006 mM +S9: 0.03, 0.04, 0.045 and 0.05 mM Solvent: DMSO	neg	neg		Gordon L.R. (1983)

<sup>1</sup> Due to a dilution error, the two lowest concentrations tested without S9 in the repeat assay were 0.0003 and 0.0004 mM

### 5.7.2 In vivo data

Table 5.7-2 summarises in-vivo genotoxicity studies. Abamectin does not induce cytogenic damage in male mouse bone marrow cells.

Table 5.7-2 Genotoxicity studies: In vivo

Test system Method Guideline	species strain(s)	Dose levels tested (give range)	Result	Remark give information on cytotoxicity and other	Reference
Structural chromosome aberration OECD 475	Mouse (male CD-1)	0, 1.2, 4.0 and 12.0 mg/kg bw	negative		Blazak, WF (1983)

### 5.7.3 Human data

No data available.

### 5.7.4 Other relevant information

None.



### 5.7.5 Summary and discussion of mutagenicity

### 5.7.6 Conclusion

Abamectin did not induce gene mutations in either bacterial or mammalian cells at any of the tested concentrations either with or without metabolic activation. There was no evidence of a clastogenic effect at any tested concentration either in vitro or in vivo. It is concluded that abamectin and/or its metabolites are not genotoxic.

Abamectine needs not to be classified for mutagenicity according to Directive 67/548/EEC or Regulation EC 1272/2008.

## 5.8 Carcinogenicity

### 5.8.1 Carcinogenicity: oral

Table 5.8-1 summarises carcinogenicity studies. Long-term toxicity and carcinogenicity studies were performed in the rat and mouse. There was no evidence of carcinogenicity in either the rat or the mouse at any of the dose levels employed.

Table 5.8-1 Carcinogenicity study

Route	Species Strain Sex no/group	dose levels frequency of application	Tumours	Reference
Oral in diet	Rat (CD(SD)BR)	105 weeks, oral 0, 0, 0.75, 1.5 and 2.0 mg/kg bw/day	none	Gordon, LR (1985b)
Oral in diet	Mouse (CD-1)	93 weeks, oral 0, 0, 2.0, 4.0 and 8.0 mg/kg bw/day	none	Gordon, LR (1985c)

### 5.8.2 Carcinogenicity: inhalation

No data available.

### 5.8.3 Carcinogenicity: dermal

No data available.

### 5.8.4 Carcinogenicity: human data

No data available.

### 5.8.5 Other relevant information

None.

### 5.8.6 Summary and discussion of carcinogenicity

Abamectin is unlikely to pose a carcinogenic hazard.

Abamectin needs not to be classified for carcinogenicity according to Directive 67/548/EEC or to Regulation EC 1272/2008.

## 5.9 Toxicity for reproduction

### 5.9.1 Effects on fertility

Table 5.9-1 summarises the available fertility studies.

A rat 2-generation reproductive toxicity study with abamectin was available. For the plant protection evaluation the registrant provided additional data to the study report (see revised addendum to pesticides Draft Assessment Report, February 2008).

The original study report did not include all relevant parameters. Furthermore, the study report incorrectly suggested that in this study fertility of the rats was affected by abamectin treatment. However, the registrant provided additional information on the fertility study and recalculated some reproduction parameters. The additional information and the recalculated reproduction parameters were evaluated by the rapporteur in 2008. The conclusion of the re-evaluation of this 2-generation reproductive toxicity study was reported in an addendum to the DAR. Thus it was concluded that in this multigeneration reproductive toxicity study in the rat the NOAEL for parental and reproduction toxicity is 0.4 mg/kg bw/day, i.e. the highest dose tested (see revised addendum to pesticides Draft Assessment Report, February 2008).

In the 2-generation study, pup mortality for both F1a and F1b litters was significantly increased at 0.4 mg/kg bw/day, with most pups dying days 5-15 postpartum. Post mortem examination of F1b weanlings showed retinal anomalies (single or multiple retinal folds of many layers of the retina) in 3 out of 4 males in the highest dose group.

Group mean body weights of F1 males and females at 0.4 mg/kg bw/day and the females at 0.12 mg/kg bw/day were significantly reduced at the start of treatment, due to retarded pre-weaning growth. Treatment-related reduced weight gain continued in males at 0.4 mg/kg bw/day for 4 weeks, after which weight gain was enhanced and terminal body weights were comparable to controls. This temporary effect on body weight is considered not a relevant endpoint for determination of the LOAEL. Retinal anomaly was observed in pups only, and appeared to be transient, and was not observed in the adult F1 animals.

In both F2a and F2b litters treated at 0.4 mg/kg bw/day pup mortality significantly increased during the course of lactation, and the associated viability and lactation indices significantly decreased. Pup weight in the high dose group was unaffected by treatment directly after birth and for the first few days, but were significantly reduced from day 7 to day 21. This was associated with increased numbers of pups that were thin, weak and not nursing. The number of male pups was decreased in the high dose group (F2a and F2b). Post mortem examination of F2b weanlings showed retinal anomalies, with characteristics identical to those observed in F1b animals in 10/63 males and 18/66 females in the highest dose group. As in the F1 pups, it is considered that these retinal anomalies are transient and confined to the pup stage.

Based on the occurrence of increased pup mortality and retarded weight gain in both F1 and F2 generation progeny, increased incidence of total litter loss, decreased lactation index and reduced

weight gain in the F1 and F2 generation weanlings at the highest dose, the NOAEL for pup toxicity in this study is 0.12 mg/kg bw/day.

Significant neonatal mortality seen in rat pups is likely to be the result of a lack of p-glycoprotein expression in the neonatal rat brain. P-glycoprotein dependent xenobiotic efflux in the blood brain barrier is considered to play an important role in attenuating neurotoxicity of avermectins. However, brain p-glycoprotein expression starts early in human development, having been detected in human foetal brain microvessels as early as week eight of pregnancy. Expression of p-glycoprotein in the cerebrum and cerebellum of rats is not fully developed in neonate rats and expression in the jejunal epithelial brush borders does not start before post-natal day 8. Adult levels are reached at post-partum day 20 or 28. Therefore, the increased postnatal mortality is considered not to be relevant to human risk assessment and also not for classification and labelling. For further information on the role of p-glycoproteins in abamectin toxicity see section 5.10.1.

Table 5.9-1 Fertility study

Species	Study type or duration; Dose levels	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Effect at LOAEL	Reference
rat	Two-generation study in rats with abamectin ; OECD 416. 0, 0.05, 0.12 and 0.4 mg/kg bw/day	0.4	--	<b>Parent:</b> No effects	Gordon, LR (1984e)
		0.4	--	<b>Reproduction:</b> No effects	
		0.12	0.4	<b>Fetes/pups:</b> increased postnatal pup mortality, retarded weight gain pups (F1 and F2), increased incidence of total litter loss, decreased lactation index, increased incidence of retinal anomaly in the eyes of pups (F1 and F2)	

## 5.9.2 Developmental toxicity

Two developmental toxicity studies (rat and rabbit) with abamectin were available.

### *Rats*

In a teratogenicity study in rats the sex ratio (m:f) was lower at 1.6 mg/kg bw/day. Since exposure to abamectin was from days 6-19 of gestation, abamectin could not have affected the sex of the fetuses directly. Apparently, abamectin exposure in the highest dose group affected resorption in a sex-specific way (more effect on female fetuses), resulting in a lower m:f ratio but within the historical control range of 1 : 0.69 to 1 : 1.22. However, there was no increase in the total number of resorptions.

In the 0.8 mg/kg bw/day group a significant higher incidence of resorptions and decreased fetal weight were observed. Similar effects were not observed at 1.6 mg/kg bw/day, and therefore these effects are considered incidental. At 1.6 mg/kg bw/day, exencephaly was observed paired with a conjoined twin, which is a spontaneous congenital abnormality, and thus it is likely that this effect is not substance-related. The observed incidence of one animal with cleft palate in the highest dose groups is considered treatment-related, since this effect is also observed in the developmental toxicity study with abamectin in the rabbit and in the developmental toxicity study with the main isomer of abamectin in CD-1 strain mice. Furthermore, historical control data provided by the registrant in 2005 showed that in 23 studies only one fetus with cleft palate was observed.

In the highest dose group, the number of pups with lumbar rib and with lumbar count variation had increased but remained within the historic control data.

The developmental effects in the study in rats are summarized in the table below.

Table 5.9-2 Developmental effects in the rat study.

Dose (mg/kg bw/day)		0	0.4	0.8	1.6
	No. of dead fetuses/no. of fetuses studied	0/319	0/320	0/279	0/326
	Malformations -exencephaly -cleft palate			1 <sup>b</sup>	1 <sup>a</sup> 1
	Skeletal deviations -lumbar rib (no / %) -lumbar count variation (no / %) - no. of litters with fetal variations / no. of litters examined	44 / 14 1 / 0.3 13 / 23	41 / 13 1 / 0.3 18 / 24	45 / 16 1 / 0.4 14 / 24	72 / 22 5 / 1.5 16 / 24

a: conjoined twin

b: anasarca, micrognathia, cleft palate, protruding tongue, ectromelia

Based on the absence of effects in the highest dose group, the NOAEL for maternal toxicity in this study is 1.6 mg/kg bw/day.

Based on the occurrence of cleft palate, changed sex ratio and increased number of fetuses with lumbar rib and lumbar count variation in the highest dose group, the NOAEL for developmental toxicity in this study is 0.8 mg/kg bw/day.

### Rabbits

In a teratogenicity study with rabbits two deaths and one premature sacrifice occurred in abamectin-treated groups. Death was preceded by reduced food and water consumption in 2 animals and by blood-stained urine in the cage of the other animal. The relationship of these deaths to treatment with abamectin is equivocal since a dose-related increase in incidence did not occur. There were no clinical signs of toxicity at any dose level.

The food and water consumption of all groups was variable, but by subjective assessment, the periods of reduced food and water consumption in the group treated at 2.0 mg/kg bw/day were more prolonged and pronounced than in the other groups. This treatment-related maternal toxicity at 2.0 mg/kg bw/day manifested as decreased food and water consumption resulted in weight loss during the dosing period which was statistically significant between day 6 and 18 of gestation compared to control. The average weight loss at 2.0 mg/kg bw/day over this period was 64 g compared to a weight increase of 64 g in the control group. The effects observed at 2.0 mg/kg bw/day are considered as evidence for maternal toxicity but not as marked maternal toxicity because the differences in maternal body weight gain are only small (3%) compared to the average weight of a rabbit of approximately 4 kg.

There were no treatment-related effects at any dose level on pre-implantation loss and post implantation loss, and mean foetal weight (sexes combined) at any dose level. Higher numbers of dead fetuses and an increased m/f sex ratio was observed in the group treated at 1.0 mg/kg bw/day, but not at the higher dose level. Therefore, these effects are considered incidental.

In the high dose group, the number of resorptions and the % malformed fetuses were increased. In the high dose group one litter contained 2 fetuses with cleft palate and 2 fetuses with omphalocele. In this litter 3 fetuses had sternebral malformations, including one of the fetuses with cleft palate. In 3 other litters of the high dose group in total 5 fetuses with clubbed fore-feet were found. One fetus with clubbed fore-feet also had a lumbar vertebral malformation. The incidences of these malformations are higher than the concurrent and historical control groups (not available) and were considered treatment related (by the study author).

Two fetuses in one litter from a female treated at 1.0 mg/kg bw/day also had clubbed fore-feet but the occurrence is considered not to be treatment-related because higher incidences of the defect have been recorded in historical controls (not available), one fetus from a concurrent control female also had a clubbed fore-foot, and no other malformations were observed at this dose.

The study report contains no information to relate the malformed pups with individual dams and their weight changes over pregnancy.

At 2.0 mg/kg bw/day, increased incidences of incomplete ossification of sternebrae and metacarpals are considered to reflect a treatment-related slight delay in ossification.

The developmental effects in the study in rabbits are summarized in the table below.

Table 5.9-3 Developmental effects in the rabbit study.

Dose (mg/kg bw/day)		0	0.5	1.0	2.0	dr
Litter response	Live fetuses	No toxicologically relevant effects				
	Fetal weight	No toxicologically relevant effects				
	Resorptions/implants (litter mean)	0.049	0.038	0.036	0.065	
	Pre implantation loss	No toxicologically relevant effects				
	Foetal implantation loss	No toxicologically relevant effects				
	Post implantation loss	No toxicologically relevant effects				
Fetus examination	No. of abnormal fetuses	No toxicologically relevant effects				
	No. of dead fetuses / no. of fetuses studied	0/97	1/91	5/100	0/121	
	Sex ratio (m:f)	1 : 0.98	1 : 1.07	1 : 1.17	1 : 1.02	
	% malformed fetuses	3.1	4.4	4.0	12.4	
	External observations and visceral deviations					
	-cleft palate	0	0	0	2 <sup>a</sup>	
	-clubbed fore-foot	1	0	2	5 <sup>a</sup>	
	-omphaloceles	1	0	0	2 <sup>a</sup>	
	Skeletal deviations					
	-sternebral malformation	0	0	0	3	
	-incompletely ossified sternebra	17	17	16	42	
	-incompletely ossified metacarpal	8	15	7	33	
	-incompletely ossified phalanx	19	27	12	31	

a: The 2 fetuses with cleft palate and 2 fetuses with omphaloceles were all from a single litter and 5 fetuses with clubbed fore-foot were from 3 other litters.

### Summary of developmental toxicity studies

Table 5.9-4 Summary of teratogenicity studies

Species	Study type or duration; Dose levels	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Effect at LOAEL	Reference
Rat	Oral developmental study (gavage); OECD 414. Day 6-19 of gestation 0, 0.25, 0.5, 1.0 and 2.0 mg/kg bw (range-finding study)	Maternal: 1.6 Developm: 0.8	>1.6 1.6	- Cleft palate, lumbar rib and lumbar count variation	Gordon, L.R.(1982a)

	0, 0.4, 0.8 and 1.6 mg/kg bw/day (main study)				
Rabbit	Oral developmental study (gavage); OECD 414. Day 6-27 of gestation 0, 0.5, 1.0, 2.0 and 3.0 mg/kg bw (range-finding study)	Maternal: 1.0	2.0	food consumption and weight loss during gestation, increased number of resorptions.	Gordon, L.R. (1982b)
	0, 0.5, 1.0 and 2.0 mg/kg bw/day (main study)	Developm: 1.0	2.0	Developmental: cleft palate, omphalocele, clubbed fore-feet and delayed ossification	

The NOAEL of abamectin for maternal toxicity in rabbits in this study is 1.0 mg/kg bw/day, based on decreased water and food consumption and weight loss during gestation at 2.0 mg/kg bw/day.

The NOAEL of abamectin for foetal toxicity was also established at 1.0 mg/kg bw/day based on the occurrence of increased number of resorptions, delayed ossification and excess incidences of cleft palate, omphalocele and clubbed fore-feet at the maternally toxic dose level of 2.0 mg/kg bw/day.

### 5.9.3 Human data

No data available.

### 5.9.4 Other relevant information

None

### 5.9.5 Summary and discussion of reproductive toxicity

#### *Fertility*

In the 2-generation study of reproductive toxicity no effect on reproductive parameters were found.

#### Classification proposals according to Directive 67/548/EEC and Regulation EC 1272/2008

Abamectine needs not to be classified for reproductive effects according to Directive 67/548/EEC or to EC 1272/2008.

#### *Developmental toxicity*

In the reproductive toxicity study no effects were observed in the pups at the time of birth. It should be noted however that this study was not designed to investigate prenatal developmental effects of abamectin.

In the developmental toxicity studies in the rat and the rabbit teratogenic effects were observed, albeit at low incidences. In the rat, developmental toxicity was observed in the absence of maternal toxicity.

#### Classification proposals according to Directive 67/548/EEC

In view of the low incidences of the developmental effects it is considered that there is some but no clear evidence of a developmental effect in rats. In the developmental toxicity study in the rat there is actually only 1 cleft palate to take into consideration for classification and labelling. As stand alone study, this would not be considered relevant for classification and labelling, but we agree that in combination with the rabbit study, this one cleft palate should be considered.

In the rabbit study there is one 'strange' litter with 2 fetuses with cleft palate and 2 fetuses with omphaloceles. The relevance of these findings for classification and labelling can be questioned. The increase in malformations (clubbed fore-foot) in rabbits at the highest dose is above the concurrent and historic controls and therefore treatment related. The increase in these malformations was small and therefore considered as evidence but no clear evidence. This effect was observed in presence of maternal toxicity including weight loss and decreased water and food consumption. The difference in body weight gain is considered small in relation to the average weight of a rabbit and is not considered as marked maternal toxicity. Also, it is considered unlikely that the increased incidence in malformation is caused by the reduced body weight gain. The increase in club fore-foot is considered to be a direct effect of the substance and not a secondary consequence of the maternal toxicity. As the time of development of this effect is unknown, it is unknown whether the differences in p-glycoprotein development between rats and humans is also important for this effect. Therefore, it is assumed that this effect is also relevant to humans. It is proposed to classify abamectin for harm to the unborn child as Repro Cat. 3; R63 based on an increase (but no clear increase) in malformations (clubbed fore-foot) which is considered not secondary to maternal toxicity and relevant to humans.

#### Classification proposals according to Regulation EC 1272/2008

The same argumentation as provided above for classification according to Directive 67/548/EEG applies also for classification according to Regulation EC 1272/2008. It is proposed to classify abamectin with Repr Cat 2 H361d.

#### *Effects on or via lactation*

The increase in post-natal mortality in the 2-generation study in rats at 0.4 mg/kg bw/day is most likely an effect on or via lactation. This is confirmed by a cross fostering study with the closely related substances ivermectine indicates that the neonatal toxicity was primarily a function of postnatal exposure ((Merck & Co., Inc., 1980f as summarised by JECFA, 1991). Therefore, these effect would be considered relevant for effects on or via lactation and not for developmental effects. Significant neonatal mortality seen in rat pups is likely to be the result of a lack of p-glycoprotein expression in the neonatal rat brain. P-glycoprotein dependent xenobiotic efflux in the blood brain barrier is considered to play an important role in attenuating neurotoxicity of avermectins. However, brain p-glycoprotein expression starts early in human development, having been detected in human foetal brain microvessels as early as week eight of pregnancy. Expression of p-glycoprotein in the cerebrum and cerebellum of rats is not fully developed in neonate rats and expression in the jejunal epithelial brush borders does not start before post-natal day 8. Adult levels are reached at post-partum day 20 or 28. Therefore, the increased postnatal mortality is considered not to be relevant to human risk assessment. For further information on the role of p-glycoproteins in abamectin toxicity see section 5.10.1.

#### Classification proposals according to Directive 67/548/EEC

No classification for effects on or via lactation is proposed because the increased post natal toxicity observed in the 2-generation study in rats is not considered relevant to humans.

#### Classification proposals according to Regulation EC 1272/2008

No classification for effects on or via lactation is proposed because the increased post natal toxicity observed in the 2-generation study in rats is not considered relevant to humans.

## 5.10 Other effects

### 5.10.1 P-glycoprotein expression and increased susceptibility to abamectin

From submitted, it was concluded (by the registrant) that CF-1 mice exhibit typical clinical signs of neurotoxicity and are increased susceptible to abamectin toxicity. From those studies it was suggested that the increased susceptibility of CF-1 mice (compared to CD-1 mice) is related to the accessibility of the 8,9-Z isomer to the target organ, and hence to the presence or absence of p-glycoprotein expression. In order to investigate this suggestion, several studies were performed to investigate the relation between p-glycoprotein and the increased sensitivity of CF-1 mice to abamectin and the 8,9-Z isomer.

Comparative studies of the acute oral toxicity of abamectin in pregnant and non-pregnant CF-1 mice and a maternal toxicity study by dietary administration during gestation have been performed. The influence of the *mdr1* genotype and p-glycoprotein levels on the expression of abamectin toxicity were investigated in two exploratory studies in CF-1 mice of known genotype and in CF-1 and CD-1 strain mice of unknown genotype. A summary of these studies is presented in the table below.

#### Summary of supplementary studies

Study/ species dose levels	NOAEL (mg/kg bw/day)	LOAEL (mg/kg bw/day)	Effects at LOEL	Reference
10-day dietary maternal toxicity; CF-1 mice;	- maternal: 0.08 (time-weighted)	- maternal: 0.24 (time-weighted)	Tremors, hunched posture, poor condition	Gordon, L.R. (1984g)
Acute oral toxicity; Pregnant / non-pregnant CF-1 mice	LD50 non-pregnant mice: >20 and <40 mg/kg bw  LD50 pregnant mice: 19 mg/kg bw	< 5  5	Deaths, tremors, bradypnea  Deaths, tremors, bradypnea	Gordon, L.R. (1986h)
Acute oral toxicity; Pregnant / non-pregnant CF-1 mice	LD50 non-pregnant mice: 15.0 mg/kg bw  LD50 pregnant mice: 11.8 mg/kg bw	< 5  < 5	Death, loss of righting reflex, bradypnea  Death, tremors, bradypnea, clonic convulsions	Gordon, L.R. (1986h)
Exploratory acute oral toxicity; CF-1 mice of known genotype for p-glycoprotein	LD50 (+/+ genotype female mice): 28 mg/kg bw  LD50 (+/- genotype female mice): 14 mg/kg bw	< 10  < 10	Tremors, bradypnea, decreased activity.  Tremors, bradypnea, decreased activity, weight loss during first week	Hall, S. (1997)



Exploratory oral toxicity; CF-1 / CD-1 mice (dose = 0.8 mg/kg bw for 4 days)	Results: All CF-1 mice showed tremors and ataxia, but 17% also showed dyspnea, lateral recumbence and coma (= sensitive to abamectin toxicity). All but one sensitive animal had no detectable p-glycoprotein in brain and small intestine.  All insensitive CF-1 mice evaluated and all CD-1 mice had detectable p-glycoprotein levels.  Control and treated CD-1 mice had similar levels of p-glycoprotein.	Lankas, G.R. (1994)
Exploratory study of p-glycoprotein development in rat fetuses and pups.	Results: the expression of p-glycoprotein in the cerebrum and cerebellum is not fully developed in neonate rats. P-glycoprotein expression reaches adult levels by post-natal day 20. Expression of p-glycoprotein in the jejunal epithelial brush border does not start before post-natal day 8. It is suggested that neonate rats with limited or no p-glycoprotein expression have an increased susceptibility to avermectin toxicity.	Cukierski, M.A. (1995), Lankas, G.R. (1996b, addendum)
Examination of developmental expression of p-glycoprotein levels in rat pups  Rats postnatal days 1,3,7,14,21,28,56,84 examined	Results: p-glycoprotein was first detected at post-natal day 7 in pups, with subsequent increases to plateau at adult levels by post-natal day 28. In the adult rat brain, p-glycoprotein was detected predominantly in the membrane fraction. Double immunostaining of p-glycoprotein and von Willebrand factor demonstrated that p-glycoprotein was co-localised with brain capillaries, suggesting a role for p-glycoprotein in the blood brain barrier.	Matsuoka, Y. et al. (1999)

Dietary administration of abamectin to pregnant CF-1 mice during organogenesis resulted in clinical signs of neurotoxicity at time-weighted average maternal dose levels above 0.08 mg/kg bw/day, whereas no treatment-related effects were observed on reproductive parameters.

In two studies with pregnant and non-pregnant CF-1 mice, singly orally exposed to abamectin at day 10, 11 or 12 of gestation, it was shown that the LD50's in pregnant animals were slightly, not statistically significantly lower (LD50 = 19 mg/kg bw and LD50 = 11.8 mg/kg bw in study 1 and 2, respectively) compared to the LD50's in non-pregnant mice (LD50 = between 20 and 40 mg/kg bw and LD50 = 15 mg/kg bw in study 1 and 2, respectively). Typical clinical signs of neurotoxicity (tremors, clonic convulsion and bradypnea) occurred in both pregnant and non-pregnant animals.

In a study with female CF-1 strain mice, heterozygous (+/-) or homozygous positive (++) for the *mdr1* gene (which codes for p-glycoprotein expression), the LD50 for abamectin in homozygous positive (++) female mice was 28 mg/kg bw, whereas the LD50 in heterozygous female mice was 14 mg/kg bw.

In a comparative study with CF-1 mice and CD-1 mice, it was demonstrated that 17% of a random population of CF-1 mice are sensitive to abamectin toxicity, showing signs of neurotoxicity (tremors, ataxia, dyspnea, lateral recumbency, coma) in response to 0.8 mg/kg bw/day abamectin for 4 days. Sensitive CF-1 individuals were shown to express no p-glycoprotein in the cerebrum, cerebellum and jejunum, whereas “non-sensitive” CF-1 mice and all CD-1 mice were shown to express p-glycoprotein in these tissues. Control and treated CD-1 mice had similar levels of p-glycoprotein.

#### *Role of p-glycoprotein in limiting avermectin toxicity*

C57BL/6 derived *abcb1a* knockout mice and some CF-1 mice were found to exhibit ivermectin sensitivity. CF-1 mouse ivermectin sensitivity exhibited classic Mendelian inheritance patterns, and has since been shown to be due to retroviral insert in exon 23 of

the *abcb1a* gene in some CF-1 mice. This results in total absence of properly transcribed, functional pgp in CF-1 mice homozygous for the disrupted form of the gene.

In both the CF-1 and the C57BL/6 *mdr1a* null mice models oral ivermectin dosing results in plasma ivermectin concentrations 2.5-fold to 3.3-fold higher in pgp null mice than in the wild type mice 24 h after dosing. Lack of pgp dependent efflux at the Blood Brain Barrier (BBB) also allows vastly increased brain penetration of avermectins. Brain ivermectin concentrations 24 h post dosing are between 33-fold and 87-fold higher in pgp null mice compared to wild type mice. Studies in our laboratory have shown similar results for two other avermectins, emamectin and abamectin, which are used predominantly as pesticides. Homozygous pgp null (*abcb1a*  $-/-$ ) mice show increased susceptibility to 0.2 mg/kg oral abamectin, while heterozygous (*abcb1a*  $+/-$ ) mice and wild type mice (*abcb1a*  $+/+$ ) are insensitive to up to 2.5 mg/kg abamectin. LD50 data indicates that at very high doses heterozygous mice are slightly more abamectin sensitive than homozygous wild type mice ( $-/+$  LD50 = 14 mg/kg,  $+/+$  LD50 = 30 mg/kg,  $-/-$  LD50 = 0.3 mg/kg). Thus although heterozygous mice express less brain pgp, a single copy of a functional *abcb1a* gene is sufficient for adequate pgp functionality in the mouse BBB at doses of avermectins used in the clinic (0.2 mg/kg), or resulting from worker pesticide exposure.

Where placental pgp activity is compromised avermectins can also exhibit developmental toxicity. In pgp null mice foetal avermectin exposure is associated with increased incidence of cleft palate. The placenta is a foetal tissue, and as such avermectin developmental toxicity is dependent on the *abcb1a* status of the fetus. CF-1 *abcb1a*  $-/-$  fetuses of mothers treated with abamectin have significantly higher concentrations of abamectin in their plasma than their *abcb1a*  $+/+$  and  $+/-$  littermates. Similarly when CF-1 dams were dosed with 1.5 mg/kg abamectin, all *abcb1a*  $-/-$  fetuses developed cleft palates, while none of their *abcb1a*  $+/+$  littermates and only 30% of their  $+/-$  littermates developed cleft palates.

Significant neonatal ivermectin neurotoxicity is seen in rat pups through a combination of ivermectin exposure of the offspring of ivermectin dosed rat dams via the dams' milk, and lack of pgp expression in the neonatal rat brain. However, this is not thought to be relevant to human risk assessment as brain pgp expression starts early in human development, having been detected in human foetal brain microvessels as early as week eight of pregnancy.

#### *P-glycoprotein human polymorphisms and pgp haplotypes*

Naturally occurring mutations that lead to non-functional pgp have been found in both the CF-1 strain of mice and dog breeds closely related to the collie. Millions of humans have received ivermectin as an anthelmintic treatment for river blindness without reports of major adverse neurological effects, although arguably adverse effect reporting may be less robust in the areas of the world where river blindness occurs. In addition, cumulatively more than 4,000 human volunteers have been genotyped for ABCB1 [although often only for known single nucleotide polymorphisms (SNPs)] without reports of major rearrangements of the ABCB1 gene similar to those in the CF-1 mouse and collie dog. Taken together this may indicate that individuals with significantly compromised pgp functionality analogous to that seen in the CF-1 pgp  $-/-$  mouse are rare.

More than 50 naturally occurring SNPs have been identified in the human ABCB1 gene. The vast majority are silent, i.e. they either do not occur in the coding region of the gene, or due to the inherent redundancy of codon usage they do not alter the amino acid sequence

of the protein. As has been extensively reviewed elsewhere there are numerous conflicting reports of the effects of individual ABCB1 SNPs on pgp expression and function in various tissues. Also the submitted publication gives an extensive overview of publications on the effects of individual ABCB1 SNPs. The conclusion is that there is no clear pattern of clinical effect of individual SNPs on pgp mediated efflux.

It is therefore suggested that combinations of human SNPs (haplotype) may be important in determining phenotype. An overview of the literature on pgp haplotypes is presented in the publication. This includes studies in which human BBB pgp function has been measured directly. Although various human ABCB1 haplotypes and/or SNPs have been reported to alter pgp function in relation to gut absorption, at present there is no conclusive data indicating that any of the known common haplotypes, including homozygosity for the most common minority haplotype, result in a significant loss of BBB pgp functionality. This would tend to indicate that the CF-1 *mdr1a*  $-/-$  mouse strain, which completely lacks pgp BBB functionality, is not a representative model for assessing risk in humans homozygous for any of the known haplotypes.

#### *Population distribution of pgp haplotypes*

Populations with different ethnicities are known to have different distributions of the various pgp haplotypes. Forty-eight and 79% of ABCB1 haplotypes found in the African American and Caucasian populations, respectively, produce a pgp identical to the reference amino acid sequence. Of the remainder, 38% of African American and 7.5% of Caucasian ABCB1 genes represent a haplotype which contains only one nonsynonymous SNP. Data from in vivo studies indicates that alleles in these two categories both produce pgp that is functional in the BBB. Given the sampled population frequencies of the commonest pgp haplotypes, and the fact that at clinically relevant doses a single functional copy of *abcb1a* is sufficient to prevent avermectin neurotoxicity in the CF-1 mouse, it is possible to calculate the proportion of the human populations that are likely to exhibit normal pgp BBB functionality (see publication for more details). >98% of people in African American and Caucasian populations will carry at least one copy of an ABCB1 haplotype that is already known to encode a pgp that is functional in the BBB and will therefore not be at risk of toxicity from the concentrations of avermectins to which humans are typically exposed. Between 1 and 2% of the population would thus carry only haplotypes with unconfirmed BBB functionality. Each individual “unconfirmed BBB functionality” haplotype is relatively rare within the population, often only having been identified in a single heterozygous individual, with each “rare” haplotype having an allelic frequency of less than 1%. As such, individuals that are homozygous for any one of the haplotypes with unconfirmed BBB functionality would be very rare within the population (<0.01%). If any of these rare haplotypes exhibited significantly compromised BBB pgp functionality it is likely that individuals homozygous for that haplotype, and thus having compromised BBB pgp function, would be extremely rare.

#### *Conclusions by MacDonald & Gledhill*

Pgp dependent xenobiotic efflux in the blood brain barrier and placental mother/fetus barrier play an important role in attenuating the known neurotoxicity of avermectins and the developmental toxicity of ivermectin and abamectin. There is currently no evidence for the existence of mutations of the ABCB1 gene in the human population that result in a loss

of function analogous to that seen in the CF-1 mouse and collie dog. Although there are numerous reports for and against the proposition that some ABCB1 SNPs and/or haplotypes exhibit reduced pgp expression and function, there are no consistent data indicating that known SNPs or haplotypes have an adverse effect on pgp function in the BBB or placenta. Where human BBB pgp function has been measured directly the most common haplotypes were found to have equal functionality. Since heterozygous pgp +/- mice and dogs do not exhibit ivermectin neurotoxicity at clinically relevant doses it is likely that humans carrying at least one functional copy of ABCB1 will not be more susceptible to avermectin toxicity at clinically relevant doses or at the low exposure levels resulting from pesticide use. Calculations using allelic frequencies of known haplotypes indicate that homozygosity for any as yet uncharacterised haplotypes with severely reduced BBB functionality is likely to be very rare in human populations.

### **Concluding remarks**

Based on the recent publication which reviews the relevant literature on p-glycoprotein polymorphism, showing that non-functional p-glycoprotein has not been identified in humans, and the supplementary studies in the DAR which show that only the -- CF-1 mouse is more sensitive to abamectin toxicity, it can be concluded that the studies with the unique polymorphic CF-1 mouse are not relevant for human risk assessment.

Also the JMPR meeting (JMPR, 1997) concluded that the CF-1 strain mouse is not appropriate for establishing an ADI for abamectin. The EU Committee for Veterinary Medicinal Products (CVMP) reached the same conclusion in 2002 upon reviewing the abamectin toxicity data and the JMPR position.

### **5.11 Derivation of DNEL(s) or other quantitative or qualitative measure for dose response**

*Not relevant for this type of report.*

**6 HUMAN HEALTH HAZARD ASSESSMENT OF PHYSICO-CHEMICAL PROPERTIES**

**6.1 Explosivity**

Abamectin has no explosive properties. No classification is required.

**6.2 Flammability**

Abamectin is considered not highly flammable. No classification is required.

**6.3 Oxidising potential**

Abamectin has no oxidising properties. No classification is required.

## 7 ENVIRONMENTAL HAZARD ASSESSMENT

The environmental hazard properties assessment for abamectin is based on the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands) and the Competent Authority Report (CAR; July 2008; RMS The Netherlands) on the inclusion of abamectin in Annex I to Directive 98/8/EC concerning the placing biocidal products on the market.

All tables in the present assessment are copied from the DAR or CAR. The tables are renumbered in accordance with the paragraph numbers in chapter 7.

### 7.1 Aquatic compartment (including sediment)

#### 7.1.1 Toxicity test results

##### 7.1.1.1 Fish

###### Short- and long-term toxicity to fresh water fish

The acute and chronic toxicity of abamectin and avermectin B<sub>1a</sub> to fresh water fish is summarised in the table below. Note that effect concentrations are given in µg/L. The toxicity of avermectin B<sub>1a</sub> to fish is similar to that of abamectin.

Table 7.1-1 Acute and chronic toxicity of abamectin and avermectin B<sub>1a</sub> to fish

Guideline / Test method	Species	Endpoint / Type of test	Exposure		Results [µg/L]		Remarks	Reference
			design	duration	L/EC <sub>50</sub>	NOEC		
EPA 1975	<i>Oncorhynchus mykiss</i>	LC <sub>50</sub> mortality	static	96 h	3.6		abamectin; nominal	LeBlanc & Sousa, 1981
OECD 203	<i>Oncorhynchus mykiss</i>	LC <sub>50</sub> mortality	flow-through <sup>a</sup>	96 h	8.7		abamectin; nominal	Peither, 2003
OECD 203	<i>Cyprinus carpio</i>	LC <sub>50</sub> mortality	flow-through	96 h	42		abamectin; nominal	Douglas & Pell, 1985
OECD 203	<i>Ictalurus punctatus</i>	LC <sub>50</sub> mortality	static	96 h	24		abamectin; nominal	McAllister et al., 1985
OECD 203	<i>Pimephales promelas</i>	LC <sub>50</sub> mortality	flow-through	96 h	14.7		abamectin; actual	Bätscher, 2003
EPA 1975	<i>Lepomis macrochirus</i>	LC <sub>50</sub> mortality	flow-through	96 h	7.2		avermectin B <sub>1a</sub> ; nominal	Forbis, 1983
OECD 204; OECD 215 (draft)	<i>Cyprinus carpio</i>	NOEC mortality; weight	flow-through	28 d		6.1	abamectin; actual	Ruffli, 2000
ASTM 1983	<i>Oncorhynchus mykiss</i>	NOEC Early Life Stage	flow-through	72 d		0.52	abamectin; actual	McAllister, 1986

a: modified exposure: gradually diminishing concentrations

##### 7.1.1.2 Aquatic invertebrates

###### Short- and long-term toxicity to fresh water invertebrates

The acute and chronic toxicity of abamectin and avermectin B<sub>1a</sub> to freshwater invertebrates is summarised in the table below. Note that effect concentrations are given in µg/L. The toxicity of avermectin B<sub>1a</sub> to daphnids is similar to that of abamectin.

Table 7.1-2 Acute and chronic toxicity of abamectin and avermectin B<sub>1a</sub> to invertebrates

Guideline /	Species	Endpoint /	Exposure	Results [µg/L]	Remarks	Reference
-------------	---------	------------	----------	----------------	---------	-----------

ANNEX VI REPORT – HARMONISATION OF C&L FORMAT

			design	duration	L/EC <sub>50</sub>	NOEC		
EPA 1975	<i>Daphnia magna</i>	LC <sub>50</sub> mortality	static	48 h	0.34		abamectin; nominal	LeBlanc & Surprenant, 1981
EPA	<i>Daphnia magna</i>	EC <sub>50</sub> immobilisation	static	48 h	0.37		abamectin; actual	Forbis, 1989a
EPA	<i>Daphnia magna</i>	EC <sub>50</sub> immobilisation	static; sediment spiked	48 h	0.26		abamectin; actual water	Forbis, 1989b
EPA; OECD 202	<i>Daphnia magna</i>	EC <sub>50</sub> immobilisation	static	48 h	0.56		abamectin; actual initial	Rufli, 1998
EPA	<i>Daphnia magna</i>	EC <sub>50</sub> immobilisation	static	48 h	0.3		abamectin; nominal	Naimie et al., 1985
EPA	<i>Daphnia magna</i>	EC <sub>50</sub> immobilisation	static	48 h	0.63		avermectin B <sub>1a</sub> ; nominal	Naimie et al., 1985
EPA; OECD 202	<i>Daphnia longispina</i>	EC <sub>50</sub> immobilisation	static	48 h	0.38		abamectin; actual	Knauer, 2001b
EPA; OECD 202	<i>Daphnia pulex</i>	EC <sub>50</sub> immobilisation	static	48 h	0.12		abamectin; actual	Knauer, 2001b
EPA; OECD 202	<i>Daphnia pulex</i>	EC <sub>50</sub> immobilisation	static	48 h	0.28		abamectin; nominal	Knauer, 2001c
EPA; OECD 202	<i>Daphnia galeata</i>	EC <sub>50</sub> immobilisation	static	48 h	0.55		abamectin; nominal	Knauer, 2001a
EPA; OECD 202	<i>Simocephalus sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	0.30		abamectin; actual	Knauer, 2001b
EPA; OECD 202	<i>Diaphanosoma sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	0.53		abamectin; nominal	Knauer, 2001c
EPA; OECD 202	<i>Thamnocephalus platyurus</i>	EC <sub>50</sub> immobilisation	static	48 h	30		abamectin; nominal	Knauer, 2001d
EPA; OECD 202	<i>Thamnocephalus platyurus</i>	EC <sub>50</sub> immobilisation	static	48 h	2.8		abamectin; actual	Knauer, 2001e
EPA; OECD 202	<i>Brachionus calyciflorus</i>	EC <sub>50</sub> immobilisation	static	48 h	4000		abamectin; actual	Knauer, 2001e
EPA; OECD 202	<i>Chaoborus sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	190		abamectin; actual	Knauer, 2001f
EPA; OECD 202	<i>Chaoborus sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	41		abamectin; nominal	Knauer, 2001g
EPA; OECD 202	<i>Cloeon sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	2.9		abamectin; nominal	Knauer, 2001g
EPA; OECD 202	<i>Gammarus sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	6.2		abamectin; nominal	Knauer, 2001h
EPA; OECD 202	<i>Gammarus sp.</i>	EC <sub>50</sub> immobilisation	static	48 h	8.6		abamectin; actual	Knauer, 2001i
EPA; OECD 202	<i>Lymnea stagnalis</i>	EC <sub>50</sub> immobilisation	static	48 h	55		abamectin; actual	Knauer, 2001j
OECD 211	<i>Daphnia magna</i>	NOEC mortality	semi-static	21 d		0.010	abamectin; nominal	Pfeifle, 2001a
OECD 211	<i>Daphnia magna</i>	NOEC mortality	flow-through	21 d		0.030	avermectin B <sub>1a</sub> ; actual	Surprenant & Mastone, 1983

### 7.1.1.3 Algae and aquatic plants

No data available

### 7.1.1.4 Sediment organisms

The toxicity of abamectine to *Chironomus riparius* is summarised in table 7.1-3.

Table 7.1-3 Chronic toxicity to sediment organisms

Guideline / Test method	Species	Endpoint / Type of test	Exposure		Results NOEC	Remarks	Reference
			design	duration			
draft BBA draft OECD	<i>Chironomus riparius</i>	NOEC emergence	static; sediment	28 d	3.3 µg/kg dw	avermectin B <sub>1a</sub> ; nominal initial	Grade, 2002

### 7.1.1.5 Other aquatic organisms

Accepted data for marine fish and invertebrates are summarised in Table 7.1-4. The large difference between the static LC<sub>50</sub> of 0.21 µg/L for the saltwater species *Mysidopsis bahia* and the results of the flow-trough experiments (LC<sub>50</sub> 0.020 and 0.022 µg/L) may be due to the fact that the first value is based on nominal concentrations.

Table 7.1-4 Acute and chronic toxicity to saltwater organisms

Guideline / Test method	Species	Endpoint / Type of test	Exposure		Results [µg/L]		Remarks	Reference
			design	duration	L/EC <sub>50</sub>	NOEC		
ASTM 1982	<i>Cyprinodon variegatus</i>	LC <sub>50</sub> mortality	static	96 h	15		abamectin; nominal	Ward, 1985
EPA 1970, APHA 1980	<i>Mysidopsis bahia</i>	LC <sub>50</sub> mortality	static	96 h	0.21		abamectin ; nominal	Forbis & Burgess, 1985
EPA 1970, APHA 1980	<i>Mysidopsis bahia</i>	LC <sub>50</sub> mortality	flow-through	96 h	0.022		abamectin ; actual	Suprenant, 1988
EPA 1970, APHA 1980	<i>Mysidopsis bahia</i>	LC <sub>50</sub> mortality	flow-through	96 h	0.020		abamectin ; actual	Suprenant, 1988
BMRL	<i>Crassostrea virginica</i>	EC <sub>50</sub> larval development	static	48 h	430		abamectin ; nominal	Ward, 1983
BMRL	<i>Penaeus duorarum</i>	LC <sub>50</sub> mortality	static	96 h	1.6		abamectin ; nominal	Ward, 1983
BMRL	<i>Callinectes spidus</i>	LC <sub>50</sub> mortality	static	96 h	153		abamectin ; nominal	Ward, 1983
	<i>Mysidopsis bahia</i>	NOEC reproduction	flow-through	28 d		0.0035	abamectin ; actual	Suprenant, 1988

### 7.1.2 Calculation of Predicted No Effect Concentration (PNEC)

Not relevant for this type of report.

## 7.2 Terrestrial compartment

### 7.2.1 Toxicity test results

#### 7.2.1.1 Toxicity to soil macro organisms

Not applicable for this type of report.

#### 7.2.1.2 Toxicity to terrestrial plants

Not applicable for this type of report.

#### 7.2.1.3 Toxicity to soil micro-organisms

Not applicable for this type of report.

#### 7.2.1.4 Toxicity to other terrestrial organisms

Not applicable for this type of report.

### 7.2.2 Calculation of Predicted No Effect Concentration (PNEC<sub>soil</sub>)

Not relevant for this type of report.



### **7.3 Atmospheric compartment**

No data available

### **7.4 Microbiological activity in sewage treatment systems**

#### **7.4.1 Toxicity to aquatic micro-organisms**

Not applicable for this type of report.

#### **7.4.2 PNEC for sewage treatment plant**

Not relevant for this type of report.

### **7.5 Calculation of Predicted No Effect Concentration for secondary poisoning (PNEC<sub>oral</sub>)**

Not relevant for this type of report.

### **7.6 Conclusion on the environmental classification and labelling**

Abamectine is hydrolytically stable. Abamectin is not readily biodegradable as only 3% degradation was observed in an OECD301F test after 28 days. Abamectin is not readily degradable as the DT<sub>50,system</sub> obtained in aerobic and anaerobic simulation studies in water/sediment systems was 87-91 days and 230-312 days, respectively.

Abamectin has a log K<sub>ow</sub> of 4.4. In a BCF study, a BCF value of 69 was obtained based on plateau total radioactive residue in whole fish and average total radioactive residue in water, whereas a BCF value of 52 was obtained based on uptake and elimination rate constants.

Abamectin generally produces LC<sub>50</sub> and EC<sub>50</sub> values in the µg/l range in fish and crustaceans. The lowest LC<sub>50</sub> value obtained for abamectin in freshwater fish, freshwater crustaceans and marine crustaceans is 3.6 µg/l, 0.12 µg/l and 0.020 µg/l, respectively. The available EC<sub>50</sub> values ranged from 0.0035 µg/l to 6.1 µg/l.

#### Conclusion of environmental classification according to Directive 67/548/EEC

In acute aquatic toxicity studies, L(E)C<sub>50</sub> values in fish and crustaceans were obtained at abamectine concentrations <1 mg/l. Abamectin is not readily biodegradable. Abamectin therefore fulfills the criteria for classification with N; R50/53.

#### Conclusion of environmental classification according to Regulation EC 1272/2008

In acute aquatic toxicity studies, L(E)C<sub>50</sub> values in fish and crustaceans were obtained at abamectin concentrations <1 mg/l. Abamectin is not rapidly degradable based on 3% degradation in a ready biodegradability study, and a DT<sub>50,system</sub> of 87-91 days and 230-312 days in aerobic and anaerobic water/sediment simulation studies. Abamectin therefore fulfills the criteria for classification as aquatic environmental hazard acute category 1, H400 and aquatic environmental hazard chronic category 1, H410.

The M-factor for abamectin is 10,000. This value is based on two LC50 values of 0.020 µg/l and 0.022 µg/l obtained for the marine crustacean *Mysidopsis bahia* in a 96-h flow-through study.

## **JUSTIFICATION THAT ACTION IS REQUIRED ON A COMMUNITY-WIDE BASIS**

Harmonised classification is required because this substance is an active substance in the meaning of Directive 91/414/EEC and Directive 98/8/EC (Regulation EC 1272/2008 article 38: 1a)

## REFERENCES

The classification and labeling proposal is based on the data provided in:

- the Draft Assessment Report and Proposed Decision of the Netherlands prepared in the context of the possible inclusion of abamectin in Annex I of Council Directive 91/414/EEC (DAR October 2005 + addendum February 2008, RMS The Netherlands)
- the Competent Authority Report (CAR; July 2008; RMS The Netherlands) on the inclusion of abamectin in Annex I to Directive 98/8/EC concerning the placing biocidal products on the market.
- JECFA, 1991, Toxicological evaluation of certain veterinary drug residues in food, IPCS, WHO Food additive series 27, Geneva, 1991.