



**Recommendation from the Scientific  
Committee on Occupational Exposure Limits  
for amitrole**  
SCOEL/SUM/157  
February 2009





## Table of Contents

1. Occurrence/use and occupational exposure .....	4
2. Health significance.....	4
2.1 Toxicokinetics.....	4
2.1.1 Human data .....	4
2.1.2. Animal data .....	4
2.1.3. Biological monitoring.....	5
2.2. Acute toxicity .....	5
2.2.1. Human data .....	5
2.2.2. Animal data .....	5
2.3. Irritation and corrosivity.....	5
2.3.1. Human data .....	5
2.3.2. Animal data.....	5
2.4. Sensitisation.....	6
2.4.1. Human data .....	6
2.4.2. Animal data.....	6
2.5. Repeated dose toxicity .....	6
2.5.1. Human data .....	6
2.5.2. Animal data.....	6
2.6. Genotoxicity .....	8
2.6.1. In vitro.....	8
2.6.2. In vivo – Human data .....	8
2.6.3. In vivo – Animal data.....	8
2.7. Carcinogenicity .....	8
2.7.1. Human data .....	8
2.7.2. Animal data.....	8
2.8. Reproductive toxicity .....	10
2.8.1. Human data .....	10
2.8.2. Animal data.....	10
Recommendations.....	13
References.....	15




## Recommendation from the Scientific Committee on Occupational Exposure Limits for amitrole

Eight-hour TWA:	0.2 mg/m <sup>3</sup>
STEL (15 minutes):	–
Notation:	–
Biological Limit Value (BLV):	
SCOEL carcinogenic group:	D ("non-genotoxic carcinogen")

### Substance identification

Amitrole:



Synonyms: 1*H*-1,2,4-Triazol-3-amine; delta2-1,2,4-Triazoline 5-imino-; s-Triazole, 3-amino-; 3-Amino-1,2,4-triazole

EC No.: 200-521-5

Annex I Index No.: 613-011-00-6

EU Classification: Repr.Cat.3; R63 - Xn; R48/22 - N; R51-53

CAS No.: 61-82-5

MWt: 84.080

Conversion factor (20 °C, 101 kPa): 1 ppm = 3.50 mg/m<sup>3</sup>; 1 mg/m<sup>3</sup> = 0.286 ppm

This evaluation is based on ACGIH (2001), ECB (2000), Greim (1998, 1999, 2002), Henschler (1983), IARC (2001), WHO (1993, 1994a,b), FAO (1997) and the references cited in these reviews.

### Physico-chemical properties

Amitrole is a crystalline, colourless solid. The melting point of the substance is 157 -159 °C and the vapour pressure is  $5.79 \cdot 10^{-7}$  hPa at 20 °C. The water solubility of amitrole is 280 - 335 g/l at 20 - 25 °C and the log *P*<sub>ow</sub> is -0.969. The substance has a density of 0.6 - 0.85 g/cm<sup>3</sup> at 20 °C (ECB, 2000).



# 1. Occurrence/use and occupational exposure

Amitrole is used as a herbicide and as a growth regulator for plants (ACGIH, 2001; IARC, 2001).

## 2. Health significance

### 2.1 Toxicokinetics


#### 2.1.1 Human data

After intentional ingestion of approximately 20 mg/kg amitrole, a female subject excreted the unchanged compound at a concentration of about 1 g/l in urine (Geldmacher-von Mallingkrodt and Schmidt, 1970).

#### 2.1.2. Animal data

The fraction of amitrole absorbed by the rat lung was 50% (ECB, 2000). The fraction of oral absorption in rats was 70 - 95% (Fang et al., 1964, 1966; WHO, 1993; FAO, 1997; IARC, 2001). After dermal exposure of rabbits for 24 h, 70% remained at the application site and about 30% was absorbed (WHO, 1993). However, EC (2001) reported a dermal absorption rate of only 1% (no further details). After oral or intravenous administration to mice, amitrole was found first in the bone marrow, spleen, thymus and gastrointestinal tract. After 3 days, amitrole was identified only in the livers of rats and mice (less than 3% of the administered dose). Little metabolic transformation of amitrole occurs in mammals. The two main urinary metabolites in rats (3-amino-5-mercapto-1,2,4-triazole and 3-amino-1,2,4-triazolyl-(5)-mercapturic acid) represented less than 10% of the administered dose. When rats were exposed by inhalation to aqueous aerosols of amitrole (25.8 or 49.2 mg/m<sup>3</sup>, 1 h), urine was the major excretion route. About 75% of the eliminated material was found in the urine within 12 h. After oral administration, most of the compound was eliminated unchanged in urine within 24 h. Minor amounts were found in faeces and expired air. Following dermal exposure, rabbits excreted amitrole in urine and faeces in similar amounts (WHO, 1993; FAO 1997; IARC, 2001).

Figure 1. Metabolic pathway of amitrole in rats





### 2.1.3. Biological monitoring

There are no data available.

## 2.2. Acute toxicity

### 2.2.1. Human data

A single case report described the development of a severe alveolar damage after a single 2 h exposure to a spray containing 19% amitrole, 17% ammonium thiocyanate, less than 1% sodium diethylsulfosuccinate and less than 1% ethylene oxide (Balkisson et al., 1992).

No toxic symptoms appeared following the intentional ingestion of 20 mg/kg amitrole by a female subject (Geldmacher-von Mallinkrodt and Schmidt, 1970).

Astwood (1960) reported that a single oral dose of 100 mg amitrole inhibited radioiodine uptake by the thyroid for 24 hours in both normal subjects and subjects with hyperthyroidism. A dose of 10 mg (0.15 - 0.2 mg/kg, Henschler, 1983) had only a slight effect on iodine uptake.

### 2.2.2. Animal data

The 4 h-inhalation  $LC_{50}$  in rats was  $> 439 \text{ mg/m}^3$ . The oral  $LD_{50}$  was  $> 4000 - 25000 \text{ mg/kg}$  in rats and  $11000 - 14700 \text{ mg/kg}$  in mice. The dermal  $LD_{50}$  values in rats and rabbits were  $> 2500 \text{ mg/kg}$  and  $> 10000 \text{ mg/kg}$ , respectively (Henschler, 1983; WHO, 1993, 1994b).

Signs of toxicity at high doses included depression, dyspnoea, diarrhoea, ataxia, convulsions, coma and death. Macroscopic findings were irritation and haemorrhage of the gastrointestinal tract (WHO, 1993). The presence of sodium carbonate, sodium bicarbonate and wetting agents considerably increased the toxicity of amitrole (WHO, 1993).

## 2.3. Irritation and corrosivity

### 2.3.1. Human data

Amitrole was not irritating in a patch test with a human volunteer exposed for 4 or 8 hours. A slight irritating effect was observed in 3 out of 6 subjects after 24 hours of exposure (Hecht, 1954).

Amitrole-exposed workers in different plants occasionally developed mild dermatitis, which was interpreted as a primary irritant effect (WHO, 1993).

### 2.3.2. Animal data

#### Skin

Dermal exposure of rabbits to  $10000 \text{ mg/kg}$  amitrole or rats to  $2500 \text{ mg/kg}$  produced mild and reversible erythema (Elsea, 1954; IPCS, 1994).

#### Eyes

A single 4 h exposure by inhalation to  $439 \text{ mg/m}^3$  did not produce eye irritation in rats (ACGIH, 2001). The application of 3 mg into the conjunctival sac of rabbits caused mild eye irritation (Elsea, 1954; IPCS, 1994).

#### Respiratory tract

A single 4 h exposure by inhalation to  $439 \text{ mg/m}^3$  did not irritate the respiratory tract of rats (ACGIH, 2001).



## 2.4. Sensitisation

### 2.4.1. Human data

A case study of a weed control operator with contact dermatitis was reported. Patch testing with 1% amitrole resulted in a strong positive vesicular reaction on days 2 and 4, indicating an allergic contact dermatitis (English et al., 1986).

### 2.4.2. Animal data

A Magnusson-Kligman maximisation test with Freund's adjuvant on guinea pigs yielded positive results after intradermal induction with 2.5% amitrole, dermal induction with 25% and 2 challenges with 12% (Bayer AG, 1984). No sensitisation was observed in a Klecak open epicutaneous test on guinea pigs, exposed 20-times dermally to 3 - 30% amitrole and challenged twice with 1 - 30% amitrole (Bayer AG, 1985).

## 2.5. Repeated dose toxicity

### 2.5.1. Human data

No medical findings except skin irritation were reported in workers in different plants with chronic amitrole exposure (Greim, 2001; WHO, 1994a). No effects on thyroid function (up to 14 days after the end of exposure) were observed in men who had sprayed amitrole for 10 days. Their exposure was estimated to about 340 mg/d (5 mg/kg • d) (Baughter et al., 1982). In a study by Bayer AG (1983), thyroid function was examined in 5 employees exposed to unknown level of amitrole who had worked for prolonged periods in the production and packaging of the substance. There were no indications of thyroid dysfunction.

### 2.5.2. Animal data

Rodents, especially rats, are more prone to thyroid effects than dogs or humans (see section "recommendations"). These effects are reversible after cessation of exposure (Greim, 1983).

#### Inhalation

No published studies are available.

In an unpublished study by Cox and Re (1978), F-344 rats (15 per sex and dose) were exposed to amitrole aerosols at concentrations of 0, 100, 320, 990 and 4050 mg/m<sup>3</sup> on 5 h/d, 5 d/w for 4 weeks. At 320 mg/m<sup>3</sup> and above, hyperplasia of the thyroid was evident and triiodothyronine (T<sub>3</sub>) levels were significantly depressed. Thyroxine (T<sub>4</sub>) levels were significantly depressed at 990 and 4050 mg/m<sup>3</sup>. The NOAEC of this study was 100 mg/m<sup>3</sup>.

Fischer rats (75 per sex and dose) were exposed by inhalation to 0, 15.8 - 32.2 and 97.9 - 376.4 mg/m<sup>3</sup> (ranges for the two dose groups) on 5 h/d, 5 d/w on study weeks 1 - 13, 40 - 52 and 78 - 90, interrupted by recovery periods (Becci, 1983 cited in WHO). The exposure of the high-dose animals was terminated early at week 51 (high mortality due to technical problems, but not substance-related). The most prominent findings of exposure were alterations of the thyroid in both dose groups, including reduced T<sub>3</sub> and T<sub>4</sub> levels, increased organ weights and follicular hyperplasia. At the terminal sacrifice (after 24 months) there was also an increased incidence of thyroid tumours.

ECB (2000) refers to a rat inhalation study with duration of 2 years, in which the animals were exposed once a week (duration not stated) to 2000 mg/m<sup>3</sup> amitrole aerosol. There were no effects on mortality, body weights and tumour incidences. No histological alterations in liver and thyroid were observed (no further details).



### Oral

Numerous studies examined the effects of short-term and long-term oral exposure to rats and mice (overviews in ECB, 2000; Greim, 1983; IARC, 2001; WHO, 1993, 1994a). Only the most relevant studies concerning the effects on the thyroid are described here.

Fregly (1968) performed a study on rats (Blue Spruce Farm strain, 10 males per dose) in order to establish the minimum dose affecting thyroid activity. Amitrole was administered in the diet at concentrations of 0, 0.25, 0.5, 2, 10 or 50 mg/kg for 11 - 13 weeks. At 2 mg/kg feed (0.1 mg/kg/d) and above, the uptake of iodine by the thyroid was reduced in a dose-related manner. A decrease in the serum protein-bound iodine level was also observed, but was not dose-related. Histological changes in the thyroid were noted at 10 and 50 mg/kg feed. The NOAEL of this study is 0.5 mg/kg feed (0.025 mg/kg/d).

This above NOAEL was corroborated by an unpublished rat study, which revealed a NOAEL of 0.5 mg/kg feed (0.025 mg/kg • d) and a threshold of 1 mg/kg feed (0.05 mg/kg/d) for decreased iodine uptake, reduced levels of T<sub>3</sub> and T<sub>4</sub> and increased thyroid weight (Henschler, 1983). Further studies have confirmed the antithyroid effect of amitrole at higher doses. EC (2001) reported a NOAEL of 0.1 mg/kg/d for thyroid effects in a 90-day rat study, but no further details are given.

Thyroid effects were also seen in rat studies on reproduction, but at higher doses (see section "reproductive toxicity").

Beagle dogs (4 per sex and group) were exposed for one year to amitrole in the diet at concentrations of 0, 10, 500 and 1500 mg/kg (corresponding to doses of 0.3, 13 and 32 mg/kg/d) (Bayer AG, 1994). The predominant finding was an effect on the thyroid starting at 500 mg/kg feed (enlargement, ectopic tissue, slight hypertrophy, follicular hyperplasia, decreased levels of T<sub>3</sub> and T<sub>4</sub>). Thrombosis, pigmentation, and haemorrhage were observed in the thyroids at the highest dose. Haematological alterations typical of hyperthyroidism included lower erythrocyte, haemoglobin and haematocrit counts as well as slightly lower mean cell volume and mean cell haemoglobin. Pituitary hyperplasia (in males at 1500 mg/kg) and hypertrophy (in males at 500 mg/kg feed and in both sexes at 1500 mg/kg feed) were also evident. Males at the high dose group showed a slight decrease in food consumption and body weight as well as an increase in cholesterol levels and platelet counts. No effects were observed at 10 mg/kg feed (0.3 mg/kg/d).

Rabbits developed cataracts after oral exposure to amitrole (0.2% in the diet or 0.2% in drinking water) for up to 25 weeks (Bhuyan et al., 1973).

The thyroid effects in subchronic rat studies were reversible within 2-4 weeks after cessation of exposure (Greim, 1981).

### Dermal

Toxicity after dermal exposure was only observed at high doses.

No systemic toxicity was observed after dermal exposure of rats to 2.39 mg/kg/d once a week for 23 months (ECB, 2000).

No local or systemic effects were observed following dermal exposure of rabbits to doses of 0, 25 or 100 mg/kg/d for 6 h/d for 15 days (six animals per group, three animals per group tested with abraded skin (Mihail and Schilde, 1984).

EC (2001) reported a NOAEL of 100 mg/kg/d for a 28-day dermal exposure study in rats, but further details are not given.

Local effects and systemic maternal toxicity was observed in a dermal teratogenicity study on rabbits. The LOAEL was 2000 mg/kg/d in rabbits, the NOAEL 1500 mg/kg/d (see section "reproductive toxicology").



## 2.6. Genotoxicity

### 2.6.1. In vitro

Most of the *in vitro* tests on the genotoxicity of amitrole – either with and without metabolic activation – yielded negative results, including tests in bacteria (*Salmonella typhimurium* of different strain, *Bacillus subtilis*, *E. coli*), yeasts and fungi (Greim, 1998; IARC, 2001; WHO, 1994a). Tests with positive responses were criticised due to the use of inappropriate methods (use of S9 from fish or mussels) or lack of documentation. However, it was stated that *in vitro* tests may be insensitive because an *in vivo* activation could occur by peroxidases, which are not present *in vitro* (Greim, 1998). No gene mutations were induced in L5187Y mouse lymphoma cells or human fibroblasts but in Syrian hamster embryo cells *in vitro* (HPRT- and Na/K ATPase locus). These cells are also sensitive to morphological transformation by amitrole. The prostaglandin-H-synthetase activity, present in Syrian hamster embryo cells, is suggested to be responsible for the sensitivity of these cells. No chromosomal aberrations were observed in Chinese hamster lung cells or human lymphocytes. One test on the induction of sister chromatid exchange in Chinese hamster lung cells yielded a positive response; but without a dose-response-relationship (Greim, 1998; IARC, 2001; WHO, 1994a).

### 2.6.2. In vivo – Human data

Human data on the genotoxic effects *in vivo* are not available.

### 2.6.3. In vivo – Animal data

In a host-mediated assay with *Salmonella typhimurium* TA1530 in mice, a positive, but not dose-related effect was observed. Amitrole did not induce dominant lethal mutations or micronuclei in the bone marrow of mice (oral or intraperitoneal exposure). Unscheduled DNA synthesis was not observed in rat hepatocytes *in vivo* (Greim, 1998; IARC, 2001; WHO, 1994a). It did not induce mutations in *Drosophila* after oral exposure (Pontecorvo and Fantaccione, 2005).

## 2.7. Carcinogenicity

### 2.7.1. Human data

Axelsson and Sundell (1974) and Axelsson et al. (1980) performed a cohort study on 348 railroad workers with exposure to amitrole, other herbicides and chemicals. In a sub-cohort exposed to amitrole in combination with chlorophenoxy herbicides and other chemicals, there was a significant increase in cancer deaths. However, in a sub-cohort mainly exposed to amitrole alone, no significant increase in tumours was reported.

### 2.7.2. Animal data

In most of the available long-term inhalation and oral studies on mice and rats, amitrole induced tumours of the thyroid (follicular-cell adenomas and carcinomas) at doses of about 2.5 mg/kg/d and above. This effect was attributed to a specific sensitivity of rodents (see “recommendations”) (Greim, 1998; IARC, 2001; WHO, 1993, 1994a).

Other target organs were the liver and pituitary. In the study by Napalkov (1962), Albino rats were exposed for their whole life to amitrole via drinking water (resulting in a dose of 20 - 25 mg/animal per day) or feed (doses of 250 and 500 mg/animal per day). The incidence of liver tumours in surviving animals at the time of appearance of the first tumour were 6/8 (drinking water study), 8/10 and 10/11 (250 and 500 mg group in the feed study, respectively), for thyroid tumours 3/8, 2/10 and 5/11. This study lacked control groups.

Steinhoff et al. (1983) fed Wistar rats, NMRI mice and Golden hamsters with a diet containing 0, 1, 10 and 100 mg/kg feed for their whole lifespan. They were estimated induction of tumors and thyroid function test (in 5 animals for each species): thyroid





weights, percentage of accumulation of radioiodine and protein-bound radioiodine. Increased incidences of malignant thyroid tumors and adenomas of the pituitary were observed in female rats of the high dose group (36/41 vs. 14/59 in controls). No carcinogenic effects were seen in mice or hamsters, but the administered doses were low. The dose below which there was neither long-lasting interference with thyroid function nor induction of tumors (NOAEL) for female rats was 10 mg/kg feed equivalent to 1.5 mg/kg bw/day.

Table 1. Numbers of animals with tumor and distribution of tumors observed in liver, pituitary, and thyroid gland (Steinhoff et al., 1983)

Species	Dose mg/kg	Sex	Animals with tumor	Animals with malignant tumor	Primary tumors of					
					Liver cells		Pituitary		Thyroid	
					Benign	Malignant	Benign	Malignant	Benign	Malignant
Hamster (76/sex/group)	0	M	20	7	-	-	1	-	2	-
		F	13	1	1	-	-	-	1	-
	1	M	11	2	-	-	-	-	2	-
		F	10	2	1	-	-	-	2	-
	10	M	13	5	-	-	-	-	-	-
		F	12	3	-	-	-	1	-	-
	100	M	11	2	-	-	-	-	1	-
		F	11	3	-	-	-	-	-	-
Mouse (75/sex/group)	0	M	52	27	4	1	1	-	-	-
		F	60	46	-	-	10	-	-	-
	1	M	56	38	2	-	-	-	-	-
		F	48	26	-	-	4	-	-	-
	10	M	47	26	3	1	2	-	-	-
		F	59	46	-	-	6	-	-	-
	100	M	46	32	2	1	-	-	-	-
		F	55	38	-	-	10	-	-	-
Rat (75/sex/group)	0	M	36	19	-	-	4	-	5	3
		F	59	20	-	-	14	1	7	-
	1	M	41	20	-	-	9	1	9	-
		F	67	34	-	-	20	2	12	1
	10	M	44	23	-	1	10	-	4	3
		F	60	29	-	-	15	4	8	4
	100	M	53	23	-	-	10	3	45	18
		F	71	45	-	-	36	5	44	28

B6C3F1 mice were exposed by Vesselinovitch (1983) to amitrole in the diet at concentrations of 500 mg/kg feed from weaning for 90 weeks. The tumour incidences in the liver were significantly increased. There was one hepatocellular adenoma in 98 control males, 9 adenomas and 11 carcinomas in 55 exposed males as well as 5 adenomas and 4



carcinomas in 49 exposed females. Other studies on mice confirmed these findings, but the mechanism of the formation of liver tumours is non-genotoxic (Greim, 1998; IARC, 2001).

No increased tumour incidences were observed after dermal exposure of rats to low doses of 2.39 mg/kg/d once a week for 23 months (ECB, 2000).

The International Agency for Research on Cancer (IARC) in 1987 classified amitrole as Group 2B (*possibly carcinogenic to humans*). The Working Group reported that amitrole had been tested for carcinogenicity in mice by oral administration, skin application and transplacental exposure, in rats by oral and subcutaneous administration and in hamsters by oral administration. After oral administration, it produced thyroid tumors and benign and malignant liver tumors in mice of each sex, benign and malignant thyroid tumors in male and female rats and benign pituitary tumors in female rats. There was *sufficient evidence* in experimental animals for the carcinogenicity of amitrole. No data were available on the genetic and related effects of amitrole in humans.

The IARC Working Group in 2001 changed the classification and concluded that amitrole is *not classifiable as to its carcinogenicity to humans* (Group 3). In making its evaluation, the Working Group concluded that amitrole produces thyroid tumors in mice and rats by a non-genotoxic mechanism, which involves interference with the functioning of thyroid peroxidase, resulting in a reduction in circulating thyroid hormone concentrations and increased secretion of thyroid-stimulating hormone. Consequently, amitrole would not be expected to produce thyroid cancer in humans exposed to concentrations that do not alter thyroid hormone homeostasis. An additional consideration of the Working Group, based on the lack of genotoxicity of amitrole, was that the liver tumors in mice and benign pituitary tumors in rats were also produced by a non-genotoxic mechanism. Epidemiological studies and toxicological studies in experimental animals provide compelling evidence that rodents are substantially more sensitive than humans to the development of thyroid tumors in response to thyroid hormone imbalance.

## 2.8. Reproductive toxicity

### 2.8.1. Human data

Human data on reproductive or developmental effects are not available.

### 2.8.2. Animal data

#### *Fertility*

Rats (Sherman strain) were exposed to amitrole in a one-generation study by Gaines et al. (1973) at concentrations of 0, 500 and 1000 mg/kg feed (for F<sub>0</sub> females: 31 - 43 mg/kg/d and 68 - 86 mg/kg/d; for F<sub>0</sub> males: 28 - 42 mg/kg/d and 60 - 87 mg/kg/d) from 55 days before mating until weaning of the F<sub>1</sub> generation. There was no effect on fertility, but the body weights of the exposed F<sub>0</sub> and F<sub>1</sub> animals were reduced and spleen and thymus were atrophic. A marked increase in offspring mortality was observed within the first week after weaning.

Gaines et al. (1973) performed a subsequent two-generation study with 0, 25 and 100 mg/kg feed (for F<sub>0</sub> females: 1.3 - 2.5 mg/kg/d and 4.8 - 9.6 mg/kg/d; for F<sub>0</sub> males: 1.1 - 2.5 mg/kg/d and 4.2 - 9.5 mg/kg/d). There were no effects on reproduction or development, but hyperplasia of the thyroid was observed in all treated groups of both studies.

More recent studies on reproductive toxicity by Savary (1994) and Richard (1995) are available (unpublished, but extensively documented in FAO, 1997).

In a one-generation study by Savary (1994), Sprague-Dawley rats were exposed to amitrole in the diet at levels of 0, 2, 10, 40 or 160 mg/kg (F<sub>0</sub> females: 0.29, 1.4, 5.3 and 28 mg/kg/d; F<sub>0</sub> males: 0.19, 0.94, 3.7 and 12 mg/kg/d) for 29 days before mating until



weaning of the F<sub>1</sub> pups. At the highest dose, clinical signs of maternal toxicity were evident and the mortality of F<sub>1</sub> pups was increased in this group. Decreased body weights and decreased feed consumption were seen at 160 mg/kg in F<sub>0</sub> and F<sub>1</sub> animals as well as in F<sub>1</sub> males at 40 mg/kg. Females at the highest dose had a lower implantation rate (12.1 implantation sites) than controls (16.6 sites), which was reflected in significantly lower birth rates and litter sizes. The viability indices were not affected by the treatment. The number of females with a normal oestrus cycle was decreased in a dose-related fashion in animals at 40 or 160 mg/kg. At doses of  $\geq 10$  mg/kg in males and at  $\geq 40$  mg/kg in females, the thyroid glands of F<sub>0</sub> animals showed a dose-related increase in size and reddish colouration. All animals exposed to 160 mg/kg also revealed other organ lesions (reddened pituitary glands, greyish-white foci on the lungs, reduced adrenal glands and spleen sizes). No treatment-related histopathological lesions were seen in pups killed on day 14 post partum. Male and female F<sub>1</sub> adults (40 mg/kg or above) had enlarged, reddened thyroid glands and reddened pituitary glands. Histopathological examination showed dose-related effects in the liver and pituitary gland in F<sub>0</sub> and F<sub>1</sub> animals, namely decreased colloid content in females of both generations at 10 mg/kg and above, vascular ectasia in all animals of both generations treated with 40 or 160 mg/kg, and periadenitis in a small number of F<sub>0</sub> males at 160 mg/kg.

In a subsequent two-generation study (Richard, 1995), groups of 30 male and 30 female Sprague-Dawley rats were fed diets containing amitrole at levels of 0, 0.5, 2, 15 or 110 mg/kg (F<sub>0</sub> females: 0.04, 0.16, 1.2 and 7.8 mg/kg/d; F<sub>0</sub> males: 0.03, 0.12, 0.9 and 5.9 mg/kg/d) from 72 days before mating throughout gestation and lactation for two generations (Richard, 1995). Increased mortality and/or clinical signs of toxicity were observed in F<sub>1</sub> males and females of the highest dose group. Both F<sub>0</sub> and F<sub>1</sub> parental animals at this dose had relevant decreases in feed consumption and body weight throughout the study, the F<sub>1</sub> generation being more affected. Animals of the 110 mg/kg group showed widespread systemic toxicity in the form of increased weights and histological alterations of the thyroid, pituitary and spleen, increased weights of testes, epididymis, seminal vesicles (both generations), prostate and uterus (only F<sub>1</sub>) and decreased relative ovarian weights (F<sub>0</sub>). Histopathological effects were also seen in reproductive organs (testis and epididymis of F<sub>1</sub> animals and in the ovaries, uterus, and vagina in both generations). Many of the histopathological effects on the reproductive tissues were considered to be related to immaturity of the animals at this dose. Other affected organs in animals of the highest dose groups were the adrenals (decreased weight, histopathological alterations), kidneys (increased weight, renal lesions) and liver (hepatocellular hypertrophy). Owing to a low survival rate of F<sub>2</sub> pups in this group, treatment-related organ changes could not be identified. Significantly decreased mating indices and decreased fertility indices were evident among F<sub>1</sub> males and females, as well as increased length of gestation in F<sub>1</sub> females and a low implantation rate in F<sub>0</sub> and F<sub>1</sub> females of the highest dose. In the offspring there was a low prenatal survival in the F<sub>2</sub> generation, significantly decreased mean litter size at day 1 (F<sub>1</sub> and F<sub>2</sub>) and day 4 or 21 (F<sub>2</sub> only) post partum, very low viability indices for the F<sub>2</sub> generation and decreased F<sub>1</sub> and F<sub>2</sub> pup body weights during lactation. In the F<sub>2</sub> generation at 110 mg/kg, there was a high postnatal mortality. There appeared to be no effects on the development of the F<sub>1</sub> pups and the F<sub>2</sub> generation could not be examined due to poor survival. At 15 mg/kg, the only effect observed was a slight increase in the severity of some histopathological changes in the thyroid. The NOAEL was 2 mg/kg (0.12 mg/kg/d) on the basis of thyroid effects. The NOAEL for reproductive toxicity was 15 mg/kg feed (0.9 mg/kg/d).

Both studies showed similar effects. The pup survival was markedly reduced in the F<sub>2</sub> generation of the two-generation study. The cause for the discrepancy with regard to liver and pituitary effects at lower doses in the one-generation study than in the two-generation study is unknown.



### Developmental toxicity

Tyl (1986a) exposed Sprague-Dawley rats to amitrole in doses of 0, 100, 500 or 1000 mg/kg/d by gavage on days 6 - 15 of gestation. Foetuses were examined on day 21 of gestation. Furthermore, 14 females per dose level were allowed to litter and wean, and were maintained until day 21 post partum. Food consumption and body weight gain of dams were reduced at the two higher doses, and maternal thyroid weights were increased at these dose levels. At the highest dose, foetotoxicity was observed, including reduced foetal body weight per litter, increased incidences of foetuses with unossified or poorly ossified bones and an increased number of foetuses with enlarged and/or dark thyroid. These latter findings in the thyroid were also observed in the 500 mg/kg/d dose group. No teratogenicity was observed. The postnatal evaluations indicated no other effects of the treatment. The NOAEL for maternal toxicity and developmental effects was 100 mg/kg/d (Tyl, 1986a).

In another developmental studies on rats (Long-Evans) with a similar design, a NOAEL of 1000 mg/kg/d was obtained (Bayer, 1977). The discrepancies in the NOAEL values may be due to the use of different rat strains.

Tjälve (1974) exposed NMRI mice on gestation days 6 - 18 to amitrole in drinking water (0, 500, 1000, 2500 and 5000 mg/l). At 1000 mg/l (140 mg/kg/d), maternal body weight gain and foetal weights were decreased and skeletal development was retarded. The highest dose produced an increased rate of resorptions. No teratogenic effects were observed. The NOAEL for maternal toxicity and developmental effects was 500 mg/l drinking water (70 mg/kg/d). Another study on mice reported a decrease of maternal and foetal body weights and an increased foetal mortality after oral administration of 215 mg/kg/d from days 6 - 14 of gestation (Bionetics Research Laboratories, 1968).

New Zealand white rabbits were exposed to amitrole by gavage at doses of 0, 4, 40, and 400 mg/kg/d on days 6 - 18 of gestation (Tyl, 1986b). The two higher doses caused a reduced maternal body weight gain. At 400 mg/kg/d, there were also increased liver weights. A dose-related increase in the number of abortions was observed (0, 1, 3 and 5 for controls, the 4, 40 and 400 mg/kg/d dose groups, respectively). The number of non-viable implants per litter was increased and the percentage of live foetuses per litter as well as the foetal weights per litter were decreased at 40 and 400 mg/kg/d (all statistically significant only at the highest dose). At 40 and 400 mg/kg/d, there were significant and dose-related increases in the incidence of malformations, especially of the head and limbs, as well as visceral and skeletal variations. Effects on the foetal thyroid (enlargement, discolouration) were also evident at these doses. The NOAEL for maternal and developmental toxicity was 4 mg/kg/d (Tyl, 1986b).

Kolb (1994) exposed Russian rabbits to amitrole in doses of 0, 5, 20 or 80 mg/kg/d by gavage on days 6 - 18 of gestation. Adverse effects were evident in the highest dose group, consisting of reduced maternal feed consumption and body weight gain as well as decreased foetal body weights and litter weights. There was no evidence of teratogenicity. The NOAEL for maternal and foetal toxicity was 20 mg/kg/d. The discrepancies between the rabbit studies may be due to the use of different strains and the higher doses used in the study by Tyl (1986b).

Subcutaneous exposure of C57/BL6 mice to 0, 215 and 464 mg/kg/d from days 6 -14 of gestation resulted in a decrease of maternal body weight and an increased foetal mortality at the higher dose (Bionetics Research Laboratories, 1968).

Dermal exposure of rabbits to 0, 1000, 1500 or 2000 mg/kg/d amitrole on days 7 to 19 of gestation (6 h/d) produced dose-related dermal irritation of the treated skin. Anorexia, decreased feed intake and decreased body weight gain as well as slightly reduced uterine weights at term were observed at 2000 mg/kg/d. At this dose, there were also effects on the offspring, consisting of reduced foetal weights, increased incidence of total resorptions (mainly early resorptions) and malformations. The NOAEL for maternal and developmental toxicity was 1500 mg/kg/d (Henwood, 1988).



## Recommendations

There is no sufficient human data to establish NOAEL/LOAEL or on which IOELV could be based for occupational exposure for amitrole. This value will protect about disturbance of thyroid.

In a patch test conducted with a human volunteer, amitrole exerted no primary dermal irritant effect. Patch testing with 1% amitrole on a weed control operator showed a strong positive vesicular reaction at 2 and 4 days, indicative of allergic contact dermatitis. Single oral dose of 100 mg amitrole inhibited radioiodine uptake by the thyroid of both normal and thyrotoxic subjects for 24 hours. A dose of 10 mg (0.15 - 0.2 mg/kg corresponding to about 1 mg/m<sup>3</sup>, Henschler, 1983) had a slight effect on iodine uptake. An epidemiological study was conducted on Swedish railway workers exposed to various herbicides. A statistically significant increase in the incidence of total tumors and lung tumors was found among workers exposed to amitrole and combinations of other herbicides (Axelson et al. 1974). However, in a follow-up study which combined data from the earlier study, results did not show a statistically significant increase in cancer incidence among those workers exposed to amitrole alone (Axelson et al. 1980). The thyroid function of 5 employees exposed to unknown level of amitrole between 3 and 16 years did not change (Bayer 1983). However, no data for repeated human exposure are available.

Amitrole has a low acute toxicity when tested in a several species by various routes of administration. The substance is rapidly absorbed from the gastrointestinal tract and lung. It is rapidly distributed throughout most body tissues, but with a slight accumulation in those tissues with a rapid cell turnover (bone marrow, spleen, thymus, gastrointestinal tract). Amitrole passes through the placenta into the fetuses with the same distribution pattern as in the mothers. Excretion is rapid after oral exposure. Within 24 hours, 70-95% of the administered radioactively-labelled compound is excreted via the urine, mainly as the parent compound. Based on experiences in animal studies, amitrole does not demonstrate irritating properties (ACGIH, 2001). Evidence of a moderate sensitizing potential was observed in a Magnusson-Kligman test but not in a Klecak open epicutaneous test. Therefore the data is too weak for sensitization assessment. Also, no reports on respiratory sensitization are available.

Exposure to amitrole leads to effects on the thyroid, which is also responsible for the induction of thyroid cancers and for reproductive toxicity in animals. The mechanism of this action is non-genotoxic.

Rodents, especially rats, are generally considered to be an especially sensitive species as Henschler (1983) showed a slight effect at 0.15 – 0.2 mg/m<sup>3</sup>. Rats are more prone to disturbances of the thyroid hormone balance than humans or dogs due to the lack of specific binding proteins in serum. Thyroid effects were reversible in the rat after cessation of exposure (Greim, 1998; Henschler, 1983; IARC, 2001; WHO, 1994a,b).

Several rat studies, highlighted in Table 2, showed that the uptake of iodine in the thyroid gland and other thyroid gland effects appeared to be the critical effect with a NOAEL around 0.1 mg/kg/d.

If we take the lowest NOAEL (0.025 mg/kg) and directly make a route-to route extrapolation (0.025 x 70/10) the OEL should be 0.2 mg/m<sup>3</sup>. As the lowest NOAEL is used and rats are assumed to be the most sensitive species, no assessment factor is needed.

If departure from the dog study with a NOAEL of 0.3 mg/kg/d for the route-to-route extrapolation and accepting an assessment factor of 10 as dog are not considered an especially sensitive species, the occupational exposure limit estimate is also estimated to be 0.2 mg/m<sup>3</sup>. This value will protect against the disturbance of thyroid function in human, where a slight effect occurred at  $\geq 0.15$  mg/kg, which corresponds to about 1 mg/m<sup>3</sup> from a route-to-route extrapolation assuming a 70 kg person inhaling 10 m<sup>3</sup> per day.

With respect to the greater sensitivity of rodents to thyroid effects, amitrole would not be expected to produce thyroid cancer in humans exposed to concentrations that do not alter thyroid homeostasis (IARC, 2001). On the basis of the lack of genotoxicity, the benign



pituitary tumours in rats and liver tumours in mice may be considered to be of non-genotoxic origin (IARC, 2001).

A "skin" notation is not proposed. Although one study reported a relevant dermal absorption of 30% in rabbits after exposure for 24 h (WHO, 1993), EC (2001) documented a dermal absorption of only 1%. No indication exists for a relevant skin sensitising potency of amitrole.

There are no data available for biological monitoring.

At the recommended IOELV, no analytical difficulties are expected.

Table 2. Summary of lowest-observed-adverse-effects levels (LOAELs) and no-observed-adverse-effect levels (NOAELs) for amitrole.

Species	Type of study	Effects	Reference
Humans	Uptake of iodine in the thyroid gland	Slight effect: 0.15-0.2 mg/kg	Henschler, 1983
Rats	Exposures were through weeks: 1-13, 40-52 and 78-90 for 5h/d, 5d/w.	At about 25 mg/m <sup>3</sup> : decreased T <sub>3</sub> and T <sub>4</sub> , increased thyroid weight and follicular hyperplasia.	Becci, 1983
Rats	Amitrole in the diet for about three months. Uptake of iodine in the thyroid gland.	Reduced iodine uptake at 0.1 mg/kg/d (LOAEL). NOAEL: 0.025 mg/kg/d Another study: NOAEL: 0.1 mg/kg/d	Fregly, 1968 Unpublished study
Rats	Life-long amitrol in the diet. Tumours and thyroid function.	NOAEL: 1.5 mg/kg/d	Steinhoff et al., 1983
Rats	Amitrole in the diet. Two-generation study.	About 2 mg/kg/d were without effect on reproduction and development, but thyroid gland hypertrophy appeared, i.e. LOAEL.	Gaines et al., 1973
Rats	One-generation study	About 1 mg/kg/d showed dose-related effects in the liver and pituitary gland in F <sub>0</sub> and F <sub>1</sub> animals with decreased colloid content in females of both generations, i.e. LOAEL.	Savary, 1994
Rats	Two generation study	About 7 mg/kg/d caused increased death (F <sub>1</sub> and F <sub>2</sub> ) and multiple organ effects. The NOAEL was 0.12 mg/kg/d on the thyroid effect and the NOAEL for reproductive toxicity was 0.9 mg/kg/d.	Richard, 1995
Beagle dogs	Amitrole in the diet for 1 year.	Thyroid gland effects (T <sub>3</sub> and T <sub>4</sub> , hypertrophy, follicular hyperplasia) LOAEL: 13 mg/kg/d NOAEL: 0.3 mg/kg/d	Bayer, 1994



## References

- ACGIH, American Conference of Governmental Industrial Hygienists, 2001  
Amitrole  
In: ACGIH, American Conference of Governmental Industrial Hygienists, Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices, 2004. Cincinnati, OH, 2001
- Axelsson, O.; Sundell, L., 1974  
Herbicide exposure, mortality and tumor incidence. An epidemiological investigation on Swedish railroad workers  
Work, Environment, Health, **11**, 1974, 21-28, cited in IARC, 2001
- Axelsson, O.; Sundell, L.; Andersson, K.; Edling, C.; Hogstedt, C., 1980  
Herbicide exposure and tumor mortality. An updated epidemiologic investigation on Swedish railroad workers  
Scandinavian Journal of work, Environment, and Health, **6**, 1980, 73-79, cited in IARC, 2001
- Astwood, E.B., 1960  
Cranberries, turnips, and goiter  
Journal of the American Medical Association, **172**, 1960, 1219-1220
- Becci, P., 1983  
Evaluation of the chronic inhalation toxicity and carcinogenicity of amitrole in rats. Unpublished report No 5821 (dated: 13-01-1983) from Food and Drug Research Laboratories, Inc., Submitted to WHO by Bayer AG, cited in WHO, 1993
- Balkissoon, R.; Murray, D.; Hoffstein, V., 1992  
Alveolar damage due to inhalation of amitrole-containing herbicide  
Chest, **101**, 1992, 1174-1176, cited in IARC, 2001
- Baughner, D.G.; Bookbinder, M.G.; Blundell, K.R., 1982  
Final Report – Medical monitoring and assessment of exposure of workers applying amitrole to utility rights-of-way  
Rockville, Maryland, Dynamic Corporation, Enviro Control Division. Unpublished report, cited in WHO, 1994a
- Bayer AG, 1977  
Amitrole. Dominant-Letal-Test an der männlichen Maus zur Prüfung auf mutagene Wirkung  
Bayer AG, Wuppertal, Nr. 6679, unpublished study, 1977, cited in Greim, 1998
- Bayer AG, 1983  
Produktion von Aminotriazol: Gesundheitsüberwachung. Brief vom 22.04.1983 von Dr. Mischke, Ärztliche Abteilung, Bayer AG, Leverkusen, unpublished, cited in Greim, 1998
- Bayer AG, 1984  
Aminotriazol. Untersuchungen auf hautsensibilisierende Wirkung bei Meerschweinchen. Bayer AG, Wuppertal, No. 12607, unpublished study, cited in Greim, 1998 and WHO, 1993 (cited as Mihail, 1984)
- Bayer AG, 1985  
Aminotriazol (c.n.: Amitrol). Untersuchung auf hautsensibilisierende Wirkung im offenen Epicutan-Test an Meerschweinchen. Bayer AG, Wuppertal, No. 13775, unpublished study, cited in Greim, 1998 and WHO 1993 (cited as Mihail, 1985)
- Bayer AG, 1994  
Technical grade amitrole: A chronic toxicity feeding study in the Beagle dog. No. R6321 from Miles Inc., Agriculture Division Toxicology, South Metcalf, Stilwell, Kansas, USA, Unpublished study, cited in Greim, 1998



- Bhuyan, K.C.; Bhuyan, D.K.; Katzing, H.M., 1973  
Amitrole-induced cataract and inhibition of lens catalase in rabbit  
*Ophthalmol Research*, **5**, 1973, 236-247, cited in WHO, 1994a
- Bionetics Research Laboratories, 1968  
Evaluation of carcinogenic, teratogenic, and mutagenic activities of selected pesticides and industrial chemicals. Volume II: Teratogenic study in mice and rats. Report PB-223 160 (dated: August 1968) from Bionetic Research Labs, Inc., prepared for National Cancer Institute, cited in WHO, 1993
- Cox, G.E.; Re, T.A., 1978  
Inhalation toxicity study with 3-amino-1,2,4-triazole (Amitrole) in adult Fisher 344 rats. Unpublished report No 5492 (dated: 08-09-1978) from Food and Drug Research Laboratories, Inc., Submitted to WHO by Bayer AG, cited in WHO, 1993 and 1994a
- EC, European Commission, 2001  
Review report for the active substance amitrole, 22 March 2001  
online:  
[http://ec.europa.eu/food/plant/protection/evaluation/existactive/list1\\_amitrol\\_en.pdf](http://ec.europa.eu/food/plant/protection/evaluation/existactive/list1_amitrol_en.pdf)
- ECB, European Chemicals Bureau, 2000  
IUCSID, International Uniform Chemical Information Database. Edition II  
EUR 19559 EN, European Commission, 2000
- Elesa, J.R., 1954  
Aminotriazole: acute dermal application, acute eye application  
Falls Church, Virginia, Hazleton Laboratories, unpublished report, cited in WHO, 1994a
- English, J.S.C.; Rycroft, R.J.G.; Calnan, C.D., 1986  
Allergic contact dermatitis from aminotriazole  
*Contact Dermatitis*, **14**, 1986, 255-256, cited in IARC, 2001 and Greim, 1998
- Fang S.C., George M., Yu T.C. (1964): Metabolism of 3-amino-1,2,4-triazole-5-C14 by rats.  
*J. Agric. Food Chem.* **12**, 219-223.
- Fang S.C., Khanna S., Rao A.V. (1966): Further study on the metabolism of labelled 3-amino-1,2,4-triazole (ATA) and its plant metabolites in rats. *J. Agric. Food Chem.* **14**, 262-265.
- FAO, Food and Agriculture Organization of the United Nations, 1997  
Amitrole  
In: FAO, Food and Agriculture Organization of the United Nations, Pesticide Residues in Food - 1997. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group, Lyon 22 September - 1 October 1997  
<http://www.inchem.org/documents/jmpr/jmpmono/v097pr03.htm>
- Fregly, M.J., 1968  
Effect of aminotriazole on thyroid function in the rat  
*Toxicology and Applied Pharmacology*, **13**, 1968, 271-286
- Gaines, T.B.; Kimbrough, R.D.; Linder, R.E., 1973  
The toxicology of amitrole in the rat  
*Toxicology and Applied Pharmacology*, **26**, 1973, 118-129
- Geldmacher von Mallinckrodt, M.; Schmidt, H.P., 1970  
Toxicity and metabolism of aminotriazole in man  
*Archives of Toxicology*, **27**, 1970, 13-18, cited in WHO, 1993 and 1994a
- Greim, H., 1983  
Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten, Loseblattsammlung, 9. Lfg.  
DFG, Deutsche Forschungsgemeinschaft, WILEY-VCH Verlag, Weinheim, 1983





- Greim, H., 1998  
Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten, Loseblattsammlung, 26. Lfg.  
DFG, Deutsche Forschungsgemeinschaft, WILEY-VCH Verlag, Weinheim, 1998
- Greim, H., 1999  
Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten, Loseblattsammlung, 28. Lfg.  
DFG, Deutsche Forschungsgemeinschaft, WILEY-VCH Verlag, Weinheim, 1999
- Greim, H., 2002  
Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten, Loseblattsammlung, 34. Lfg.  
DFG, Deutsche Forschungsgemeinschaft, WILEY-VCH Verlag, Weinheim, 2002
- Hecht, P., 1954  
Aminotriazol - Letter report of May 11, 1954 from Bayer AG Institute of Toxicology. Submitted to WHO by Bayer AG, unpublished report, cited in WHO, 1993 and 1994a
- Henschler, D., 1983  
Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten, Loseblattsammlung, 9. Lfg.  
DFG, Deutsche Forschungsgemeinschaft, VCH Verlag Weinheim, 1983
- Henwood, S.M., 1988  
Percutaneous teratology study with amitrole technical in rabbits. Unpublished report on number HLA 6224-107 from Hazelton Laboratories America, Inc., submitted to WHO by Rhône-Poulenc Ag. Company, cited in WHO, 1993
- IARC, International Agency for Research on Cancer, 2001  
IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol. 79. Some Thyrotropic Agents WHO, World Health Organization, Geneva, 2001
- IPCS (1994): International Programme on Chemical Safety. Health and Safety Guide No. 85. Amitrole. UNEP, ILO, WHO, Geneva.
- Kolb, J., 1994  
Amitrole: Developmental toxicity study in rabbits after oral administration. Unpublished report No. 23486 (study No. T5044250) from Bayer AG, Department of Toxicology, Wuppertal, Germany. Submitted to WHO by Bayer AG, Leverkusen, Germany, cited in FAO, 1997
- Mihail, F.; Schilde, B., 1984  
Aminotriazole - Study for subacute dermal toxicity to rabbits. Unpublished report No 12369 (dated: 09-01-1984) from Bayer AG Institute of Toxicology. Submitted to WHO by Bayer AG, cited in WHO, 1993
- Napalkov, N.P., 1962  
[Blastomogenic action of 3-amino-1,2,4-triazole]  
Gigiena Truda i Professional'nye Zabolevaniya, **6**, 1962, 48-51, cited in IARC, 2001
- Pontecorvo, G.; Fantaccione, S., 2006  
Recombinogenic activity of 10 chemical compounds in male germ cells of *Drosophila melanogaster*  
Ecotoxicology and Environmental Safety, **65**, 2006, 93-101
- Richard, J., 1995  
Two generation study by oral route (dietary admixture) in rats. Unpublished report No. 6432 (study No. T5041262) from Centre International de Toxicologie, Evreux, France. Submitted to WHO by Bayer AG, Leverkusen, Germany, cited in FAO, 1997
- Savary, M.-H., 1994  
Preliminary study to a two-generation study by oral route (dietary admixture) in rats. Unpublished report No. 6306 (study No. T6041263) from Centre International de



- Toxicologie, Evreux, France. Submitted to WHO by Bayer AG, Leverkusen, Germany, cited in FAO, 1997
- Schneider K., Oltmanns J., Hassauer M., 2004  
Allometric principles for interspecies extrapolation in toxicological risk assessment – empirical investigations. *Regulatory Tox and Pharm*, 2004, 39(3), 334-347.
- Steinhoff, D.; Weber, H.; Mohr, U.; Boehme, K., 1983  
Evaluation of amitrole (aminotriazole) for potential carcinogenicity in orally dosed rats, mice, and golden hamsters  
*Toxicology and Applied Pharmacology*, **69**, 1983, 161-169, cited in IARC, 2001
- Tjälve, H., 1974  
Fetal uptake and embryogenetic effects of aminotriazole in mice  
*Archives of Toxicology*, **33**, 1974, 41-48, cited in Greim, 1999 and WHO, 1994a
- Tyl, R.W., 1986a  
Teratogenicity evaluation of aminotriazole technical administered by gavage to CD rats. Unpublished report No 49-45 (dated: 29-05-1986) from Union Carbide Bushy Run Research Center. Submitted to WHO by Bayer AG and Rhône-Poulenc Ag. Company, cited in WHO, 1993
- Tyl, R.W., 1986b  
Teratogenicity evaluation of aminotriazole technical administered by gavage to New Zealand White rabbits. Unpublished report No 49-66 (dated: 30-05-1986) from Union Carbide Bushy Run Research Center. Submitted to WHO by Rhône-Poulenc Ag. Company, cited in WHO, 1993
- Vesselinovitch, S.D., 1983  
Perinatal hepatocarcinogenesis  
*Biological Research in Pregnancy and Perinatology*, **4**, 1983, 22-25
- WHO, World Health Organization, 1993  
Amitrole (856)  
In: WHO, World Health Organization, *Pesticide Residues in Food - 1993. Evaluations. Part II - Toxicology*, 1993  
<http://www.inchem.org/documents/jmpr/jmpmono/v93pr02.htm>
- WHO, World Health Organization, 1994a  
Environmental Health Criteria 158, Amitrole  
IPCS, International Programme on Chemical Safety; World Health Organization, Geneva, 1994
- WHO, World Health Organization, 1994b  
WHO/FAO Data Sheet on Pesticides No. 79. Amitrole  
[http://www.inchem.org/documents/pds/pds/pest79\\_e.htm](http://www.inchem.org/documents/pds/pds/pest79_e.htm)