ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

BRODIM LIGHT

Product type(s)

PT14: Rodenticides

Authorisation number: IT/2018/00446/AUT

R4BP asset number: IT-0018474-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	RATOMIDE PASTA
	BRODIM LIGHT

1.2. Authorisation holder

Name and address of the authorisation holder	Name	COLKIM S.R.L.
	Address	Via Piemonte, 50 40064 Ozzano dell'Emilia Italy
Authorisation number		IT/2018/00446/AUT
R4BP asset number		IT-0018474-0000
Date of the authorisation		05/02/2014
Expiry date of the authorisation		31/12/2026

1.3. Manufacturer(s) of the product

Name of manufacturer	Colkim S.r.l.
	Via Piemonte, 50 40064 Ozzano dell'Emilia (BO) Italy
Location of manufacturing sites	Via Piemonte, 50 40064 Ozzano dell'Emilia (BO) Italy

Name of manufacturer	Industrialchimica S.r.l.
Address of manufacturer	Via Sorgaglia, 40 35020 Arre (PD) Italy
Location of manufacturing sites	Via Sorgaglia, 40 35020 Arre (PD) Italy

1.4. Manufacturer(s) of the active substance(s)

Active substance	Brodifacoum
Name of manufacturer	Activa S.rl
Address of manufacturer	via Feltre 32 20132 milano Italy
Location of manufacturing sites	Tezza S.r.l-via Tre Ponti 22 37050 S. Maria di Zevio Italy

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro -1-napthyl]-4-hydroxycoumarin	active substance	56073-10-0	259-980-5	0,0025

2.2. Type(s) of formulation

RB Bait (ready for use)

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H373: May cause damage to organs (blood) through prolonged or repeated exposure {1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard:}.
Precautionary statements	P102: Keep out of reach of children. P301+P310: IF SWALLOWED: Immediately call a POISON CENTER or a doctor/physician.
	P308+P313: IF exposed or concerned: Get medical advice.
	P501: Dispose of contents to according to the national rules.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice – professional – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: house mouse Development stage: juveniles to adults
Field(s) of use	indoor use
Application method(s)	Method: bait application Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 20 g - 40 g pasta bait per bait point. Dilution (%): 0,0025 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	professional
Pack sizes and packaging material	Minimum pack size of 1,5 kg Bait formulations: Paste
	Number of packed bags per packaging: 3 Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 15 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 1,5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 4 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: Grams/kg of bait per packed bag: 5 Kg (containing rodenticide

29/05/2024 - IUCLID 6

Polyesther, cardboard

baits of 10-20 g in sachets of pasta bait)

Packaging material: Plastic composite, Plastic, PE/Aluminium/

4.1.1. Use-specific instructions

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- **4.1.2.** Use-specific risk mitigation measures
- 4.1.3. Use-specific risk mitigation measures
- 4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment
- 4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- **4.1.6.** Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage
- 4.2. Use description

Table 2. Rats - professional - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: brown rat Development stage: juveniles to adults
Field(s) of use	indoor use
Application method(s)	Method: bait application Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g - 200 g per bait station. Dilution (%): 0,0025 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	professional

Pack sizes and packaging material Minimum pack size of 1,5 kg. - Bait formulations: Paste Number of packed bags per packaging: 3 Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

Number of packed bags per packaging: 15 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 1,5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 4 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1

Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
Number of packed bags per packaging: 5 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

4.2.1. Use-specific instructions

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

- 4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment
- 4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- 4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.3. Use description

Table 3. House mice and Rats – professionals – around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides

Target organism(s) (including development	Scientific name: Mus musculus
stage)	Common name: house mouse
	Development stage: juveniles to adults
	Scientific name: Rattus norvegicus
	Common name: brown rat Development stage: Juveniles to adults
77.11() 0	
Field(s) of use	Other
	Around buildings
Application method(s)	Method: bait application
	Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: House mice: 20 – 40 g pasta bait per bait point Brown rat: 100 g - 200 g pasta bait per bait point
	Dilution (%): 0,0025
	Number and timing of application: The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
Category(ies) of users	professional
Pack sizes and packaging material	
	Minimum pack size of 1,5 kg.
	- Bait formulations: Paste
	Number of packed bags per packaging: 3
	Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
	Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 15 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 1,5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2


Grams/kg of bait per packed bag: 5 Kg (containing rodenticide

29/05/2024 - IUCLID 6

baits of 10-20 g in sachets of pasta bait)

Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 4 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)



4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice at least every 2 to 3 days at] [for rats only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

Do not apply this product directly in the burrows.

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.4. Use description

Table 4. House mice – general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: house mouse Development stage: juveniles to adults
Field(s) of use	indoor use
Application method(s)	Method: bait application Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 20 g - 40 g pasta bait per bait point. Dilution (%): 0,0025 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Up to 150 g (containing rodenticide baits of 10-20 g in sachets of pasta bait). Packaging material: Plastic, Plastic Composite, PE/Aluminium/ Polyesther

4.4.1. Use-specific instructions

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- 4.4.2. Use-specific risk mitigation measures
- 4.4.3. Use-specific risk mitigation measures
- 4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment
- 4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- **4.4.6.** Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.5. Use description

Table 5. Rats –general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles to adults
Field(s) of use	indoor use
Application method(s)	Method: bait application
	Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g - 200 g per bait point. Dilution (%): 0,0025 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Up to 150g (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic, Plastic Composite, PE/Aluminium/Polyesther.

4.5.1. Use-specific instructions

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- 4.5.2. Use-specific risk mitigation measures
- 4.5.3. Use-specific risk mitigation measures
- 4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment
- 4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- 4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.6. Use description

Table 6. Rats – professional and general public – around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage) Field(s) of use	Scientific name: Rattus norvegicus Common name: brown rat Development stage: Juveniles to adults Other
	Around buildings.
Application method(s)	Method: bait application
	Detailed description: Ready-to-use pasta bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g - 200 g pasta bait per bait point. Dilution (%): 0,0025 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Up to 150g (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic, Plastic Composite, PE/Aluminium/Polyesther.

4.6.1. Use-specific instructions

⁻ Place the bait stations in areas not liable to flooding.

- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

For professional use around building: Do not apply this product directly in the burrows.

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

For professional use: When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.7. Use description

Table 7. House mice and/or rats – trained professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Common name: house mouse Development stage: juveniles to adults
	Scientific name: Rattus norvegicus Common name: brown rat Development stage: juveniles to adults
Field(s) of use	indoor use Indoor
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting point as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations Application of the product in pulsed baiting

Application rate(s) and frequency	Application rate: House mice: $20 - 40$ g pasta bait per bait point; Brown rat: 100 g - 200 g pasta bait per bait point
	Dilution (%): 0 - Ready-to-use pasta bait
	Number and timing of application: See section 5.1
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size of 1,5 kg Bait formulations: Paste
	Number of packed bags per packaging: 3
	Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
	Number of packed bags per packaging: 2
	Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
	Number of packed bags per packaging: 10
	Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/

Number of packed bags per packaging: 15 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 1,5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 4 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1

Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 5 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

4.7.1. Use-specific instructions

Remove the remaining product at the end of the treatment period.

Additional specific instructions for use for application in covered and protected baiting points

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Additional specific instructions for use "pulsed baiting"

- Replace eaten bait only after 3 days and then at maximum 7 days intervals. Collect any spilled bait and dead rodents.

4.7.2. Use-specific risk mitigation measures

4.7.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

4.7.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.7.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.7.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage


Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (Trained professional)" are implemented.

4.8. Use description

Table 8. Mice and/or rats – trained professionals – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides

Target organism(s) (including development stage)	Scientific name: Mus musculus Common name: house mouse Development stage: juveniles to adults Scientific name: Rattus norvegicus
	Common name: brown rat Development stage: juveniles to adults
Field(s) of use	outdoor use
Application method(s)	Around buildings Method: bait application
1 application method(g)	Detailed description: - Ready-to-use bait to be used in tamper- resistant bait stations or covered and protected baiting point as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations Direct application of ready-to-use bait into the burrow - Application of the product in pulsed baiting
Application rate(s) and frequency	Application rate: House mice: 20 – 40 g pasta bait per bait point; Brown rat: 100 g - 200 g pasta bait per bait point
	Dilution (%): 0 - Ready-to-use pasta bait
	Number and timing of application: See section 5.1
Category(ies) of users	trained professional
Category(ies) of users Pack sizes and packaging material	trained professional Minimum pack size of 1,5 kg Bait formulations: Paste
	Minimum pack size of 1,5 kg.
	Minimum pack size of 1,5 kg Bait formulations: Paste
	Minimum pack size of 1,5 kg Bait formulations: Paste Number of packed bags per packaging: 3 Grams/kg of bait per packed bag: 500 gr (containing rodenticide





Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 5 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)

Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard	

4.8.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.

Additional specific instructions for use for application in covered and protected baiting points

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Additional specific instructions for use for direct application in the burrows

- Baits must be placed to minimise the exposure to non-target species and children.
- Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

Additional specific instructions for use "pulsed baiting"

- Replace eaten bait only after 3 days and then at maximum 7 days intervals. Collect any spilled bait and dead rodents.

4.8.2. Use-specific risk mitigation measures

4.8.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Additional specific risk mitigation measures for direct application in the burrow

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign

4.8.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.8.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.8.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage


Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (Trained professional)" are implemented.


4.9. Use description

Table 9. Rats – trained professionals – Outdoor open areas & waste dumps

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development	Scientific name: Rattus norvegicus Common name: brown rat
stage)	Development stage: juveniles to adults
Field(s) of use	Outdoor use Outdoor open areas & waste dumps
Application method(s)	Method: bait application
	Detailed description: - Ready-to-use bait to be used in tamper- resistant bait stations or covered and protected baiting point as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations Direct application of ready-to-use bait into the burrow - Application of the product in pulsed baiting

Application rate(s) and frequency	Application rate: 100 g - 200 g pasta bait per bait point
	Dilution (%): 0 - Ready-to-use pasta bait
	Number and timing of application: See section 5.1
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size of 1,5 kg Bait formulations: Paste
	Number of packed bags per packaging: 3
	Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
	Number of packed bags per packaging: 2
	Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard
	Number of packed bags per packaging: 10
	Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)





Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 20 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 25 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 5 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

4.9.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.

Additional specific instructions for use for application in covered and protected baiting points

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

Additional specific instructions for use for direct application in the burrows

- Baits must be placed to minimise the exposure to non-target species and children.

- Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. **Additional specific instructions for use "pulsed baiting"**
- Replace eaten bait only after 3 days and then at maximum 7 days intervals. Collect any spilled bait and dead rodents.

4.9.2. Use-specific risk mitigation measures

4.9.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Additional specific risk mitigation measures for direct application in the burrow

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign

4.9.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

- 4.9.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- 4.9.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (Trained professional)" are implemented.

4.10. Use description


Table 10. Rats – trained professionals – sewers

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development	Scientific name: Rattus norvegicus
stage)	Common name: brown rat
	Development stage: juveniles to adults
Field(s) of use	outdoor use
	Sewers
Application method(s)	Method: bait application
	Detailed description: - Ready-to-use bait to be used in tamper- resistant bait stations or covered and protected baiting point as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: 100 g - 200 g pasta bait per bait point
	Dilution (%): 0 - Ready-to-use pasta bait
	Number and timing of application: See section 5.1
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size of 1,5 kg Bait formulations: Paste
	Number of packed bags per packaging: 3
	Grams/kg of bait per packed bag: 500 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait)
	Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 750 gr (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 15 Grams/kg of bait per packed bag: 1 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 1,5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard

Number of packed bags per packaging: 4 Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 1 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 2 Grams/kg of bait per packed bag: 10 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait) Packaging material: Plastic composite, Plastic, PE/Aluminium/ Polyesther, cardboard Number of packed bags per packaging: 10



Number of packed bags per packaging: 5

Grams/kg of bait per packed bag: 5 Kg (containing rodenticide baits of 10-20 g in sachets of pasta bait)

Packaging material: Plastic composite, Plastic, PE/Aluminium/Polyesther, cardboard

4.10.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- Follow any additional instructions provided by the relevant code of best practice.

Additional specific instructions for use for application in covered and protected baiting points

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.


Additional specific instructions for use in sewers

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

4.10.2. Use-specific risk mitigation measures

4.10.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.



Additional specific risk mitigation measures for use in sewers

- Place baits only in sewer systems which are connected to the sewage treatment plant
- Do not use this product in pulsed baiting treatment

4.10.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

- 4.10.5. Where specific to the use, the instructions for safe disposal of the product and its packaging
- 4.10.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (Trained professional)" are implemented.

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

PROFESSIONAL USE ONLY

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-nticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.
- Instruction for use that are "bait specific":
- Bait in sachets: Do not open the sachets containing the bait.

PROFESSIONAL AND TRAINED PROFESSIONAL USE

- Read and follow the product information as well as any information ccompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.

TRAINED PROFESSIONAL USE ONLY

- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

GENERAL PUBLIC USE ONLY

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Do not place bait stations near water drainage systems where they can come into contact with water.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

PROFESSIONAL AND TRAINED PROFESSIONAL USE

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Dispose dead rodents in accordance with local requirements.

PROFESSIONAL USE ONLY

- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week)
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that:
- the product shall not be supplied to the general public (e.g. "for professionals only").
- the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product ecommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.

TRAINED PROFESSIONAL USE ONLY

- The product information (i.e. lable and/or leaflet) shall clearly show that the product shall only be subblied t90 trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only")
- Do not use in areas where resistance to the active substance can be suspected.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensiles used in covered and protected bait points with water between applications.

GENERAL PUBLIC USE ONLY

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).
- The product information (i.e. label and/or leaflet) shall clearly show that:
- the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
- users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.
- Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet.

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre.
- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent, a dye and a flavouring.