

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

Contrac Blox

Product type(s)

PT14: Rodenticides

Authorisation number: IE/BPA 70551

R4BP asset number: IE-0006240-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	Conrac Blox Tomcat Block Bait Tomcat Blox Tomcat 2 Blox Hawk Blox
---------------	---

1.2. Authorisation holder

Name and address of the authorisation holder	Name	Bell Laboratories Netherlands B.V.
	Address	De Cuserstraat 93 1081 CN Amsterdam Netherlands
Authorisation number	IE/BPA 70551	
<i>R4BP asset number</i>	IE-0006240-0000	
Date of the authorisation	30/01/2014	
Expiry date of the authorisation	01/07/2025	

1.3. Manufacturer(s) of the product

Name of manufacturer	Bell Laboratories Inc - European Division
Address of manufacturer	3699 Kinsman Boulevard 53704 Madison, Wisconsin United States (the)
Location of manufacturing sites	3699 Kinsman Boulevard 53704 Madison, Wisconsin United States (the)

1.4. Manufacturer(s) of the active substance(s)

Active substance	Bromadiolone
Name of manufacturer	Bell Laboratories Inc
Address of manufacturer	3699 Kinsman Boulevard 53704 Madison, Wisconsin United States (the)
Location of manufacturing sites	3699 Kinsman Boulevard 53704 Madison, Wisconsin United States (the)

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[3-(4'-Bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	active substance	28772-56-7	249-205-9	0,005

2.2. Type(s) of formulation

RB Bait (ready for use)

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H360D: May damage the unborn child. H372: Causes damage to organs blood through prolonged or repeated exposure { 1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard: }.
Precautionary statements	P201: Obtain special instructions before use. P202: Do not handle until all safety precautions have been read and understood. P281: Use personal protective equipment as required. P270: Do not eat, drink or smoke when using this product. P264: Wash hands thoroughly after handling. P308+P313: IF exposed or concerned: Get medical attention. P405: Store locked up.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. Use #1 - House mice and/or rats - professional user - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: all
Field(s) of use	indoor use Indoors
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Use against rats: Up to 225 g bait per baiting point at 5 to 10 metre intervals Use against mice: 5 – 28 g bait per baiting point at 2 to 4 metre intervals
Category(ies) of users	professional
Pack sizes and packaging material	Minimum pack size 2.5 kg For use against mice only: Loose 5 g and 28 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 10 kg For use against rats only: Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 9 kg 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to 10 kg

	1 x 28 g block in prefilled polypropylene mouse bait station/ Bait stations in multiples - 2.5 to 10 kg
--	---

4.1.1. Use-specific instructions

- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- [*When available*] Follow any additional instructions provided by the relevant code of best practice.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

See section 5.2

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

4.2. Use description

Table 2. Use #2 - House mice and/or rats - Trained professionals - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: all
Field(s) of use	indoor use Indoors
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected baiting points
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Use against rats: Up to 225 g bait per baiting point at 5 to 10 metre intervals Use against mice: 5 – 28 g bait per baiting point at 2 to 4 metre intervals
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size 2.5 kg For use against mice only: Loose 5 g and 28 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 10 kg For use against rats only: Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 9 kg 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to 10 kg 1 x 28 g block in prefilled polypropylene mouse bait station/ Bait stations in multiples - 2.5 to 10 kg

--	--

4.2.1. Use-specific instructions

- Remove the remaining product at the end of treatment period.

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product in pulsed baiting treatments.

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

4.3. Use description

Table 3. Use #3 – Mice and/or rats – Professional users – Outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: all
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Use against rats: Up to 225 g bait per baiting point at 5 to 10 metre intervals Use against mice: 5 – 28 g bait per baiting point at 2 to 4 metre intervals
Category(ies) of users	professional
Pack sizes and packaging material	Minimum pack size 2.5 kg For use against mice only: Loose 5 g and 28 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 10 kg For use against rats only: Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 9 kg 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to 10 kg 1 x 28 g block in prefilled polypropylene mouse bait station/ Bait stations in multiples - 2.5 to 10 kg

--	--

4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas non-labile to flooding.

-
For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

-
For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [*When available*] Follow any additional instructions provided by the relevant code of best practice.

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

4.4. Use description

Table 4. Use #4 – Mice and/or rats – Trained professionals – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: all
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected baiting points
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Use against rats: Up to 225 g bait per baiting point at 5 to 10 metre intervals Use against mice: 5 – 28 g bait per baiting point at 2 to 4 metre intervals Permanent Baiting (Rats): Up to 225 g bait per baiting point
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size 2.5 kg For use against mice only: Loose 5 g and 28 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 10 kg For use against rats only: Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 9 kg 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to 10 kg 1 x 28 g block in prefilled polypropylene mouse bait station/ Bait stations in multiples - 2.5 to 10 kg

--	--

4.4.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species
- Where possible, it is recommended that the area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.4.2. Use-specific risk mitigation measures

4.4.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation

4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.4.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

4.5. Use description

Table 5. Use #5 – Rats – Trained professionals – Outdoor open areas & waste dumps

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All
Field(s) of use	outdoor use Outdoor open areas and waste dumps
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected baiting points
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Use against rats: Up to 225 g bait per baiting point at 5 to 10 metre intervals Permanent Baiting (Rats): Up to 225 g bait per baiting point
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size 2.5 kg Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5to 9 kg. 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to10 kg

--	--

4.5.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
 - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
 - Remove the remaining product at the end of treatment period
 - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species
- Where possible, it is recommended that the area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population
- [When available] Follow any additional instructions provided by the relevant code of best practice

4.5.2. Use-specific risk mitigation measures

4.5.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
 - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
 - Do not use this product in pulsed baiting treatments.
 - Do not apply this product directly in the burrows.
- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation

4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

4.6. Use description

Table 6. Use #6 – Rats – Trained professionals – sewers

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All
Field(s) of use	Other Sewers
Application method(s)	Method: bait application Detailed description: - Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. - Covered and protected baiting points
Application rate(s) and frequency	Application rate: See below Dilution (%): 0 Number and timing of application: Up to 225 g per manhole
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size 2.5 kg Loose 225 g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box or chipboard box - 2.5 to 9 kg. 2 to 10 x 28 g or 1 x 225 g blocks in prefilled polypropylene rat bait station supplied in a polyethylene pail, chipboard box, cardboard box, or polyethylene plastic pouch with ziploc seal. Bait stations in multiples - 2.5 to 10 kg

4.6.1. Use-specific instructions

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

- Do not use this product in pulsed baiting treatments.

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See section 2.5.3

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

See 2.5.4

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See 2.5.5

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

Trained professional users

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Professional users

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 2.3 of First Renewal PAR for the information to be shown on the label).
- [If national policy or legislation require it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

Trained professional users

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements
- Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site.

Permanent baiting: Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient. The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation. Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting.

- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

Professional users

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

-
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").
 - Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
 - Antidote: Vitamin K1 administered by medical/veterinary personnel only.
 - In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Medical professionals should contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-809 2166), for further advice.
- Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison centre".
 - Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

Trained professional users

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.
- Use of gloves is recommended.

Professional users

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 3 year

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.
- Use Biocides Safely and Sustainably
- It is illegal to use this product for uses or in a manner other than that prescribed on this label.
- Poison Information: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.
- Please note IRELAND applies a minimum Professional pack size of 2.5 Kg instead of 3.0 kg.