

CLH report

Proposal for Harmonised Classification and Labelling

Based on Regulation (EC) No 1272/2008 (CLP Regulation),
Annex VI, Part 2

International Chemical Identification:

2-Ethylhexyl-10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate; DOTE

EC Number: 239-622-4
CAS Number: 15571-58-1
Index Number: 050-027-00-7

Contact details for dossier submitter:

BAuA
Federal Institute for Occupational Safety and Health
Federal Office for Chemicals
Friedrich-Henkel-Weg 1-25
44149 Dortmund, Germany

Version: 2.0
Date: October 2017

CONTENTS

1	IDENTITY OF THE SUBSTANCE	1
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE.....	1
1.2	COMPOSITION OF THE SUBSTANCE	2
2	PROPOSED HARMONISED CLASSIFICATION AND LABELLING.....	3
2.1	PROPOSED HARMONISED CLASSIFICATION AND LABELLING ACCORDING TO THE CLP CRITERIA	3
3	HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING	5
4	JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL.....	5
5	IDENTIFIED USES	6
6	DATA SOURCES.....	6
7	PHYSICOCHEMICAL PROPERTIES.....	7
8	EVALUATION OF PHYSICAL HAZARDS	9
9	TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)	10
9.1	SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED TOXICOKINETIC INFORMATION ON THE PROPOSED CLASSIFICATION(S).....	10
10	EVALUATION OF HEALTH HAZARDS.....	11
10.1	ACUTE TOXICITY - ORAL ROUTE	12
10.2	ACUTE TOXICITY - DERMAL ROUTE	12
10.3	ACUTE TOXICITY - INHALATION ROUTE	12
10.4	SKIN CORROSION/IRRITATION	13
10.5	SERIOUS EYE DAMAGE/EYE IRRITATION	13
10.6	RESPIRATORY SENSITISATION.....	13
10.7	SKIN SENSITISATION	13
10.8	GERM CELL MUTAGENICITY	13
10.9	CARCINOGENICITY	13
10.10	REPRODUCTIVE TOXICITY.....	13
10.10.1	<i>Adverse effects on sexual function and fertility.....</i>	<i>13</i>
10.10.2	<i>Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility.....</i>	<i>14</i>
10.10.2.1	<i>Summary of effects on fertility.....</i>	<i>14</i>
10.10.3	<i>Comparison with the CLP criteria</i>	<i>18</i>
10.10.4	<i>Adverse effects on development.....</i>	<i>18</i>
10.10.5	<i>Short summary and overall relevance of the provided information on adverse effects on development</i>	<i>21</i>
10.10.6	<i>Comparison with the CLP criteria</i>	<i>28</i>
10.10.7	<i>Adverse effects on or via lactation</i>	<i>30</i>
10.10.8	<i>Short summary and overall relevance of the provided information on effects on or via lactation</i>	<i>30</i>
10.10.9	<i>Comparison with the CLP criteria</i>	<i>30</i>
10.10.10	<i>Conclusion on classification and labelling for reproductive toxicity.....</i>	<i>30</i>
10.11	SPECIFIC TARGET ORGAN TOXICITY-SINGLE EXPOSURE	34
10.12	SPECIFIC TARGET ORGAN TOXICITY-REPEATED EXPOSURE	34
10.12.1	<i>Short summary and overall relevance of the provided information on specific target organ toxicity – repeated exposure.....</i>	<i>35</i>
10.12.2	<i>Comparison with the CLP criteria</i>	<i>37</i>
10.12.3	<i>Conclusion on classification and labelling for STOT RE</i>	<i>37</i>

10.13	ASPIRATION HAZARD.....	38
11	EVALUATION OF ENVIRONMENTAL HAZARDS.....	39
11.1	RAPID DEGRADABILITY OF ORGANIC SUBSTANCES	39
11.1.1	<i>Ready biodegradability</i>	39
11.1.2	<i>BOD₅/COD</i>	40
11.1.3	<i>Hydrolysis</i>	40
11.1.4	<i>Other convincing scientific evidence</i>	40
11.1.4.1	Field investigations and monitoring data (if relevant for C&L).....	40
11.1.4.2	Inherent and enhanced ready biodegradability tests.....	40
11.1.4.3	Water, water-sediment and soil degradation data (including simulation studies)	40
11.1.4.4	Photochemical degradation.....	40
11.2	ENVIRONMENTAL TRANSFORMATION OF METALS OR INORGANIC METALS COMPOUNDS.....	40
11.2.1	<i>Summary of data/information on environmental transformation</i>	40
11.3	ENVIRONMENTAL FATE AND OTHER RELEVANT INFORMATION.....	40
11.4	BIOACCUMULATION	41
11.4.1	<i>Estimated bioaccumulation</i>	41
11.4.2	<i>Measured partition coefficient and bioaccumulation test data</i>	41
11.5	ACUTE AQUATIC HAZARD.....	41
11.5.1	<i>Acute (short-term) toxicity to fish</i>	42
11.5.2	<i>Acute (short-term) toxicity to aquatic invertebrates</i>	43
11.5.3	<i>Acute (short-term) toxicity to algae or other aquatic plants</i>	43
11.5.4	<i>Acute (short-term) toxicity to other aquatic organisms</i>	43
11.6	LONG-TERM AQUATIC HAZARD	44
11.6.1	<i>Chronic toxicity to fish</i>	44
11.6.2	<i>Chronic toxicity to aquatic invertebrates</i>	44
11.6.3	<i>Chronic toxicity to algae or other aquatic plants</i>	44
11.6.4	<i>Chronic toxicity to other aquatic organisms</i>	45
11.7	COMPARISON WITH THE CLP CRITERIA	45
11.7.1	<i>Acute aquatic hazard</i>	45
11.7.2	<i>Long-term aquatic hazard (including bioaccumulation potential and degradation)</i>	46
11.8	CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS.....	46
12	EVALUATION OF ADDITIONAL HAZARDS.....	46
13	REFERENCES.....	46
14	ANNEXES.....	50

Tables

Table 1: Substance identity and information related to molecular and structural formula of the substance	1
Table 2: Constituents (non-confidential information)	2
Table 3: Impurities (non-confidential information) if relevant for the classification of the substance	2
Table 4: Additives (non-confidential information) if relevant for the classification of the substance	2
Table 5: Test substances (non-confidential information) (this table is optional)	2
Table 6: Proposed harmonised classification and labelling according to the CLP criteria	3
Table 7: Reason for not proposing harmonised classification and status under public consultation	4
Table 8: Summary of physicochemical properties	7
Table 9: Summary table of toxicokinetic studies	10
Table 10: Summary table of animal studies on adverse effects on sexual function and fertility	13
Table 11: Summary table of animal studies on adverse effects on development	18
Table 12: Summary table of human data on adverse effects on development	20
Table 13: Summary table of other studies relevant for developmental toxicity	20
Table 14: Summary of Dose-Response: DOTI/MOTI 80:20 vs DOTE in Developmental Toxicity Study with Mice. This is a dose-to-dose comparison stepping down from highest to lowest dose.	32
Table 15: Summary of Dose-Response: DOTI/MOTI 80:20 vs DOTE in Developmental Toxicity Study with Rabbits. This is a dose-to-dose comparison stepping down from highest to lowest dose.	33
Table 16: Summary table of animal studies on STOT RE	34
Table 17: Summary table of human data on STOT RE	35
Table 32: Summary of relevant information on rapid degradability	39
Table 33: Summary of relevant information on bioaccumulation	41
Table 34: Summary of relevant information on acute aquatic toxicity	41
Table 35: Summary of relevant information on chronic aquatic toxicity	44
Table 36: Comparison with criteria for acute aquatic hazards	45
Table 37: Comparison with criteria for long-term aquatic hazards	46

1 IDENTITY OF THE SUBSTANCE

1.1 Name and other identifiers of the substance

Table 1: Substance identity and information related to molecular and structural formula of the substance

Name(s) in the IUPAC nomenclature or other international chemical name(s)	8-Oxa-3,5-dithia-4-stannatetradecanoic acid, 10-ethyl-4,4-dioctyl-7-oxo-, 2-ethylhexyl ester
Other names (usual name, trade name, abbreviation)	Diocyltin bis(2-ethylhexyl)thioglycolate; Diocyltin bis(2-ethylhexyl)mercaptoacetate; DOTE; DOT(EHMA) ₂
EC number (if available and appropriate)	239-622-4
EC name (if available and appropriate)	2-ethylhexyl 10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate
CAS number (if available)	15571-58-1
Other identity code (if available)	
Molecular formula	C ₃₆ H ₇₂ O ₄ S ₂ Sn
Structural formula	
SMILES notation (if available)	<chem>O=C(CS[Sn](SCC(=O)OCC(CC)CCCC)(CCCCCCCC)CCCCCCCC)OCC(CC)CCCC</chem>
Molecular weight or molecular weight range	751.79 g/mol
Information on optical activity and typical ratio of (stereo) isomers (if applicable and appropriate)	The molecule contains two chiral carbon atoms in the 2-ethylhexyl substituent of the acetic ester group. The substance is not known to show optical activity. It is unlikely that one of the possible enantiomers shows different reactivity.
Degree of purity (%) (if relevant for the entry in Annex VI)	> 90 % - < 99.9 %

1.2 Composition of the substance

Table 2: Constituents (non-confidential information)

Constituent (Name and numerical identifier)	Concentration range (% w/w minimum and maximum in multi- constituent substances)	Current CLH in Annex VI Table 3.1 (CLP)	Current self- classification and labelling (CLP)
2-Ethylhexyl-10-ethyl-4,4- dioctyl-7-oxo-8-oxa-3,5- dithia-4- stannatetradecanoate EC no.: 239-622-4 CAS No 15571-58-1	> 90 % / <99.9 % (w/w)	Repr. 1B H360D GHS08 Dgr	Acute Tox. 4 (H302) Skin Sens. 1A (H317) Repr. 1B (H360d) STOT RE 1 (H372) Aquatic Acute 1 (H400) Aquatic Chronic 1 (H410)

Table 3: Impurities (non-confidential information) if relevant for the classification of the substance

Impurity (Name and numerical identifier)	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self- classification and labelling (CLP)	The impurity contributes to the classification and labelling
see conf. Annex				

Table 4: Additives (non-confidential information) if relevant for the classification of the substance

Additive (Name and numerical identifier)	Function	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self- classification and labelling (CLP)	The additive contributes to the classification and labelling
-					

Table 5: Test substances (non-confidential information) (this table is optional)

Identification of test substance	Purity	Impurities and additives (identity, %, classification if available)	Other information	The study(ies) in which the test substance is used
-				

2 PROPOSED HARMONISED CLASSIFICATION AND LABELLING

2.1 Proposed harmonised classification and labelling according to the CLP criteria

Table 6: Proposed harmonised classification and labelling according to the CLP criteria

	Index No	International Chemical Identification	EC No	CAS No	Classification		Labelling			Specific Conc. Limits, M-factors	Notes
					Hazard Class and Category Code(s)	Hazard statement Code(s)	Pictogram, Signal Word Code(s)	Hazard statement Code(s)	Suppl. Hazard statement Code(s)		
Current Annex VI entry	050-027-00-7	2-ethylhexyl 10-ethyl-4,4- dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate	239-622-4	15571-58-1	Repr. 1B	H360D	GHS08 Dgr	H360D			
Dossier submitters proposal					Modify Repr. 2	Modify H361d	GHS08 GHS09 Dgr	H361d H372 (thymus) H411			
Resulting Annex VI entry if agreed by RAC and COM					Repr. 2 STOT RE 1 Aquatic Chronic 2	H361d H372 (thymus) H411	GHS08 GHS09 Dgr	H361d H372 (thymus) H411			

Table 7: Reason for not proposing harmonised classification and status under public consultation

Hazard class	Reason for no classification	Within the scope of public consultation
Explosives	<i>hazard class not assessed in this dossier</i>	No
Flammable gases (including chemically unstable gases)	<i>hazard class not assessed in this dossier</i>	No
Oxidising gases	<i>hazard class not assessed in this dossier</i>	No
Gases under pressure	<i>hazard class not assessed in this dossier</i>	No
Flammable liquids	<i>hazard class not assessed in this dossier</i>	No
Flammable solids	<i>hazard class not assessed in this dossier</i>	No
Self-reactive substances	<i>hazard class not assessed in this dossier</i>	No
Pyrophoric liquids	<i>hazard class not assessed in this dossier</i>	No
Pyrophoric solids	<i>hazard class not assessed in this dossier</i>	No
Self-heating substances	<i>hazard class not assessed in this dossier</i>	No
Substances which in contact with water emit flammable gases	<i>hazard class not assessed in this dossier</i>	No
Oxidising liquids	<i>hazard class not assessed in this dossier</i>	No
Oxidising solids	<i>hazard class not assessed in this dossier</i>	No
Organic peroxides	<i>hazard class not assessed in this dossier</i>	No
Corrosive to metals	<i>hazard class not assessed in this dossier</i>	No
Acute toxicity via oral route	<i>hazard class not assessed in this dossier</i>	No
Acute toxicity via dermal route	<i>hazard class not assessed in this dossier</i>	No
Acute toxicity via inhalation route	<i>hazard class not assessed in this dossier</i>	No
Skin corrosion/irritation	<i>hazard class not assessed in this dossier</i>	No
Serious eye damage/eye irritation	<i>hazard class not assessed in this dossier</i>	No
Respiratory sensitisation	<i>hazard class not assessed in this dossier</i>	No
Skin sensitisation	<i>hazard class not assessed in this dossier</i>	No
Germ cell mutagenicity	<i>hazard class not assessed in this dossier</i>	No
Carcinogenicity	<i>hazard class not assessed in this dossier</i>	No
Reproductive toxicity	harmonised classification proposed	Yes
Specific target organ toxicity-single exposure	<i>hazard class not assessed in this dossier</i>	No
Specific target organ toxicity-repeated exposure	harmonised classification proposed	Yes
Aspiration hazard	<i>hazard class not assessed in this dossier</i>	No
Hazardous to the aquatic environment	harmonised classification proposed	Yes
Hazardous to the ozone layer	<i>hazard class not assessed in this dossier</i>	No

3 HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING

		CLP Regulation (EC) No1272/2008
2 October 2013	944/2013/EU, Amendment of Annex VI, Tables 3.1 and 3.2 CLP as follows	Repr 1B H360D GHS08 Dgr
8 June 2012	RAC adopted opinion that DOTE should be classified and labelled as follows	Repr 1B H360D GHS08 Dgr
25 March 2011	Arkema on behalf of ETINSA submitted CLH dossier with the following proposed classification.	Repr. 2 H361d

4 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

Reason for a need for action at Community level:

- New data
- Differences in self-classification (STOT RE, Aquatic Toxicity)

Further detail on need of action at Community level

According to Article 36(1), a substance that fulfils the criteria set out in Annex I of the CLP regulation for the following shall normally be subject to harmonized classification and labelling in accordance with Article 37:

(d) Reproductive toxicity, Category 1A, 1B or 2 (Annex I, section 3.7).

According to Article 37(6), manufacturers, importers and downstream users who have new information which may lead to a change of the harmonised classification and labelling elements of a substance in Part 3 of Annex VI shall submit a proposal in accordance with the second subparagraph of paragraph 2 to the competent authority in one of the Member States (MSCA) in which the substance is placed on the market.

CLH dossiers proposing a revision of a specific hazard class from an existing entry in Annex VI to the CLP Regulation can only be submitted directly to ECHA by an MSCA (Article 37(1), CLP). A revision of an existing entry can be justified in the event that new data has become available since the harmonised classification was agreed. The new data could, for example show that classification in a different category is justified.

Currently the harmonized classification of DOTE by the Commission Regulation (EU) No 944/2013 is Repr. 1B (H360D). The RAC assessment and the resulting opinion from 2012 are based solely on read across data.

Two new key studies are available which investigated the developmental toxicity with the DOTE substance itself. These new GLP studies complement earlier flawed studies with the most sensitive species, mice and rabbits. The new studies were conducted with 96 % purity DOTE and are fully complete guideline studies (OECD 414) following GLP which evaluated gestational integrity, external anomalies, soft tissue anomalies and skeletal anomalies.

Recent *in vitro* hydrolysis studies at gastric pH value of 1.2 and pH values of 9, 7 and 4, using ¹¹⁹Sr-NMR to identify the reaction products, clearly show that Dioctyltinchloro 2-ethylhexylmercaptoacetate (DOTEC) is the only identifiable hydrolysis product of Dioctyltin bis(2-ethylhexylmercaptoacetate) (DOTE). No Dioctyltin dichloride (DOTC) could be detected under the conditions of the study.

Thus, industry found that the read across from DOTC to DOTE is no longer considered appropriate without restrictions. To their view the studies with DOTC (since DOTC is not formed during acidic

hydrolysis) may have low significance for the hazard assessment and studies with dioctyltin bis(isooctyl mercaptoacetate)/Monooctyltin tris(isooctyl mercaptoacetate) 80:20 (DOTI/MOTI 80:20) on mice and rabbits and lower significance for the hazard assessment than studies with DOTE.

This CLH report was initially drafted by Industry (proposing no classification on reproductive toxicity) and modified by the German Competent Authority according to their evaluation of the data.

5 IDENTIFIED USES

DOTE is mostly used as a stabiliser in plastic. The identified uses of the substance cover uses at industrial sites.

6 DATA SOURCES

REACH dossiers (12/2016)

7 PHYSICOCHEMICAL PROPERTIES

Table 8: Summary of physicochemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
Physical state at 20°C and 101,3 kPa	Liquid, clear colourless to slightly yellow	Baltussen E, 2010, Determination of physico-chemical properties of dioctyltin bis(2-ethylhexylmercaptoacetate), NOTOX B.V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands	
Melting/freezing point	-39 °C	Elf Atochem NA. 1999. Material Safety Data Sheet. Thermolite (R) 890 Stabilizer. Revision 2. Issued 31 August 1999.	
Boiling point	275 °C	Baltussen E, 2010, determination of the boiling temperature of DOTE by differential scanning calorimetry, NOTOX BV, Hambakenwetering 7, 5231 DD Hertogenbosch, NL	The substance decomposes at T >275°C and normal pressure without boiling.
Relative density	1.07 g/cm ³ at 20 °C	Elf Atochem NA. 1999. Material Safety Data Sheet. Thermolite (R) 890 Stabilizer. Revision 2. Issued 31 August 1999.	
Vapour pressure	< 2.50 x 10 ⁻⁴ Pa	Baltussen E, 2010, Determination of physico-chemical properties of dioctyltin bis(2-ethylhexylmercaptoacetate), NOTOX B.V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands	Due to the behaviour of the test material in the equipment, an exact value for the vapour pressure could not be calculated. Three tests were performed. Significant differences between the individual measurements were observed. The vapour pressure was therefore reported to be lower than the highest measured value at < 2.50 x 10 ⁻⁴ Pa
Surface tension			not technically feasible as the water solubility of the substance is less than 0.1 mg/l.
Water solubility	0.000000001 µg/l	EPI-QSAR	study technically not feasible
Partition coefficient n-octanol/water	Log Pow = 15.34	EPI-QSAR	study technically not feasible
Flash point	182 °C	Baltussen E, 2010, Determination of physico-chemical properties of dioctyltin bis(2-	Pensky-Martens closed cup method.

Property	Value	Reference	Comment (e.g. measured or estimated)
		ethylhexylmercaptoacetate), NOTOX B.V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands	
Flammability	Not flammable		
Explosive properties	Not explosive		Expert judgement based on physico-chemical properties and the substance's structure
Self-ignition temperature	390 °C at 989.6 -999.2 hPa.	Baltussen E, 2010, Determination of physico- chemical properties of dioctyltin bis(2- ethylhexylmercaptoacetate), NOTOX B.V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands	
Oxidising properties	No oxidising properties		Expert judgement based on physico-chemical properties and the substance's structure
Granulometry	Not relevant		Substance is liquid
Stability in organic solvents and identity of relevant degradation products	Substance is stable in apolar solvents		
Dissociation constant			Technically not feasible
Viscosity	57.9 mm ² /s at 20 °C 24.0 mm ² /s at 40 °C	Baltussen E, 2010, Determination of the kinematic viscosity of dioctyltin bis (2- ethylhexylmercaptoacetate), NOTOX B.V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands	

The following QSAR estimations were done for water solubility and Log Pow:

Water Sol: 2.94e-011 mg/L

SMILES : O=C(CS[Sn](SCC(=O)OCC(CC)CCCC)(CCCCCCCC)CCCCCCCC)OCC(CC)CCCC

CHEM: MOL FOR: C36 H72 O4 S2 Sn1 MOL WT: 751.80

WSKOW v1.42 Results: Log Kow (estimated): 15.35 Log Kow (experimental): not available from
database Log Kow used by Water solubility estimates: 15.35 Equation Used to Make Water Sol estimate:

Log S (mol/L) = 0.693-0.96 log Kow-0.0092(Tm-25)-0.00314 MW + Correction Melting Pt (Tm) = -39.00
deg C (Use Tm = 25 for all liquids) Correction(s): Value

No Applicable Correction Factors Log Water Solubility (in moles/L): -16.408

Water Solubility at 25 deg C (mg/L): 2.94e-011

Log Kow(version 1.68 estimate): 15.35

SMILES : O=C(CS[Sn](SCC(=O)OCC(CC)CCCC)(CCCCCCCC)CCCCCCCC)OCC(CC)CCCC

CHEM: DOTE MOL FOR: C36 H72 O4 S2 Sn1 MOL WT: 751.80

CLH REPORT FOR DOTE

TYPE	NUM	LOGKOW FRAGMENT DESCRIPTION	COEFF	VALUE
Frag	6	-CH3 [aliphatic carbon]	0.5473	3.2838
Frag	26	-CH2- [aliphatic carbon]	0.4911	12.7686
Frag	2	-CH [aliphatic carbon]	0.3614	0.7228
Frag	2	-C(=O)O [ester, aliphatic attach]	-0.9505	-1.9010
Frag	2	-S- [aliphatic attach]	-0.4045	-0.8090
Frag	1	Tin [Sn]	1.0600	1.0600
Const	-	Equation Constant	-	0.2290

Log Kow = 15.3542

8 EVALUATION OF PHYSICAL HAZARDS

Physical hazards are not evaluated in this dossier

9 TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)

Table 9: Summary table of toxicokinetic studies

Method	Results	Remarks	Reference
<i>in vitro</i> study DOTE- Hydrolysis as a Function of pH (OECD TG 111)	The study showed that DOTE at pH 9, 7 and 4 can be considered hydrolytically stable. After 5 days at 50 °C less than 10 % DOTE was hydrolyzed ($t_{0.5}$ 25 °C > 1 year). Under the simulated gastric conditions (0.1 M HCl / pH 1.2 / 37 °C, 5 days) DOTE was hydrolyzed to DOTE _C , its monochloride ester.	2 (reliable with restrictions) key study experimental result Test material (EC name): 2-ethylhexyl 10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate	Anonymous (2015)
<i>in vitro</i> study rat and human epidermis dermal Exposure regime: 24 hour(s) Doses/conc.: 17,007 µg tin/cm ² OECD Draft Guideline for Dermal Delivery and Percutaneous Absorption: In Vitro Method [OECD TG 428]	Main ADME results: Absorption: Absorption of tin from DOTE through rat epidermis significantly overestimates absorption through human epidermis. Evaluation of results: bioaccumulation potential cannot be judged based on study results	2 (reliable with restrictions) key study experimental result Test material: DOTE (EC name): 2-ethylhexyl 10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate	Ward, R.J. (2003)

9.1 Short summary and overall relevance of the provided toxicokinetic information on the proposed classification(s)

Hydrolysis:

A new *in vitro* study simulating mammalian gastric conditions, using 119-Sn-NMR spectroscopy to identify the reaction products, clearly shows that Dioctyltin chloro 2-ethylhexyl mercaptoacetate (DOTE_C) is the only identifiable hydrolysis product of DOTE (Dioctyltin bis(2-ethylhexyl mercaptoacetate)). No Dioctyltin dichloride (DOTC) could be detected under the conditions of the study.

This *in vitro* study can be assumed to reasonably predict the behaviour of DOTE within the gastric contents, but it does not elucidate the full mode-of-action for oral toxicity studies because neither the full spectrum of possible *in vivo* metabolites, nor the toxicologically active species *in vivo*, are known. No definitive conclusion can be drawn for the mode-of-action from this study for repeated dose, *in vivo* genotoxicity, reproduction, and developmental effects, when they are assessed using oral administration.

The *in vitro* study leaves open the question whether any DOTE or DOTE_C formed in the gastric environment might be taken up systemically and then be further metabolised into other dioctyltin species, monoctyltin species or non-organotin species. However, in the absence of any additional data read across from studies using DOTC as test material, must be considered carefully in the discussion of the reproductive toxicity endpoints.

The same study showed that DOTE at pH 9, 7 and 4 can be considered hydrolytically stable. After 5 days at 50 °C less than 10 % DOTE was hydrolyzed ($t_{0.5}$ 25 °C > 1 year). However the study only considers abiotic conditions. The situation in an organism where metabolizing enzymes are present might be different.

Inhalation and dermal absorption:

With respect to inhalation and dermal mammalian toxicity, the thioesters have much higher molecular weights and considerably lower volatility than the chloride. The high molecular weights of the esters reduce their potential for absorption via the dermal route, and their volatility reduces their potential for absorption via the inhalation route relative to the chloride.

The absorption of DOTE was measured *in vitro* (Ward 2003) through both occluded and unoccluded human and rat epidermis. The absorption through rat epidermis was much faster than through human epidermis:

HUMAN EPIDERMIS: A dose of undiluted liquid DOTE, corresponding to 17,007 $\mu\text{g tin}/\text{cm}^2$ was determined to slightly reduce the measured electrical resistance across rat skin. Electrical resistance is one indicator of the integrity of the barrier function of the epidermis. The measured reduction was minimal [3.13 ohms versus ≤ 3.00 ohms indicative of undamaged skin]. Because human skin is typically more robust than rat skin, the authors chose to continue with the 17,007 $\mu\text{g tin}/\text{cm}^2$ dose, which was the highest dermal dose achievable, and both the rat and the human skin samples were judged to be entirely adequate for the integrity of the dermal penetration test.

From the occluded and unoccluded applications, the rates of tin absorption over the 0-24 h exposure period were below the limit of quantification (0.001 $\mu\text{g}/\text{cm}^2/\text{h}$). In terms of percent applied tin, 0.0001 % was absorbed from the occluded dose, and 0.0001 % was absorbed from the unoccluded dose after 24 hours of exposure.

RAT EPIDERMIS: Absorption of tin through rat epidermis was much faster than through human epidermis. From the occluded application, the maximum rate of tin absorption (0.035 $\mu\text{g}/\text{cm}^2/\text{h}$) occurred during 16-24 hours of exposure, and the mean rate of tin absorption over the whole 24-h exposure period was 0.021 $\mu\text{g}/\text{cm}^2/\text{h}$. From the unoccluded application, the maximum rate of tin absorption occurred during 12-24 hours of exposure and was 0.033 $\mu\text{g}/\text{cm}^2/\text{h}$. The mean rate of tin absorption over the whole 24-h exposure period was 0.025 $\mu\text{g}/\text{cm}^2/\text{h}$. In terms of percent applied tin, 0.003 % was absorbed from the occluded dose, and 0.004 % was absorbed from the unoccluded dose after 24 hours of exposure. The overall recovery of tin from the test system after 24-h exposure was low and may be due to adsorption of the test substance to the glass equipment used. The recovery was 45.5 % (human) and 25.2 % (rat) of the applied occluded doses, and 29.6 % (human) and 30.5 % (rat) were recovered from the unoccluded test systems. Of the recovered tin, 2.1 % (human) and 5.5 % (rat) were obtained from the surface of the epidermis and donor chamber. The mean amounts of tin absorbed by 24 hours were 0.010 $\mu\text{g}/\text{cm}^2$ (unoccluded) and 0.011 $\mu\text{g}/\text{cm}^2$ (occluded) through human epidermis and 0.641 $\mu\text{g}/\text{cm}^2$ (unoccluded) and 0.547 $\mu\text{g}/\text{cm}^2$ (occluded) through rat epidermis.

These results show that the absorption of tin from DOTE through rat epidermis significantly overestimated the absorption from human epidermis. However, by 24 hours only a small amount of the applied tin (3 % in human and 1 % in the rat) is associated with the epidermis and is not regarded as systemically available. Thus, based on the low recovery the reliability of the study is highly questionable.

10 EVALUATION OF HEALTH HAZARDS

Read across approach for repeated exposure assessment:

The CLH dossier of 25 March 2011 and the RAC opinion of 8 June 2012 applied read across among three chemicals:

- DOTE
- DOTI
- DOTC

These substances are all members of the dioctyltin family of compounds, and the read across characteristics for this family were discussed in depth under the HPV program: SIDS Initial Assessment Reports "Dioctyltin

dichloride and selected thioesters". The dioctyltins (DOT) are tetravalent tin compounds comprised of two octyl groups bound to tin through tin-carbon bonds, and two other groups bound to tin.

The rationale for the read across was based on a simulated gastric hydrolysis study proposing that DOTE (dioctyltin bis(2-ethylhexylmercaptoacetate)) would readily hydrolyse under physiological conditions to DOTC (dioctyltin dichloride). Thus DOTC was selected as an anchor compound and surrogate for the mammalian toxicology endpoints of repeated dose, *in vivo* genetic toxicity, reproduction, and developmental effects, when they are assessed using oral administration.

An inherent uncertainty exists in this approach due to the fact that the resulting tin species cannot be analysed and characterized as such by gas chromatography. The analytically necessary derivatisation modifies the chemical structure into tetraalkyltin derivatives, which are available to the gas phase. Thus a distinction between different ligands bound to the dioctyltin moiety during the simulated gastric hydrolysis is no longer possible.

Recent *in vitro* hydrolysis studies under simulated mammalian gastric conditions, using 119-Sn-NMR to identify the reaction products, clearly show that dioctyltinchloro 2-ethylhexylmercaptoacetate (DOTE) is the only identifiable hydrolysis product. No dioctyltin dichloride (DOTC) could be detected under the conditions of the study.

Recent *in vitro* hydrolysis studies at gastric pH value of 1.2 and pH values of 9, 7 and 4, using 119-Sn-NMR to identify the reaction products, show that dioctyltinchloro 2-ethylhexylmercaptoacetate (DOTE) is the only identifiable hydrolysis product. No dioctyltin dichloride (DOTC) could be detected under the conditions of the study (detection limit 0.5 % on pure DOTC).

However, the study does not allow to conclude, if and which other dioctyl species might be formed after systemic uptake of DOTE and/or DOTE) under *in vivo* conditions. Thus, read across to DOTC cannot be disregarded based on the results of this hydrolysis study. In addition to the read across to DOTC, the CLH dossier of 25 March 2011 and the RAC opinion of 8 June 2012 used a read across from the structural analogue substance DOTI, diisooctyl 2,2'-[(dioctylstannylene)bis(thio)]diacetate (CAS No. 26401-97-8).

Furthermore, read across at an "analogue level" as described in the SIDS report (SIDS Initial Assessment Report "Esters of Thioglycolic Acid" prepared for SIAM 23 (2006)) was applied to data on diisooctyl 2,2'-[(dioctylstannylene)bis(thio)]diacetate (CAS No. 26401-97-8, also named dioctyltin bis(isooctylmercaptoacetate), dioctyltin bis(IOMA), DOTI). DOTI and DOTE are isomers differing only slightly in the structure of the C-8 alcohol of the mercaptoester ligand (either *iso*-octanol or 2-ethylhexanol, respectively). Since these alcohols are so close in structure, their respective mercaptoacetate esters are expected to have very similar physicochemical and toxicological properties, including hydrolysis products.

The recently conducted developmental toxicity studies with DOTE in two species may have higher significance than those with DOTI for reproduction hazard assessment. Therefore the significance of any effects read across from DOTI to DOTE is regarded in a weight-of-evidence assessment and the significance weighted accordingly (see Summary Tables of Dose-Response (Table 14 and Table 15) in Chapter 10.10.10, which suggest that DOTI might have a higher toxicological potency with respect to reproductive toxicity).

10.1 Acute toxicity - oral route

This point is not proposed for harmonization.

10.2 Acute toxicity - dermal route

This point is not proposed for harmonization.

10.3 Acute toxicity - inhalation route

Not evaluated in this dossier.

10.4 Skin corrosion/irritation

This point is not proposed for harmonisation.

10.5 Serious eye damage/eye irritation

This point is not proposed for harmonisation.

10.6 Respiratory sensitisation

Not evaluated in this dossier.

10.7 Skin sensitisation

This point is not proposed for harmonisation.

10.8 Germ cell mutagenicity

This point is not proposed for harmonisation.

10.9 Carcinogenicity

Not evaluated in this dossier

10.10 Reproductive toxicity**10.10.1 Adverse effects on sexual function and fertility**

Table 10: Summary table of animal studies on adverse effects on sexual function and fertility

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
Method: OECD Guideline 416 (Two-Generation Reproduction Toxicity Study) GLP key study 2 (reliable with restrictions) Rat (Sprague-Dawley) 25 male/25 female per group two-generation study oral: feed	Test material: DOTI/MOTI 80:20 In original study named: Dioctyltin bis(IOMA) [CAS no. 26401-97-8] : Octyltin tris(IOMA) [CAS no. 26401-86-5] (purity 78.8 : 16.9 %) 0, 20, 60, and 200 ppm (nominal in diet) (P: ~1.5, 4.4, 15 mg test material/kg bw/d) (F1: ~1.6, 4.7, 16 mg test material/kg bw/d) Exposure: Duration of dosing of F0 generation males - 10 weeks prior to mating, during mating (3 weeks), and post mating until sacrifice; females - 10 weeks prior to mating and during mating. Mated females continued to receive test diets during gestation and lactation; unmated females received test diets until sacrifice. Test diets were prepared weekly	NOAEL (P): 20 ppm based on test material (~1.5 mg/kg bw/d male/female) (based on a reduction in the relative thymus weight of males) NOAEL (F1): 20 ppm based on test material (male/female) (The NOAEL for the F1 generation until weaning was 20 ppm (~1.6 mg/kg bw/d), based on a decrease in relative thymus weights in male and female pups at 60 ppm (~4.7 mg/kg bw/d). The NOAEL for the F1 generation post lactation was 20 ppm (~1.6 mg/kg bw/d, based on a slight decrease in the relative thymus weight of males and an increase in stillbirths at 60 ppm (~4.7 mg/kg bw/d). NOAEL (reproductive organs, fertility, gestational integrity) 200 ppm based on test material. No effect on reproductive organs or	Anonymous (1997)

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
	and analysed for homogeneity and stability. Duration of dosing of F1 generation: males - 14 weeks (starting at the end of lactation prior to mating), during mating (3 weeks), and post mating until sacrifice; females - 14 weeks (starting at the end of lactation prior to mating) and during mating (3 weeks). (continuously (in diet))	reproductive capacity was observed up to and including the highest dose tested. NOAEL (teratogenicity): 200 ppm based on test material (No teratogenic effect was observed within the limits of the experimental design up to and including the highest dose tested)	

10.10.2 Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility

10.10.2.1 Summary of effects on fertility

In the two generation study performed under GLP and according to OECD 416 (Anonymous, 1997), the test material consisting of 78.8 % DOTI (CAS No. 26401-97-8) and 16.9 % MOTI (Mono-n-octyltin tris(2-ethylhexylmercaptoacetate), CAS No. 26401-86-5, also named Octyltin tris(IOMA)) was administered to the F0 generation 10 weeks prior to mating, during mating (3 weeks) and post-mating. DOTI and DOTE are isomers, they are expected to have similar reactivity and to be toxicologically equivalent. However, the recently conducted developmental toxicity studies with DOTE in two species (see Table 11, Section 10.10.4) support that DOTI might have a higher toxicological potency with respect to reproductive toxicity (see Summary Tables of Dose-Response (Table 14 and Table 15) in Chapter 10.10.10).

The F₁ generation was treated 14 weeks during pre-mating, 3 weeks during mating. Females continued to receive the test material during gestation and lactation.

The following treatment-related effects were observed:

P generation:

- Mortality: 1 male died at 200 ppm test material in the diet
- No substance-related mortality or changes in behaviour or external appearance
- Absolute food consumption reduced in females at 200 ppm (-6 % on lactation days 7-14, -9 % on lactation days 14-21)
- No substance-related changes in Mean pre-coital time, Mating Index, Pregnancy Rate, Fertility Index, Gestation Index, Mean pregnancy duration, live-born pups, or stillbirths
- Viability index slightly increased at 200 ppm (96.2 % in the controls vs. 98.6 % s).
- Lactation index significantly decreased at 200 ppm diet (88.6 % vs. 94.4 % in controls, $p < 0.05$) after 21 days lactation.
- Pup body weights not significantly decreased (approx. -3-4 % at 200 ppm) at birth, but significantly decreased at 200 ppm in both sexes after 14 and 21 days lactation (-19 to -21 %, $p < 0.01$; male: 34.58 g vs. 42.59 g for controls; female: 33.54 g vs. 42.27 g for controls).
- Slight delay in vaginal opening at 200 ppm (35.4 days vs. 33.2 days for controls).
- Slight decrease in relative thymus weight in males at 60 ppm (approx. 4.4 mg test material/kg bw/day); significant decrease in relative thymus weight in both sexes at 200 ppm.
- Increased incidence of thymic involution at 200 ppm (1/25 for control males; 9/25 for 200 ppm males; 0/25 for control females; 4/25 for 200 ppm females; significant for males only) at microscopic examination.
- Functional tests and examination of morphological landmarks revealed no substance-related findings at all dose-levels except for a slightly delayed in vaginal opening at 200 ppm.

- Microscopic examination of the other organs found no substance-related changes.

F₁ generation:

- No mortality.
- Body weight: significant reduction in F₁ males during their growth phase at 200 ppm [89 % of controls on test week 1 (TW1); 90 % of controls TW8; 91 % of controls TW22].
- Food consumption: generally within historical control values for both sexes with occasional statistical differences; not reduced in females during gestation, but reduced at 200 ppm and significant on lactation days 14-21 [-12 % for LD1-7, -15 % for LD7-14, -17 % for LD14-21].
- No substance-related changes in Mean pre-coital time, Mating Index, Pregnancy Rate, Fertility Index, Gestation Index, Mean pregnancy duration, or live-born pups
- Statistically significant increase in number of stillbirths at 200 ppm (26 vs. 5 in controls).
- Viability index: decreased at 200 ppm (82.0 % vs. 95.7 % in controls).
- Pup mortality: no significant differences from birth to PND4 prior to culling; significantly increased mortality at 200 ppm from PND4 after culling through PND21.
- Lactation index: decreased at 200 ppm (82.3 % vs. 94.4 %).
- Pup body weight for F₂ generation: reduced at 200 ppm; no significant differences at birth; significantly reduced at 200 ppm for males and females on PND 4, significantly reduced at 200 ppm for females on PND 7, 14, and 21 (male pups between approx. 3 % and 19 %; female pups between approx. 4 % and 21 %, $p < 0.01$).
- Morphological changes for F₂ generation: pinna unfolding, eye and ear opening were slightly, but not significantly, delayed (approx. 0.5 days) at 200 ppm, likely related to reduced post-natal body weight.
- Relative thymus weight of F₁ generation: showed a tendency towards a decrease [approx. 86 % of controls] in male and female rats at 60 ppm (approx. 4.7 mg test material/kg bw/day) and was significantly decreased in both sexes [males: 78 % of controls; females: 61 % of controls] at 200 ppm ($p < 0.01$).
- Relative thymus weight of F₂ generation: no significant differences in male or female rat pups at any dose on PND22.
- Relative spleen weight of F₁ generation: no significant differences for either sex at 60 or 200 ppm.
- Increased incidence of thymic involution for the F₁ generation at 200 ppm based on test material in feed (4/25 for control males; 13/25 for 200 ppm males; 1/25 for control females; 2/23 for 200 ppm females; significant for males) at microscopic examination.

The NOAEL for P males and females was 20 ppm (approx. 1.5 mg test material/kg bw/day) based on a slightly reduced relative thymus weight for males at 60 ppm (approx. 4.4 mg test material/kg bw/day).

The NOAEL for the F₁ generation was 20 ppm (approx. 1.6 mg test material/kg bw/day), based on a reduction in relative thymus weights for males and females at 60 ppm (approx. 4.7 mg test material/kg bw/day).

No teratogenic effects were observed in this study.

Interpretation

The body weight of F₁ animals showed no statistically significant differences from the controls in either sex at birth, PND4 (postnatal day 4 both before and after culling) or PND7. Beginning with PND14 there was a statistically significant decrease in both male and female pup body weights which persisted through PND21. This correlates exactly with the incidences of post-natal losses documented for these F₁ animals. Up to PND4 the number of F₁ pups was not different from controls for either sex. Pups were culled to 4/sex/litter on PND4 (a routine practice). After culling, the pup losses across the four dose groups between PND4 and PND21 were 0, 4, 11, and 20 for the control, 20, 60, and 200 ppm doses of DOTI/MOTI in feed, respectively. It may not be a reproductive effect as the effects may occur as a consequence of impaired nursing behaviour as a result of maternal toxicity and/or a consequence of a direct toxic effect of dioctyltin species on young animals who may well be receiving two sources of exposure, one from the maternal milk and a second from the diet as they begin to eat adult feed during the lactation period. The mg/kg bw/day dose

to these juveniles is likely far higher than the dose to the lactating P females (the mothers) and the dose-response of the juvenile deaths supports this conclusion. There is no indication from clinical signs or body weights in females on a significant maternal toxicity that may have caused impaired nursing. Only food consumption was found to be lowered during the lactation period. In addition, the delay in vaginal opening of the high dose females is likely due to the non-specific toxic effect and low weight gain resulting from this postnatal exposure. For the F1 generation, DOTI/MOTI did not produce an adverse effect on mating, fertility, pregnancy rates, gestation, litter size or neonatal body weight. There was an increase in stillbirths which was inexplicably marked both non-statistically significant and statistically significant at the high dose, however 12/26 still births were in a single litter which diminishes any statistical significance in the endpoint analysis. There was an effect on thymus weight of the F2 animals and the histopathology of the thymus was typical of the known target organ effect of dioctyltin substances on this gland.

The body weight of F2 animals at the mid and low doses showed no statistically significant differences from controls in either sex at birth, PND4 (postnatal day 4 both before and after culling) or PND7. The body weight of F2 animals at the high dose showed no statistically significant differences from controls in either sex at birth, however at PND4 (postnatal day 4 both before and after culling) and thereafter through PND21 body weights of both sexes were statistically decreased.

Just as in the F1 generation, the body weight outcome for the F2 animals correlated with the incidences of postnatal losses. Up to PND4 the number of F2 pups was not different from controls for either sex. Pups were again culled to 4/sex/litter on PND4. After culling, the pup losses across the four dose groups between PND4 and PND21 were 9, 26, 29, and 22 for the control, 20, 60, and 200 ppm doses of DOTI/MOTI in feed. This is not a reproductive effect for the F2 generation either. It is the second manifestation of a direct toxic effect of dioctyltin species on young animals who may well be receiving two sources of exposure, one from the maternal milk and a second from the diet as they begin to eat adult feed during the lactation period. As in the F1 pups, the mg/kg/day dose to these F2 juveniles is likely far higher than the dose to the lactating F1 females (the mothers) and the dose-response of the juvenile deaths (taking into account the slightly lower number of pups available in the F2 high dose group) supports this conclusion. The consequences of the significant maternal and paternal toxicity early in the F1 generation at the high dose can be seen to be played out in the manifestation of the reproductive results in the F2 generation. No assessment of vaginal opening or testicular descent was done in the F2 animals because the experiment was terminated at PND21, prior to the occurrence of these developmental landmarks. As with the F1 generation, the effects on the offspring are not reproductive effects as envisioned by the classification criteria.

There are no adverse reproductive effects at any dose of DOTI/MOTI in the F1 generation. The postnatal losses are most probably the consequence of dose-related direct toxic effects of dioctyltin species on the F1 animals beginning after PND4 possibly coupled with impaired nursing behaviour, which is causally related to maternal toxicity. The effects on body weight and survival of the F1 animals at the high dose support a conclusion that there is carry-over toxicity, both maternal and paternal. This is manifested again during the physiological stress of producing the F2 generation pups, which are similarly adversely affected, but only after PND4.

The reproductive NOEL is the high dose; the maternal and paternal NOEL is 20 ppm (the low dose) which is driven by the effect of the DOTI/MOTI test material on the thymus gland. The effect on the PND4-PND21 pups of the F1 and F2 generations should not be used in setting the NOEL because the delivered dose to these animals via lactation is unknown, and likely higher than that of the adults because of dual exposure via milk and diet.

Conclusion for effects on fertility

Under the experimental conditions of the two generation study on DOTI/MOTI 80:20, the NOAEL for the F0 parental generation was 20 ppm (~1.5 mg test material/kg bw/day), based on a reduction in the relative thymus weight of males at 60 ppm (~4.7 mg test material/kg bw/day). The NOAEL for the F1 generation until weaning was 20 ppm (~1.6 mg test material/kg bw/day), based on a decrease in relative thymus weight in male and female pups at 60 ppm. The NOAEL for the F1 generation post-lactation was 20 ppm, based on a slight decrease in the relative thymus weight of males and an increase in stillbirth at 60 ppm.

Indices of mating, fertility, gestation and the pregnancy rates were within the range of the control group at 20 and 60 ppm. The mean pre-coital time, duration of pregnancy in days and duration in hours did not show any substance related effects at all dose-levels. The fertility index was slightly decreased at 200 ppm but was within the range of historical control data. The viability and lactation indices were decreased at 200 ppm in both the F0 and F1 generation, however this was associated with a decrease in pup body weight (by 3 to 4 %) in the F0 generation and a significant decrease in pups weight in the F1 generation (males pups between approx. 3 % and 19 %; female pups between approx. 4 % and 21 %, at $p < 0.01$) during the lactation period that may be unrelated to in utero exposure.

It was concluded by RAC in its 2012 assessment, that the data may not be sufficiently detailed or complete for a comprehensive evaluation for the endpoint fertility.

No effects on the male or female reproductive organs or on fertility, or on reproductive capacity, or gestational integrity were observed in this study. Further, there was no evidence of a teratogenic effect observed at any dose within the limits of the design of this study.

There is a GLP-conform reproductive toxicity screening study according to OECD guideline 421 (Appel and Waalkens, 2004) performed with dioctyltin dichloride (CAS 3542-36-7) and described in detail in section 0. In this GLP study, comparable effects to the two generation study were obtained; indeed thymus effects were also recorded. Dose-related effects were seen at 10, 100 and 300 ppm (corresponding to 0.5-0.7, 4.2-6.2 and 8.4-17.0 mg DOTC/kg bw/d), with post-implantation losses in the top two dose groups. The maternal LOAEL was set at 10 ppm diet (equivalent to 0.7 mg DOTC/kg bw/day for males and 0.5-0.7 mg DOTC/kg bw/day for females) for treatment related effects to dams including moderate to very severe lymphoid depletion in the thymus, which was considered related to treatment.

Lymphoid depletion was characterized by a decrease in the size of the thymic lobules which can be ascribed to extensive loss of cortical and medullary small lymphocytes. Consequently, the distinction between the cortical and medullary area was blurred. Lymphoid depletion was observed in 5/10 of the 10 ppm group and in all animals of the 100 and 300 ppm groups.

One control animal also had very severe lymphoid depletion in the thymus. However, this was most probably associated with the fact this animal was physiologically disturbed, as was demonstrated by 12 resorptions in the uterus and an abnormal kidney (gross change: flabby and yellow patches).

Some animals of the 10 ppm group showed thymic involution as a result of pregnancy/lactation. This appearance was similar to the thymic pregnancy/lactation involution in control animals and was characterized by a decreased size of thymic lobules exhibiting normal architecture. This phenomenon is a common observation in pregnant or lactating animals. However, the lymphoid depletion in the animals of the 10 ppm group was similar to the thymic change in the animals of the 100 and 300 ppm groups. Therefore, lymphoid depletion in animals of the 10, 100 and 300 ppm groups was considered related to treatment with DOTC.

In the screening reproductive toxicity study performed with DOTC, no effects were observed on the mating index, the pre-coital time was comparable for the control and the treated groups, the female fecundity index, female fertility index and male fertility index were not affected while the gestation index was 86, 100, 71 and 50 % in the control, 10, 100 and 300 ppm groups, respectively. The livebirth index was 99, 95, 53 and 60 % in the control, 10, 100 and 300 ppm groups, respectively. Post-implantation loss was 22.3, 21.0, 49.2 and 70 % for the control, 10, 100 and 300 ppm groups, respectively.

It was concluded by RAC in its 2012 assessment that the data do not allow a comprehensive evaluation for the endpoint fertility.

10.10.3 Comparison with the CLP criteria

The discussion and comparison with the CLP Criteria for adverse effects on fertility is done in Section 10.10.6 together with the discussion and comparison of CLP criteria for developmental effects and effects via lactation

10.10.4 Adverse effects on development

Table 11: Summary table of animal studies on adverse effects on development

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
<p>OECD TG 414 (Prenatal Developmental Toxicity Study)</p> <p>GLP</p> <p>key study</p> <p>1 (reliable without restriction)</p> <p>rabbit (New Zealand White)</p> <p>24 presumed-pregnant females per dose</p>	<p>Test material: DOTE</p> <p>In original Study (EC name): 2-ethylhexyl 10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate (96.1 % purity)</p> <p>0 (vehicle control), 4, 20, 80 mg/kg bw/day (nominal conc.)</p> <p>oral: gavage</p> <p>Vehicle: peanut oil</p> <p>Exposure: Daily from gestation day 6 to gestation day 28 (inclusive). (Daily throughout treatment period.)</p>	<p>NOAEL (maternal toxicity): 20 mg/kg bw/day (actual dose received) based on test material (biologically relevant depression [$> 10\%$] in thymus weight at 80 mg/kg bw/day)</p> <p>LO(A)EL (developmental toxicity): 80 mg/kg bw/day (actual dose received) based on test material (Biologically relevant effect on foetal weight (-11.9 %) and foetal crown-rump length (-10.7 %) relative to controls, statistically significant negative trend on foetal weight and foetal crown-rump length/1 high-dose litter with foetal loss [within historical controls] excluded)</p> <p>LO(A)EL (maternal toxicity): 80 mg/kg bw/day (actual dose received) based on test material (Biologically relevant depression (-12.8 %) in thymus weight, dose-dependent: -9.6 % at mid and -5.1 % at low dose group)</p>	<p>Anonymous (2014a)</p>
<p>Method: OECD Guideline 414 (Prenatal Developmental Toxicity Study)</p> <p>GLP</p> <p>key study</p> <p>1 (reliable without restriction)</p> <p>mouse (Swiss)</p> <p>25 pregnant females per dose</p>	<p>Test material: DOTE</p> <p>In original Study (EC name): 2-ethylhexyl 10-ethyl-4,4-dioctyl-7-oxo-8-oxa-3,5-dithia-4-stannatetradecanoate (96.1 % purity)</p> <p>0 (vehicle control), 15, 30, 60 mg/kg bw/day (nominal conc.)</p> <p>oral: gavage</p> <p>Vehicle: peanut oil</p> <p>Exposure: Females received test material daily from gestation day 5 to gestation day 17 (inclusive). (Daily throughout treatment period.)</p>	<p>NOAEL (maternal toxicity): 15 mg/kg bw/day (actual dose received) based on test material (Statistically significant decrease in thymus size at 30 mg/kg bw/day (2 animals) and 60 mg/kg bw/day (8 animals), statistically significant decrease in thymus weight from 30 mg/kg bw/day)</p> <p>Developmental toxicity: statistically significant positive trend on percentage of post-implantation loss: 0.9 ± 2.8 at low, 1.5 ± 4.9 at mid, and 2.6 ± 5.6 at high dose, respectively)</p> <p>LOAEL (maternal toxicity): 30 mg/kg bw/day (actual dose received) based on test material (Statistically significant depression in thymus size, stat. sign. depression (23 %) in thymus weight, 35 % at 60 mg/kg bw/day, treatment-related statistically not significant reduction in corrected maternal body weight gain -26.5% (2.34 ± 2.41 g) in the high dose mice relative to controls, mid and low dose -16.7% and -17.7%, respectively)</p>	<p>Anonymous (2014b)</p>
<p>similar to OECD TG 414 (Prenatal</p>	<p>Test material: DOTI/MOTI 80:20.</p>	<p>NOAEL (maternal toxicity): 5 mg/kg bw/day (slight but non-significant decrease in corrected body weight and corrected body weight gain of</p>	<p>Battenfeld, R. (1991)</p>

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
<p>Developmental Toxicity Study) supporting study GLP 2 (reliable with restrictions) rat (Han-Wistar SPF) 25 per treatment group</p>	<p>In original study named: Dioclytin bis(IOMA) [CAS No. 26401-97-8] : Octyltin tris(IOMA) [CAS No. 26401-86-5] (80:20 %) (See endpoint summary for justification of read-across) 1, 5, and 25 mg/kg/day (actual ingested) oral: gavage Vehicle: peanut oil Exposure: days 6-15 of gestation (once/day x 10 days)</p>	<p>the dams indicating a marginal maternal toxic effect of the test material) NOAEL (developmental toxicity): 5 mg test material/kg bw/day (significant increase in the percentage of dead foetuses)</p>	
<p>similar to OECD Guideline 414 (Prenatal Developmental Toxicity Study) GLP 3 (not reliable) Rabbit (New Zealand White) 23-24 females/group</p>	<p>Test material: DOTI/MOTI 80:20. In original study named: Dioclytin bis(IOMA) [CAS No. 26401-97-8] : Octyltin tris(IOMA) [CAS No. 26401-86-5] (80:20 %) (See endpoint summary for justification of read-across) 1.0, 10, and 100 mg/kg bw/day (actual ingested) oral: gavage Vehicle: peanut oil Exposure: days 6-18 of gestation (once/day x 13 days)</p>	<p>NOAEL (maternal toxicity): 10 mg/kg bw/day (based on increased incidence of abortions at 100 mg/kg bw/day) NOAEL (developmental toxicity): 10 mg test material/kg bw/day: Slight non-significant increase in minor skeletal head anomalies (incompletely ossified bones in the skull). 100 mg test material/kg bw/day: Significantly increased incidence of abortions, post implantation loss, minor visceral anomalies (severely dilated renal pelves and additional small vessels originating from the aortic arch), minor skeletal head anomalies (incompletely ossified bones in the skull), and skeletal variations of the sternum and feet bones (not or incompletely ossified sternbrae and feet bones); and a significant reduction in foetal body weight.)</p>	<p>Battenfeld, R. (1992)</p>
<p>Method equivalent or similar to OECD Guideline 414 (Prenatal Developmental Toxicity Study) 3 (not reliable) Mouse (NMRI) 22 to 25 females/group</p>	<p>Test material: DOTI/MOTI 80:20. In original study named: Dioclytin bis(IOMA) [CAS No. 26401-97-8] : Octyltin tris(IOMA) [CAS No. 26401-86-5] (80:20 %) (See endpoint summary for justification of read-across) 20, 30, or 45 mg/kg bw/day (group 1); 67 or 100 mg/kg bw/day (group 2) (actual ingested) oral: gavage Vehicle: peanut oil Exposure: days 6-17 of gestation</p>	<p>NOAEL (maternal toxicity): 30 mg test material/kg bw/day (Based on a significant decrease in thymus weight [-15 %] at 45 mg/kg bw/day.) NOAEL (developmental toxicity): 45 mg test material/kg bw/day (based on an increased incidence of cleft palate [+5.5 %] in foetuses from dams exposed to 67 mg test material/kg bw/day.)</p>	<p>Faqi, A.S., H. Schweinfurth, and I. Chahoud (2001)</p>

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
	(once/day x 12 days)		

Table 12: Summary table of human data on adverse effects on development

Type of data/report	Test substance,	Relevant information about the study (as applicable)	Observations	Reference
		No human data available		

Table 13: Summary table of other studies relevant for developmental toxicity

Type of study/data	Test substance, dose levels, duration of exposure	Relevant information about the study (as applicable)	Observations	Reference
<p>OECD Guideline 421- reproduction/ developmental screening study: sub-chronic (13 week) oral toxicity study in rats (OECD Test guideline 408), including a satellite group for a reproduction/ developmental screening study (OECD Test guideline 421)</p> <p>GLP</p> <p>Rat (Wistar) male, female</p> <p>44 females (four dose groups of 10 rats/sex)</p> <p>Oral: feed</p>	<p>Test material: Dichlorodioctylstannane (CAS no 3542-36-7) (purity 94 %)</p> <p>10, 100, 300 ppm (nominal in diet) The test substance intake of the female animals of the 10, 100 and 300 ppm groups were, respectively: Premating period days 0-7: 0.6, 5.8 and 13.5 mg/kg bw/day days 7-14: 0.7, 5.9 and 16.4 mg/kg bw/day; Gestation period GD 0-7: 0.7, 6.2 and 16.6 mg/kg bw/d, GD 7-14: 0.7, 6.2 and 17.0 mg/kg bw/day, GD 14-21: 0.5, 4.2 and 11.0 mg/kg bw/day; Lactation period: PND 1-4: 0.7, 5.0 and 8.4 mg/kg bw/day.</p> <p>Duration of exposure: females: daily for 2 consecutive weeks during the pre-mating period, daily during gestation (up to 26 days after study initiation) and up to euthanasia at or shortly after postnatal day (PND) 4. (daily); males: daily for 13 weeks prior to mating</p>	<p>Read-across from supporting substance (structural analogue or surrogate)</p> <p>(See endpoint summary for justification of read-across)</p> <p>Reliability: 2 (reliable with restrictions)</p>	<p>NOAEL (reproduction toxicity): 10 ppm (0.5 — 0.7 mg/kg bw/day (female)) (Based on reproductive and developmental effects: animals showing only implantations at necropsy, animals delivering only dead pups, decreases in gestation, live birth and viability indices and increases in post-implantation loss and number of runts)</p> <p>LOAEL (general toxicity): 10 ppm (0.5 — 0.7 mg/kg bw/day (female)) (decreases in absolute and relative thymus weights associated with treatment related lymphoid depletion at 10, 100 and 300 ppm groups)</p>	<p>Appel, M.J. and D.H. Waalkens-Berendsen. (2004)</p>

10.10.5 Short summary and overall relevance of the provided information on adverse effects on development

Two new key studies, investigating the developmental toxicity of DOTE itself, are available; one study was conducted in mice and the other in rabbits. Both studies were performed under GLP conditions and in accordance with the standardised guideline OECD 414.

1/ In the new mice developmental toxicity study (Anonymous, 2014b), dams were given DOTE 96.1 % purity at 0, 15, 30, and 60 mg test material/kg bw/day (corresponding to 14.4, 28.8, and 57.7 mg DOTE/kg bw/day) during day 5 to 17 of pregnancy.

Maternal toxic effects

Reduction in thymus weight from 30 mg/kg bw/day

Embryotoxic / teratogenic effects

None

MATERNAL DATA

- Clinical Signs of Toxicity and Morbidity/Mortality

The animals did not reveal any clinical signs of toxicity and mortality at any of the tested doses throughout the experiment.

- Maternal Body Weight

There were no statistically significant differences in maternal body weights across the dose groups on any single gestation day. However, there was a clear dose-related pattern of reduced body weights beginning after GD6, the first day of dosing, and continuing for the duration of the study. The high dose was the most severely affected, though a dose-related decrement relative to the control body weights can be seen across all doses particularly from GD16 to GD18. At 30 mg/kg bw/day the maternal weight effect was marginal, but maternal body weight gain in the 60 mg/kg bw/day high dose group was 11.5 % [uncorrected] and 26.6 % [corrected] less (2.34 ± 2.41 g) than the vehicle control (3.19 ± 2.23 g). The corrected body weight gain in the low (15 mg/kg bw/day) and mid (30 mg/kg bw/day) dose groups were -16.7 % (2.62 ± 2.03 g) and -17.7 % (2.66 ± 2.46 g) when compared to the controls.

- Feed Consumption

No treatment related differences in average feed consumption were observed at any dose.

- Gross Pathology [Maternal]

There was a treatment-related macroscopic finding of reduced maternal thymus weight.

The mean maternal thymus weight was statistically significantly reduced (-23 %) in the 30 mg/kg bw/day [mid] and 35 % in the 60 mg/kg bw/day [high] dose groups. The mean maternal thymus weight in the low dose mice was reduced relative to controls, but was not statistically significant. These observations are indicative of a treatment-related specific target organ toxicity resulting from exposure to the test material. No other gross pathological findings were noted in any dose group.

PREGNANCY DATA

A total number of 21 (84 %), 21 (84 %), 20 (80 %) and 20 (80 %) mated females were confirmed pregnant at the time of caesarean section for groups G1, G2, G3 and G4, respectively.

UTERINE OBSERVATIONS

There were no statistically significant differences in these gravid uterus weights, number of implantation sites, pre- and post-implantation loss or early or late resorptions across dose groups when compared to the vehicle control.

REPRODUCTION DATA

No treatment related effects were noted in mean gravid uterus weight, no. of corpora lutea, no. of implantations in all the groups, and no. of early or late resorptions. There was a statistically significant

positive trend on percentage of post implantation loss of 0.9 ± 2.8 at the low, 1.5 ± 4.9 at the mid and 2.6 ± 5.6 at the high dose group, respectively.

There were no statistically significant differences in the sex ratio, mean litter size or the number of live foetuses per dam across dose groups when compared to the vehicle control.

FOETAL DATA

- Foetal Weight

The mean foetal weights [combined sexes] were 1.35, 1.37, 1.30 and 1.31 grams for groups G1, G2, G3 and G4, respectively. Mean foetal weights were not statistically significantly different across the dose groups when compared to controls.

- External Examination

No external abnormalities were noted during gross examinations of foetuses at any dose.

- Visceral Examination

No treatment-related abnormalities were observed during visceral examinations of foetuses at any dose. The noted findings [pale coloured kidneys and dilated renal pelvis] are common findings for foetuses of this species and strain. The observations were not dose dependent, nor was the severity of the anomaly increased with dose. This result supports the conclusion that the findings are incidental and that the test material did not produce an adverse effect during foetal development of the soft tissues.

- Skeletal Examination

There was a single incidence in one litter in the high dose group of fused sternum and a single incidence in a different high dose litter of short rib. Single incidences, even of rare malformations (which these are not), cannot be reliably attributed to treatment. The other noted anomalies (poorly ossified frontal, parietal and inter-parietal bones; ossification site at first lumbar vertebrae; supplementary ribs) are common findings for foetuses of this species and strain. These morphologic observations did not occur in a dose-dependent pattern, nor was the severity of the anomaly increased with dose. The findings were therefore considered to be incidental, and not indicative of a teratogenic effect.

- Crown-rump length

The average crown-rump lengths were 23.2, 24.0, 23.3 and 22.9 mm for groups, G1 through G4, respectively. There were no statistically significant differences in length across all dose groups when compared to the control.

2/ In the new rabbit developmental toxicity study (Anonymous, 2014a), dams were given DOTE 96.1 % purity during day 6-28 of pregnancy at 0, 4, 20 and 80 mg test material/kg bw/day (corresponding to 3.8, 14.4, and 76.9 mg DOTE/kg bw/day).

Maternal toxic effects

Biologically relevant depression [$> 10\%$] in thymus weight at 80 mg/kg bw/day

Embryotoxic / teratogenic effects

None

PREGNANCY DATA

A total number of 19, 21, 19 and 20 mated females were confirmed pregnant in groups G1, G2, G3 and G4, respectively (0, 4, 20 and 80 mg/kg bw/day, respectively). This is a pregnancy rate of 79, 88, 79 and 83 % at the time of caesarean section for groups G1, G2, G3 and G4, respectively.

MATERNAL DATA

- General Tolerability: No deaths or abortions were observed during the experimental period and there were no clinical signs recorded which were indicative of overt toxicity.

- Maternal Body weight, Body Weight Gain, and Corrected Body Weight Gain: There were no statistically significant differences in maternal body weights across the dose groups on any single gestation day, nor was there a dose-related pattern of reduced body weights beginning after GD6, the first day of dosing. There were

no statistically significant differences in uncorrected maternal body weight gain across the dose groups for any single gestational period. Treated does had higher weights than controls on GD29; this was considered to be a random occurrence and not related to test material administration.

- Food Consumption: No treatment related differences in average feed consumption were observed at any dose.

- Gross Pathology: There was a treatment-related macroscopic finding of reduced maternal thymus weight. The mean maternal thymus weights were -5.1, -9.6, and -12.8 % in the 4 mg/kg bw/day [low], 20 mg/kg bw/day [mid], and 80 mg/kg bw/day [high] dose groups, respectively when compared to controls. While these decreases were not statistically significant, the results were dose-dependent relative to controls and are consistent with data from other species [rat and mouse] which demonstrate the thymus is a target organ for octyltin compounds. These observations are indicative of a treatment-related specific target organ toxicity resulting from exposure to the test material. No other gross pathological findings were noted in any dose group.

REPRODUCTION DATA

No treatment related effects were noted in the number of corpora lutea, number of implantation sites, number of early or late resorptions, or percentage of post implantation loss across all the groups.

The mean gravid uterus weights were 230.1, 309.5, 253.7 and 196.9 g for groups G1 through G4, respectively. A statistically significant increase in the mean gravid uterus weight was noted at the low dose when compared to the vehicle control. This difference was attributed to the increase in the number of foetuses compared to the vehicle control group and other dose groups.

The mean pre-implantation losses were 0.9, 0.8, 2.3, and 4.9 % for groups G1, G2, G3 and G4, respectively. Mean post-implantation losses for these treatment groups were 3.1, 3.5, 6.4 and 5.7, respectively. These losses are within the historical control range and there is no clear evidence of a dose-response across dose groups.

There were no statistically significant differences in either the mean litter size or the number of live foetuses per dose across dose groups when compared to the vehicle control.

No treatment-related effect on the sex ratio of foetuses was noted at any dose compared to the vehicle controls.

FOETAL DATA

- Foetal Weight

The mean foetal weights [combined sexes] were 36.6, 37.3, 35.5, and 32.3 g for groups G1, G2, G3 and G4, respectively. Mean foetal weights were not statistically significantly different across the dose groups when compared to controls. At the high dose the mean foetal body weight was -11.9 % relative to controls which suggests a biologically-relevant, but marginal effect on foetal maturation. However, the mean weight for this group was disproportionately affected by low foetal body weights in a single litter in this group. There was a statistically significant negative trend on foetal weight (mean foetal weight 33.5 g for group G4 with single litter excluded).

- External Examination

No external abnormalities were noted during gross examinations of foetuses at any dose.

- Visceral Examination

No treatment-related abnormalities were observed during visceral examinations of foetuses at any dose. The noted findings [pale coloured kidneys and dilated renal pelvis] are common findings for foetuses of this species and strain. The observations were not dose dependent, nor was the severity of the anomaly increased with dose. This result supports the conclusion that the findings are incidental and that the test material did not produce an adverse effect during foetal development of the soft tissues.

- Crown rump length:

The mean foetal crown-rump length for both sexes was 92.1, 91.1, 89.3 and 82.3 mm for groups G1, G2, G3 and G4, respectively. A statistically significant reduction in the mean foetal crown-rump length was noted in the high dose group when compared to controls. This was -10.7 % relative to the vehicle control suggesting a

marginal but biologically relevant effect on foetal maturation which correlated to the degree of skeletal ossification. One litter in this group had low crown-rump lengths which were found to disproportionately affect the mean. There was a statistically significant negative trend when not considering all the litters of this group (mean foetal crown-rump length 83.8 mm for group G4 with single litter excluded). There were no correlating statistically significant differences for any other parameter.

- Skeletal Examination:

The noted skeletal anomalies in the study are common findings for foetuses for this species and strain. The morphologic observations did not occur in a dose-dependent pattern, nor was the severity linked to an increase in dose. The incidences of absent sternum No. 5, a measure of delayed ossification were found to be higher in controls compared to any other treatment group. Other indicators of delayed ossification such as the absence of proximal phalanges were found to be single incidences within a litter both in the control group and treatment groups. Poor ossification in sternum No. 5 and No. 6, an indicator of delayed ossification, was the only variation to be found to occur as multiple incidences within a litter. This variation however is considered to be less significant than the absence of sternum No. 5. The conflicting results indicate that these findings are incidental.

Supporting information is available in the form of an embryotoxicity and teratogenicity study which was conducted with a read across material, consisting of DOTI [CAS No. 26401-97-8] : MOTI [CAS No. 26401-86-5] (80:20 %). Read across is considered justified as DOTE and DOTI are isomers of the same compound and are structural analogues of each other. Based on the recently conducted developmental toxicity studies in two species with DOTE it is considered that the use of data on DOTI would be considered in a weight-of-evidence assessment of the registered substance.

3/ In the developmental toxicity study in rats (Battenfeld, 1991), dams were treated with DOTI/MOTI (80:20 %) at 1, 5 and 25 mg/kg bw/day during day 6-15 of gestation.

Maternal effects:

Alopecia was observed in single animals of all four groups and was not attributed to treatment. There was a slight (non-significant) decrease in corrected body weight and corrected body weight gain from day 6 to day 21 at 25 mg/kg bw/day dose. This reduction was attributed largely to one single dam (dam No.97).

Foetal observations:

No embryofetal effects were reliably attributed to treatment as the observed embryoletality was marginal (observed in only one dam) and is therefore considered to have occurred by chance

It should be noted that for rats exposed to DOTI/MOTI [GD6-15] the foetuses did not show the malformations or variations of bone formation as observed in mice and rabbits. For this test material the rat was the least sensitive species.

At 1 and 5 mg/kg/day dose-levels no treatment-related effects were observed. At the 25 mg/kg/day dose level there was a slight, but not statistically significant decrease in corrected body weight and corrected body weight gain of the dams. This reduction was to a great extent due to the loss of corrected body weight in one single dam (-58g in dam No. 97) Seven dead foetuses were observed, all in one litter in group 4. This led to an artefact in the statistics of the percentage of dead foetuses. All dead foetuses in the group were from a single dam (No. 97), and in addition to the high maternal weight loss, four early resorptions and an extremely low mean weight (2.2 g) of the two living foetuses were found in this animal. There were no treatment-related effects (external, visceral or skeletal malformations) at any dose-level. The frequency of all external, visceral and skeletal variations was within historical control limits and there was no dose-response.

For DOTI/MOTI, no adverse effects were observed at the low [1 mg/kg/day] or intermediate [5 mg/kg/day] doses. The high dose [25 mg/kg/day] produced marginal maternal toxicity and no embryofetal effects were reliably attributed to treatment. The effects noted in a single high dose dam were considered to have occurred by chance.

RAC position (2011):

“Indeed there is only weak evidence on developmental effects from this rat study since dead foetuses were

seen only from one dam”

4/ Two generation study performed under GLP and according to OECD 416 with Diocetyl tin bis(IOMA)/Octyltin tris(IOMA) (Anonymous, 1997)

Please refer to section 'effects on fertility' for details.

The NOAEL for P males and females was 20 ppm diet (approx. 1.5 mg test material/kg bw/day) based on a slightly reduced relative thymus weight for males at 60 ppm (approx. 4.4 mg test material/kg bw/day).

The NOAEL for the F1 generation was 20 ppm (approx. 1.6 mg test material /kg bw/day), based on a reduction in relative thymus weights for males and females at 60 ppm (approx. 4.7 mg test material/kg bw/day).

No teratogenic effects were observed in this study.

Three additional studies had been considered by RAC in its scientific opinion reached on 8 June 2012. They have been regarded for hazard classification in a weight-of-evidence assessment and the significance weighted accordingly.

5/ In the mice developmental toxicity study (Faqi, 2001), dams were given DOTI/MOTI (80:20 %) at 20, 30, 45, 67 and 100 mg test material/kg bw/day (corresponding to 16, 24, 36, 53.6, and 80 mg DOTI/kg bw/day) during day 6 to 17 of pregnancy.

Maternal effects:

There was a dose dependent decrease (-43 % at the high dose and -10 % at 67 mg test material/kg bw/day, respectively) in maternal body weight gain (mean (control): 5.1 ± 7.0 g, mean ± SD: 2.9 ± 4.8 g at 100 mg test material/kg bw/day), but differences were not significant in mice exposed to the test substance. No signs of toxicity were observed with the exception of one dam in the 100 mg/kg bw/day dose group that died. Pregnancy rates were comparable between treated groups and the control groups.

The mean maternal thymus weights in the 45 and 100 mg test material/kg bw/day dose groups were significantly lower than the control groups (-27 %, $p < 0.05$ at 100 mg test material/kg bw/day). At 67 mg/kg bw/day, the mean maternal thymus weight was slightly (-13 %) but not significantly decreased. Maternal liver weights were significantly lower in the 100 mg/kg bw/day dose group (-23 %, $p < 0.05$ at 100 mg test material/kg bw/day). The number of implantations per litter was comparable between treated groups and the control groups. Resorption rates were significantly increased in mice treated with 67 (13 %) or 100 mg test material/kg bw/day (16 %).

Foetal observations:

Foetal weights were significantly decreased in the 67 and 100 mg test material/kg bw/day groups (-9 and -18 %, respectively). There were no dead foetuses in any of the treated groups. There were no external malformations reported in the foetuses exposed to 20, 30, or 45 mg test material/kg bw/day, however a significantly increased incidence of cleft palate in the foetuses exposed to 67 or 100 mg test material/kg bw/day were observed (5.5 and 9.3 %, respectively), and incidences of bent forelimbs (18.5 %) and exencephaly (7.1 %) were significant in the foetuses exposed to 100 mg test material/kg bw/day. Skeletal variations reported in the low dose groups included unossified digit (4.4 % at 45 mg test material/kg bw/day) and supernumerary cervical ribs (significantly increased at 20 and 45 mg test material/kg bw/day, but not at 30 mg test material/kg bw/day); hind paw incompletely ossified (9.3 %), Os frontale misshapen (25.2 %), and interparietale incompletely ossified (24.4 %) (Significantly increased at 45 mg test material/kg bw/day); and supernumerary lumbar or cervical ribs (significantly increased at 20, 30, and/or 45 mg test material/kg bw/day). There was a significant increase in skeletal abnormalities in the foetuses of dams exposed to 67 or 100 mg test material/kg bw/day. Skeletal abnormalities reported in these dose groups included bent forelimbs, bent hindlimbs, dislocated sternum, fused or bent ribs (6.4 and 35.5 % at the 67 or 100 mg test material/kg bw/day groups, respectively), or bent vertebral column. Skeletal variations were observed in the low dose groups (20, 30, or 45 mg test material/kg bw/day).

The authors defined malformations as a permanent or irreversible structural change that is likely to adversely affect survival or health. The authors reported a no-observed-adverse-effect-level (NOAEL) for each endpoint examined, i. e., malformations, variations, organ toxicity.

- The embryo-foetal NOAEL for malformations was reported as 45 mg test material/kg bw/day, based on an increased incidence of cleft palate (5.5 %) in foetuses from dams exposed to 67 mg test material/kg bw/day.
- A NOAEL for skeletal variations could not be determined, but would be expected to be < 20 mg/kg bw/day, based on an increased incidence of supernumerary lumbar ribs (68.5 %) observed at 20 mg test material/kg bw/day.
- The authors reported that the NOAEL for maternal organ toxicity was 30 mg test material/kg bw/day, based on a significant decrease in thymus weight (15 %) at 45 mg test material/kg bw/day.

The mouse study reported by **Faqi (2001)** with DOTI/MOTI (80:20 %) failed to fulfil the requirements of a GLP OECD 414 guideline study since detailed information on the following were not included: maternal body weight, food consumption, housing, dosing, analysis of the dosing formulations or individual animals data. Furthermore, no information was given on foetal sex ratio, foetal weight per sex or internal malformations and no comparison to historical control data was carried out. The study was therefore considered to be deficient and the results are taken forward for hazard classification with lower significance than the new key study, investigating the developmental toxicity with the substance itself on mice.

6/ In the rabbit developmental toxicity study (Battenfeld, 1992), dams were given DOTI/MOTI (80:20 %) during day 6-18 of pregnancy at 1, 10 and 100 mg test material/kg bw/day (corresponding to 1, 8, and 80 mg DOTI/kg bw/day).

Maternal effects:

Except for a nasal haemorrhage in one dam of Group 2 (1 mg/kg bw/day), slight torticollis in one dam of Group 3 (10 mg/kg bw/day), and bloody outflow in 3 dams of Group 4 (100 mg/kg bw/day), no clinical observations were made. In total, 18 of 24 dams in Group 1, 23/23 in Group 2, 18/22 in Group 3, and 17/24 in Group 4 survived until day 28. Two dams in Group 1 and 3 dams in Group 3 died after treatment had commenced. Death resulted from infectious diseases (pneumonia or enteritis), and there was no dose-related increase. Therefore, these deaths were not attributed to the test substance. In Group 1, 3 dams were eliminated because of normal deliveries before day 28. Before start of treatment, one dam in Group 1 and one dam in Group 2 were found dead. Maternal body weight data did not reveal differences between treatment groups. Abortion was diagnosed in one dam of Group 1 and 4 dams of Group 4. All abortions occurred after termination of treatment. The high incidence of abortion in Group 4 was considered to result "at least partly from a slight maternal toxic effect of the test compound."

Foetal observation:

Total foetal death was found only in Groups 1 and 4. In both groups, total post-implantational loss occurred in 3 dams. Percentages of post-implantation losses per group were 17.7 % (control), 10.5 % (1 mg/kg bw/day), 5.7 % (10 mg/kg bw/day), and 28.4 %, $p < 0.05$ (100 mg/kg bw/day). The significant increase in post-implantation loss at the high dose-levels was explained by a significant increase of total resorptions (28.4 %, $p < 0.05$ vs. 17.1 % in controls).

External examination revealed two nasal clefts and an encephalocele in one foetus of Group 2. Umbilical hernia was found in one foetus of the control group and in one foetus each in Groups 3 and 4. These were not associated with treatment. Other findings, such as malformations of the vertebral column (one animal in Group 4) and absence of the right kidney and adrenal gland (one animal in Group 4) were regarded as chance findings and not attributed to treatment due to their single occurrence and because they represented totally different types of malformations. The lack of a statistically significant difference to the control group and inconsistency regarding the type of anomaly found did not "point towards a compound-related effect." Foetuses with minor external anomalies (flexion of digits and limbs, open eyelids, shortened tail) were observed in all four groups, and not attributed to the test substance. Minor visceral anomalies found included severely dilated renal pelves and additional small vessels originating from the aortic arch. The statistically

significant increase in the incidence of visceral anomalies of fetuses in Group 4 is an indication of retardation in foetal development. Individual body weights of the fetuses in Group 4 with minor visceral anomalies were approximately 40 % lower than the mean weight of control fetuses.

Suspected or definite compound-related changes noted included:

-1 mg/kg bw/day: No substance-related effects.

-10 mg/kg bw/day: Slight non-significant increase in minor skeletal head anomalies (incompletely ossified bones in the skull).

-100 mg/kg bw/day: clear substance-related embryotoxic effects were noted i. e. significantly increased incidence of abortions, post-implantation loss, minor visceral anomalies (severely dilated renal pelves and additional small vessels originating from the aortic arch), minor skeletal head anomalies (incompletely ossified bones in the skull), and skeletal variations of the sternum and feet bones (not or incompletely ossified sternebrae and feet bones); and a significant reduction in foetal body weight.

In conclusion, the author of the rabbit developmental toxicity study reported that the evaluation of reproduction data and foetal weights indicated a slight embryolethal and moderate retarded effect (with regard to foetal development) at the high dose level (100 mg/kg bw/day) associated with maternal toxicity (abortions).

In the rabbit study reported by **Battenfeld (1992)** with DOTI/MOTI (80:20 %) it was not possible to interpret the results of the study based on the information presented in the study report. During the study maternal disease was reported; the disease that is described in the report is common in rabbits of the age and strain used in the study. It was not possible to ascertain whether maternal effects which were observed resulted from the disease or from treatment with the test material since detailed examination of the immune system are not included in developmental toxicity studies and were not included as part of this study. There was one case of torticollis, a symptom of encephalitozoonosis, a parasitic disease occurring in immune deficient animals, which was reported. Since no effects were seen in the rangefinder study, at doses up to 30 mg test material/kg bw/day, the maternal findings observed in the main study were considered spurious. The exposure time was rather short (13 days) as also noted by RAC in its 2012 assessment. Overall the study was considered to be deficient and the results are taken forward for hazard classification with lower significance than the new key study, investigating the developmental toxicity with the DOTE substance itself on rabbits.

7/ In the Reproduction / Developmental Toxicity Screening Test (according to OECD 421) (Appel and Waalkens, 2004) rats were given DOTC > 94 % purity at 10, 100, 300 mg/kg in the diet:

At 10 ppm (equivalent 0.7 mg/kg bw/day for males and 0.5-0.7 mg/kg bw/day for females), treatment-related effects to dams included lymphoid depletion were observed in dams.

At 100 ppm (equivalent to 6.5 mg/kg bw/day for males -6.8 mg/kg bw/day for females, treatment-related effects included increased post-implantation loss (49 %), decreased gestation index (71 %) decreased live birth index (53 %), decreased viability index (74 %), increased number of runts, increased pup mortality (PND 1 and 4), and decreased absolute and relative thymus weights and lymphoid depletion in the dams.

At 300 ppm (equivalent to 19.3 mg/kg bw/day for males -19.8 mg/kg bw/day for females), treatment-related effects included increased in post-implantation loss (70 %), decreased gestation index (50 %), decreased live birth index (60 %) decreased viability index (12 %), increased number of runts, decreased pups weights (PND 1 and 4), increased pup mortality (PND 1 and 4), and decreased absolute and relative thymus weights and lymphoid depletion (dams).

Summary of litter data

- **Litter size:** The mean number of pups delivered per litter amounted to 11.7, 11.0, 10.3 and 8.6 for the control, 10, 100 and 300 ppm groups, respectively.

- **Litter weight:** Mean pup weights and pup weight changes were similar in the 10 and 100 ppm groups when compared to the control group. Pup weight of the 300 ppm group (PND 1, 3 litters and PND 4, 1 litter) was reduced.

- **Pup mortality:** 1.4, 4.5, 47 and 40 % in the control, 10, 100 and 300 ppm groups, respectively (PND 1); 5.8, 8.3, 26 and 88 % in the control, 10, 100 and 300 ppm, respectively (PND 4).
- **Number viable:** The viability index (PND 1-4) was 94, 92, 74 and 12 % in the control, 10, 100 and 300 ppm groups, respectively.
- **Number live pups per litter:** 11.5, 10.5, 7.6 and 6.5 for the control, 10, 100 and 300 ppm groups, respectively (PND 1); 10.8, 11.0, 9.3 and 3.0 for the control, 10, 100 and 300 ppm groups, respectively (PND 4).
- **Sex ratio:** No difference was observed in the sex ratio between the groups.

The above developmental effects were associated with maternal toxicity substantiated by a statistically significant decrease in absolute and relative thymus weight in the 100 (c. 62 and 67 % in male and females,) and 300 ppm group (31 and 38 % in males and females) and a moderate to very severe lymphoid depletion in dams (5/10 animals at 10 ppm and in all animals of the 100 and 300 ppm groups).

Based on reproductive and developmental effects in the screening reproductive toxicity assay (particularly severe post-implantation losses and foetal losses) observed after mating of 100 and 300 ppm female of the satellite groups with male animals of the main study, the low dose level of 10 ppm in diet (equivalent to 0.7 mg/ kg bw/day in males and 0.5-0.7 mg/kg bw/day for females) can be considered as a NOAEL for fertility and developmental effects.

Based on the treatment related histological changes in the thymus (lymphoid depletion) of the 10 mg/kg female animals of the satellite groups, 10 ppm in diet (equivalent to 0.5-0.7 mg/kg bw/day) was considered to be a LOAEL for maternal toxicity.

To assess teratogenic effects was not subject of this study. Thus, the animals were not in depth examined regarding external, soft tissue or skeletal abnormalities. However, grossly visible abnormalities were recorded.

This study reported by **Appel, M.J. and D.H. Waalkens-Berendsen. (2004)** was regarded in a weight-of-evidence assessment and the significance weighted accordingly based on the results of the recent hydrolysis study (Anonymous, 2015, see section 9.1), which indicate the substance DOTE is the only hydrolysis product of DOTE which was formed at pH 1.2 and which, however, does not exclude that no dioctyltin dichloride can be formed in vivo.

A direct comparison of the studies using DOTI/MOTI (80:20 %) with the newer studies on the substance itself is given at the end of section 10.10.10 within the conclusion on classification and labelling of DOTE.

10.10.6 Comparison with the CLP criteria

The classification criteria for reproductive toxicity for the ECHA are documented in Section 3.7.2 of the Guidance on the Application of the CLP Criteria, Guidance to Regulation (EC) No. 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures, Version 4.1, June 2015.

For the purpose of classification the hazard class Reproductive Toxicity is differentiated into:

- adverse effects
 - o on sexual function and fertility, or
 - o on development;
- effects on or via lactation.

Annex I: 3.7.1.4. defines adverse effects on development of the offspring as follows:

“Developmental toxicity includes, in its widest sense, any effect which interferes with normal development of the conceptus, either before or after birth, and resulting from exposure of either parent prior to conception, or exposure of the developing offspring during prenatal development, or postnatally, to the time of sexual maturation. However, it is considered that classification under the heading of developmental toxicity is primarily intended to provide a hazard warning for pregnant women, and for men and women of reproductive capacity. Therefore, **for pragmatic purposes of classification, developmental toxicity essentially means adverse effects induced during pregnancy, or as a result of parental exposure.** These

effects can be manifested at any point in the life span of the organism. The major manifestations of developmental toxicity include (1) death of the developing organism, (2) structural abnormality, (3) altered growth, and (4) functional deficiency.”

The hazard categories for the reproductive endpoint are presented in Table 3.7.1 (a) of the referenced document.

Rationale for classification

There are no human reproductive data on DOTE, therefore DOTE is not a candidate for Category 1A.

There is no animal evidence that DOTE interferes with sexual function or fertility, therefore DOTE should not be classified for these endpoints.

Based on recently conducted developmental toxicity studies according to OECD TG 414 (Anonymous, 2014a, b) there is animal evidence in the most sensitive species, rabbit and mouse, that DOTE interferes with gestational integrity in mice and the embryological development in rabbits. In mice, study results show a statistically significant positive trend on percentages of post implantation loss. In rabbits, at the high dose (80 mg/kg bw/day) the mean foetal body weight decreased about 12 % relative to controls, suggesting a marginal but biologically relevant effect on foetal maturation. Furthermore, a statistically significant reduction in the mean foetal crown-rump length was noted in this study (-10.7 % relative to the controls), again suggesting a marginal but biologically relevant effect on foetal maturation which moreover correlated to the degree of skeletal ossification. DOTE, however, produced no statistically significant adverse effects on foetal morphology of skeletal elements or soft tissues even in the presence of slight maternal toxicity. Observed effects were only statistically significant in trend tests, supporting the assumption that the effects were treatment- and dose-related. Furthermore, in comparison to other studies on analogues the highest dose tested using DOTE was notably lower. According to the OECD TG 414 the highest dose used in the experiment should be chosen with the aim to induce some developmental and/or maternal toxicity (clinical signs or a significant decrease in body weight) but not death or severe suffering. The descending sequence of dose levels should further be selected with a view to demonstrating any dosage-related response and NOAEL. Hence, the highest dose tested (Anonymous 2014a, b) might have been too low to be able to detect such a dose-response relationship and thus might only reflect the starting point of a potential dosage-related response.

There is evidence of postnatal toxicity from a two generation study with a related substance, DOTI/MOTI 80:20. DOTI and DOTE are isomers differing only slightly in the structure of the C-8 alcohol of the mercaptoester ligand (either *iso*-octanol or 2-ethylhexanol, respectively). Since these alcohols are so close in structure, their respective mercaptoacetate esters are expected to have very similar physicochemical and toxicological properties, including hydrolysis products. In the two generation study with DOTI/MOTI, postnatal deaths in the F1 and F2 generations occurred, however, the offspring did not die during the perinatal period. Deaths occurred after PND4 and hence might rather be attributed to postnatal exposure of lactating pups than to in utero exposure. In this case, these findings do not meet the criteria noted in bold text of the definition above.

There were relevant developmental effects in the two generation study and in the developmental toxicity studies performed with DOTI/MOTI, particularly the effects on pups such as the increased incidence of abortions, marked retardations of foetal development, increased number of runts, decreased foetal weight, decreased number of pups per litter, increased post-implantation loss, and decreased thymus weight for the F0 parent and F1 progeny.

In addition, the findings of the screening reproductive toxicity feeding study with DOTC were consistent with a part of these particular findings (increase in post-implantation loss, decreased viability index, increased number of runts, decreased pups weights). These effects were observed at dose levels inducing some maternal toxicity (statistically significant decrease in absolute and relative thymus weight and a moderate to very severe lymphoid depletion in dams). A recent *in vitro* hydrolysis study under simulated mammalian gastric conditions, admittedly only identified DOTE as a hydrolysis product and no DOTC could be detected under the particular study conditions, however this study does not allow to conclude, if and which other dioctyl species might be formed after systemic uptake of DOTE under *in vivo* conditions. Hence,

read-across from studies using DOTC as test material have to be included when discussing potential reproductive effects of DOTE and cannot be disregarded.

The above reported effects (increased post-implantation loss, increase incidence of resorption, increase pups mortality, depressed foetal weight) are indicative of developmental effects. These effects observed in all the above reported studies were associated with no or only slight maternal (thymo-)toxicity that has no bearing on the developmental toxicity. In the absence of significant/systemic maternal toxicity, DOTE should be classified for developmental toxicity.

In summary, the weight-of-evidence analysis for DOTE for toxicity studies in experimental animals there is some evidence which is not sufficiently convincing to place the substance in Category 1. The effects recorded with DOTI/MOTI are considered of toxicological significance. For DOTE per se there is some evidence of developmental toxicity occurring in two different species with low toxicological significance (trend only). Repr. 1B should only be chosen if there is clear evidence. Therefore, in accordance with the criteria for classification as defined in Annex I, Regulation (EC) No. 1272/2008 (CLP), classification of the substance with respect to developmental toxicity as Repr. 2 (H361d) is appropriate.

10.10.7 Adverse effects on or via lactation

Not evaluated in this dossier

10.10.8 Short summary and overall relevance of the provided information on effects on or via lactation

The postnatal deaths in the F1 and F2 generations of the 2G study with DOTI/MOTI occurred during the lactation period. The increase in pup mortality, respectively decrease in viability at birth and probably growth retardation during the lactational period has been discussed in Section 10.10.6 and considered as indicative of developmental effects. However, the low weight gain and deaths occurring after PND4 were explained as a consequence of repeated-dose exposure from the maternal milk and/or from the diet.

As a contribution of test substance in the milk cannot be estimated, and the (assumed) presence in the milk only is not sufficient to justify a classification for effects on/via lactation, no classification for effects on or via lactation is proposed.

10.10.9 Comparison with the CLP criteria

The discussion and comparison with the CLP Criteria for adverse effects via lactation is done in Section 10.10.6 together with the discussion and comparison of CLP criteria for developmental effects and effects on fertility.

10.10.10 Conclusion on classification and labelling for reproductive toxicity

A two generation study in rats with DOTI/MOTI (78.8:16.9, 80:20 ratio) showed maternal effects on the thymus and this result was the critical toxic effect for the adult NOAEL.

The body weight of F1 animals showed no statistically significant differences from controls in either sex at birth, PND 4 or PND 7, but showed a statistically significant decrease in both male and female pup body weights at PND 14 which persisted through PND 21. This correlates exactly with the incidences of postnatal losses documented for these F1 animals. Up to PND 4 the number of F1 pups was not different from controls for either sex. Pups were culled to 4/sex/litter on PND 4 (a routine practice). After culling, the pup losses across the four dose groups between PND4 and PND21 were 0, 4, 11, and 20 for the control, 20, 60, and 200 ppm doses. This is most probably not a reproductive effect; it can be best explained as a consequence of a direct toxic effect of dioctyltin species on these young animals who may well be receiving two sources of exposure, one from the maternal milk and a second from the diet as they begin to eat adult feed during the lactation period. It may (in theory) also be a consequence of impaired nursing behaviour as a result of maternal toxicity for which no clear indication was given. The mg/kg/day dose to these juveniles is likely

higher than the dose to the lactating F0 females (the mothers) and the dose-response of the juvenile deaths supports this conclusion. In addition, the delay in vaginal opening of the high dose females is likely due to the non-specific toxic effect and low weight gain resulting from this postnatal exposure.

For the reproductive testing of the F1 generation, DOTI/MOTI did not produce an adverse effect on mating, fertility, pregnancy rates, gestation, litter size or neonatal body weight. There was an increase in stillbirths which was inexplicably marked both non-statistically significant and statistically significant at the high dose; however 12 of the 26 still births for the group were in a single litter. This diminishes any statistical significance in the endpoint analysis. There was an effect on thymus weight of the F2 animals and the histopathology of the thymus was typical of the known target organ effect of multiple dioctyltin substances on this gland.

The body weight of F2 animals at the mid and low doses showed no statistically significant differences from controls in either sex at any time. The body weight of F2 animals at the high dose showed no statistically significant differences from controls in either sex at birth, however at PND4 (postnatal day 4 and thereafter through PND21 body weights of both sexes were statistically decreased.

This result for the F2 pups is parallel to the outcome from the F1 pups. The effect on weight correlated exactly with the incidences of postnatal losses documented for the F2 animals. Up to PND 4 the number of F2 pups was not different from controls for either sex. After culling, the pup losses across the four dose groups between PND 4 and PND 21 were 9, 26, 29, and 22 for the control, 20, 60, and 200 ppm doses. This is not a reproductive effect for the F2 generation either. It is the second manifestation of a direct toxic effect of dioctyltin species on young animals who may well be receiving two sources of exposure, one from the maternal milk and a second from the diet as they begin to eat adult feed during the lactation period. The dose-response of the juvenile deaths supports this conclusion. No assessment of vaginal opening or testicular descent was done in the F2 animals because the experiment was terminated at PND 21, prior to the occurrence of these developmental landmarks.

The reproductive NOEL is the high dose, 200 ppm. The maternal and paternal NOEL is 20 ppm (the low dose) which is driven by the effect of DOTI/MOTI test material on the thymus gland.

The outcomes for the PND4-PND21 pups of the F1 and F2 generations, however, should not be used in setting the reproductive NOEL, because the effect does not appear to be a reproductive effect, rather a direct toxic effect of repeated doses delivered dose to these animals. The actual dose is unknown, but likely higher than the adults because of the probable dual exposure routes. Taken together, this study does not meet the classification criteria of clear evidence for reproductive effects of DOTI, which suggests the study has diminished significance when extrapolating to the named substance in the classification proposal, DOTE.

Two new key studies, investigating the developmental toxicity with the DOTE substance itself focused on the most sensitive species, mice and rabbits. These new GLP studies were conducted with 96 % purity DOTE and are compliant guideline studies (OECD 414) following GLP which evaluated gestational integrity, external anomalies, soft tissue anomalies and skeletal anomalies. However, in comparison to other studies on analogues the highest dose tested using DOTE was notably lower (Table 14), maybe too low to detect a dose-response relationship and thus the highest dose tested might rather reflect the starting point of a potential dosage-related response.

A maternal and/or a developmental/reproductive LOAEL specifically for the DOTE substance can be derived from the data above. Earlier estimates of the NOAEL should be disregarded in light of the new data set for DOTE.

As noted in the text above (Chapter 10.10.5), pregnant rats exposed orally to DOTI/MOTI did not show any variations of bone formation as seen in mice and rabbits. Rat dams had no treatment-related adverse effects on their foetuses. At a high dose of 25 mg/kg/day there was marginal maternal toxicity and the adverse outcomes noted in a single high dose dam were considered to have occurred by chance.

Quantitative comparison of new data on DOTE with data on DOTI / MOTI 80:20

Table 14: Summary of Dose-Response: DOTI/MOTI 80:20 vs DOTE in Developmental Toxicity Study with Mice. This is a dose-to-dose comparison of the DOTI and DOTE content in the test material stepping down from highest to lowest dose.

<p align="center">DOTI/MOTI 80:20 Effects Observed at dose levels of 100, 67, 45, 30, and 20 mg test material/kg bw/d corresponding to the amounts of the dioctyl species given below) Mouse (Faqi et al. 2001)</p>	<p align="center">DOTE (purity 96.1 %) Effects Observed at dose levels of 60, 30, and 15 mg test material/kg bw/d corresponding to the amounts of the dioctyl species given below) Mouse (Anonymous 2014b)</p>
<p>80 mg DOTI/kg bw/day for 12 days Maternal Toxicity 1 death; resorption rate ↑ [16 %] thymus gland weight ↓ [27 %], liver weight ↓ [23 %]; corrected maternal body weight gain ↓ [43 %, mean (control): 5.1 ± 7.0 g, mean ± SD: 2.9 ± 4.8 g] Developmental Toxicity foetal body wt. ↓ [18 %]; litter size ↓ ; foetal loss ↑; major malformations related to treatment: ↑ incidence of cleft palate [9.3 %], exencephaly [7.1 %], bent ribs [35.5 %] and forelimbs [18.5 %]</p>	<p>No data for this dose level.</p>
<p>53.6 mg DOTI/kg bw/day for 12 days Maternal Toxicity resorption rate ↑ [13 %] thymus gland weight ↓ [13 % statistically not significant]; corrected body weight ↓ [10%] Developmental Toxicity– THRESHOLD* foetal body wt. ↓ [9 %]; litter size ↓; foetal loss ↑ ↑ incidence of cleft palate [5.5 %], bent ribs [6.4 %]</p>	<p>57.7 mg DOTE/kg bw/day for 13 days [higher dose of dioctyltin species, longer duration] Maternal Toxicity thymus gland weight ↓ [35 %]; corrected body weight ↓ [26.6 %] Developmental Toxicity statistically significant positive trend on percentage of post implantation loss: mean ± SD: 2.6 ± 5.6</p>
<p>36 mg DOTI/kg bw/day for 12 days Maternal Toxicity – THRESHOLD* thymus gland weight ↓ [15 %]; NO effect on body weight Developmental Toxicity – NOAEL (malformations) ↓ ossification of skull bones [25.2 %] and digits, [4.4 %] supernumerary ribs ↑</p>	<p>No data for this dose level</p>
<p>No data for this dose level</p>	<p>28.8 mg DOTE/kg bw/day for 13 days Maternal Toxicity – THRESHOLD thymus gland weight ↓ [23 %] ; marginal effect on body weight ↓ Developmental Toxicity statistically significant positive trend on percentage of post implantation loss: mean ± SD: 1.5 ± 4.9</p>
<p>24 mg DOTI/kg bw/day for 12 days Maternal Toxicity - NONE Maternal - NOAEL Developmental Toxicity supernumerary lumbar ribs ↑ [67.8 %]</p>	<p>No data for this dose level</p>
<p>16 mg DOTI/kg bw/day for 12 days Maternal Toxicity - NONE Developmental Toxicity skeletal variations: supernumerary ribs ↑</p>	<p>14.4 mg DOTE/kg bw/day for 13 days [~equal dose of dioctyltin species, longer duration] Maternal Toxicity- NONE Maternal NOAEL Developmental Toxicity statistically significant positive trend on percentage of post implantation loss: mean ± SD: 0.9 ± 2.8</p>

* In the Table the word THRESHOLD appears in capital letters, always to the right of bold text indicating either Maternal Toxicity or Developmental toxicity, e.g. in this cell of the Table 36 mg DOTI/kg bw/day is the threshold for maternal toxicity. The word THRESHOLD highlights the threshold dose for Maternal Toxicity or Developmental toxicity.

Table 15: Summary of Dose-Response: DOTI/MOTI 80:20 vs DOTE in Developmental Toxicity Study with Rabbits. This is a dose-to-dose comparison of the DOTI and DOTE content in the test material stepping down from highest to lowest dose.

<p align="center">DOTI/MOTI 80:20 Effects Observed at dose levels of 100, 10, and 1 mg test material/kg bw/d corresponding to the amounts of the dioctyl species given below) Rabbit (Battenfeld 1992)</p>	<p align="center">DOTE (purity 96.1 %) Effects Observed at dose levels of 80, 20, and 4 mg test material/kg bw/d corresponding to the amounts of the dioctyl species given below) Rabbit (Anonymous 2014a)</p>
<p>80 mg DOTI/kg bw/day for 13 days [GD6-18]</p> <p>Maternal Toxicity– THRESHOLD* incidence of abortion ↑ [4 dams], total resorptions [28.4 %]; body weight unaffected</p> <p>Developmental Toxicity mean foetal body wt. per litter ↓ [~20 %]; foetolethality [3 dams]; incidence of post implantation loss ↑ [28.4 % per group] incidence of skeletal/visceral abnormalities ↑ [~60 %]: ossification of skull bones, sternebrae and feet bones ↓; dilated renal pelvis and additional small vessels originating from the aortic arch</p>	<p>76.9 mg DOTE/kg bw/day for 23 days [GD6-28] [~equal dose of dioctyltin species, longer duration]</p> <p>Maternal Toxicity– THRESHOLD NO abortion, or resorption thymus gland weight ↓ [12.8 %]; body weight unaffected</p> <p>Developmental Toxicity – biologically relevant effect on foetal body wt. (mean -11.9 % relative to controls) and foetal crown-rump length (mean -10.7 % relative to controls) statistically significant negative trend on foetal weight and on foetal crown-rump length / 1 litter with foetal loss [within historical controls] excluded</p>
<p>8 mg DOTI/kg bw/day for 13 days [GD6-18]</p> <p>Maternal Toxicity - NOAEL Developmental Toxicity – THRESHOLD marginal retardative effect (non-significant increase): ossification of skull bones and digits ↓;</p>	<p>14.4 mg DOTE/kg bw/day for 23 days [GD6-28] [higher dose of dioctyltin species and longer duration]</p> <p>Maternal Toxicity- NOAEL Developmental Toxicity statistically significant negative trend on foetal weight and on foetal crown-rump length</p>
<p>1 mg DOTI/kg bw/day for 13 days [GD6-18]</p> <p>Maternal Toxicity - NONE Developmental Toxicity– NOEL</p>	<p>3.8 mg DOTE/kg bw/day for 23 days [GD6-28] [higher dose of dioctyltin species and longer duration]</p> <p>Maternal Toxicity- NONE Developmental Toxicity statistically significant negative trend on foetal weight and on foetal crown-rump length</p>

* In the Table the word THRESHOLD appears in capital letters, always to the right of bold text indicating either Maternal Toxicity or Developmental toxicity, e.g. in this cell of the Table 76.9 mg DOTE/kg bw/day is the threshold for maternal toxicity. The word THRESHOLD highlights the threshold dose for Maternal Toxicity or Developmental toxicity.

The NOAEL for maternal toxicity in mice is 14.4 mg DOTE/kg bw/day [GD5-17]. The NOAEL for developmental/reproductive toxicity cannot be determined as there is statistically significant positive trend on percentage of post implantation loss [GD5-17]. The NOAEL for maternal toxicity in rabbits is 76.9 mg DOTE/kg bw/day [GD6-28] and the LOAEL for developmental/reproductive toxicity is 76.9 mg DOTE/kg bw/day [GD6-28]. Therefore the overall oral NOAEL for maternal toxicity is 14.4 mg DOTE/kg/day, based on slight maternal thymus toxicity. The overall developmental NOAEL [which is defined as the highest dose producing no adverse effect] cannot be determined, but is based on increasing percentages of post implantation loss in mice and effect on foetal weight and foetal crown-rump length in rabbits.

These data support the conclusion that there are adverse developmental effects from DOTE in mice and rabbits, the most sensitive species. The developmental effects in rats with DOTI/MOTI were related to the observation that this species is less sensitive and that further testing of DOTE in rats is (a) unlikely to alter the conclusion, and (b) unnecessary for classification, since DOTI is structurally very similar to DOTE differing only slightly at the C-8 alcohol of the mercaptoester ligand and their respective mercaptoacetate esters are expected to have very similar physicochemical and toxicological properties, including hydrolysis products. However, the recently conducted developmental toxicity studies with DOTE in two species (see Table 11, Section 10.10.4) suggest that DOTI might have a higher toxicological potency with respect to developmental toxicity as supported by the Summary Tables Table 14 and Table 15 of Dose-Response.

There are no data to suggest that classification of the substance for reproductive toxicity is required, however, not all doubts could be dismissed regarding developmental toxicity of DOTE, and thus the following classification is appropriate:

CLP
Classification as Repr. 2 (H361d)

10.11 Specific target organ toxicity-single exposure

This point is not proposed for harmonization.

10.12 Specific target organ toxicity-repeated exposure

Table 16: Summary table of animal studies on STOT RE

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, route of exposure, dose levels, duration of exposure	Results	Reference
equivalent or similar to OECD TG 408 (Repeated Dose 90-Day Oral Toxicity in Rodents) supporting study 2 (reliable with restrictions) Rat (Sprague-Dawley) 4 groups of 40 rats 20 male/20 female subchronic	Test material: DOTE/MOTE Diocetyltn bis(2-EHMA) [CAS No. 15571-58-1] : Octyltin tris(2-EHMA) [CAS No. 27107-89-7] (purity 70:30 %) oral: feed 25, 50, and 100 ppm (nominal in diet) (0, 1.6, 3.3, and 6.6 mg/kg bw/day) Exposure: 90 days (continuously) Post-exposure period: 30 days	NOAEL: 25 ppm (male/female) based on: test material (At 50 and 100 ppm: significant dose-related reduction in absolute and relative thymus gland weights [~25-35 %].) 25 ppm calculated as 1.25 mg/kg bw/day, based on a food factor of 0.05.	Anonymous (1974)
equivalent or similar to OECD TG 408 (Repeated Dose 90-Day Oral Toxicity in Rodents) key study 2 (reliable with restrictions) rat (Wistar) male/female (15 rats/sex/group) subchronic	Test material: DOTE (97 %) Diocetyltn bis(2-EHMA) [CAS No. 15571-58-1] : [CAS No. 27107-89-7] Octyltin tris(2-EHMA) : Trioctyltin (2-EHMA) [CAS No. 61912-55-8] (purity 97:0.3 : 2.17 %) oral: feed 100, 500, and 1000 ppm (experiment 1) (nominal in diet) 50 and 250 ppm (experiment 2) (nominal in diet) 10 and 25 ppm (experiment 3) (nominal in diet) Exposure: 90 days (continuously)	NOAEL: 10 ppm (male/female) based on: test material (reduced thymus weight [~20 %]) 10 ppm is equivalent to 0.5 mg/kg bw/day	Anonymous (1970)

Table 17: Summary table of human data on STOT RE

Type of data/report	Test substance	Route of exposure Relevant information about the study (as applicable)	Observations	Reference
		No human data		

10.12.1 Short summary and overall relevance of the provided information on specific target organ toxicity – repeated exposure

The two subchronic studies (Anonymous, 1974 and 1970) with DOTE (CAS No. 15571-58-1) and MOTE (CAS No. 27107-89-7) at 70/30 % Dioctyltin (2-EHMA)/Monoctyltin (2-EHMA) and 97:2.17 % Dioctyltin (2-EHMA) and Monoctyltin (2-EHMA) demonstrated that the substance causes clear target effects substantiated by thymus lymphocyte depletion. A GLP deficiency for these two tests exists in that the reports do not contain information on the test substance homogeneity and stability; however other data for the octyltin family and more broadly for organotin thioesters indicate that the substances are stable in animal feed. Therefore, the studies can be considered “valid with deficiencies” and used for classification of the substance. In the summaries below, the high mortality in the 500 and 1000 ppm groups indicated these doses were too high to accurately and meaningfully assess the non-lethal adverse effects. For this reason, calculations of % change in measured parameters were done only for doses below 500 ppm.

1/ In the first subchronic diet non GLP study (Anonymous, 1970), rats were given 100, 500 and 1000 ppm (test 1), 50, 250 ppm (test 2), 10, 25 ppm (test 3) Dioctyltin (2-EHMA) during 90 days. The following effects were observed:

- Mortality: 9/15 males and 4/15 females died in the 500 ppm diet group; 15/15 males and 14/15 females died in the 1000 ppm diet group;
- Food consumption and food efficiency: slightly, but not significantly reduced at 500 and 1000 ppm.
- Haematology:
 - o Significant decrease of RBC at 100 ppm diet for males (92 % of controls), and at 500 ppm diet for females (week 6).
 - o Significant decrease in percentage of lymphocytes and neutrophils at 500 ppm diet (both sexes) (weeks 6 and 12).
 - o Significant decrease in haemoglobin content at 100 ppm diet for males (week 12) [97 % of controls], and at 500 ppm diet for females (weeks 6 and 12).
 - o Significant decrease in percentage of packed cell volume at 100 ppm diet for males [96 % of controls] and a significant increase for females [105 % of controls] at week 12.
 - o Significant decrease in percentage of packed cell volume at 500 ppm diet for females but not for males at week 12.

Because these haematological findings show small percentage differences, are not consistent across sexes, nor do they follow a clear dose-response, it is uncertain if the findings are treatment-related.

- Urinalysis: Specific gravity of the urine was significantly decreased and UGOT levels were significantly increased at 500 ppm diet (both sexes). Specific gravity of the urine of females at 100 ppm diet was also significantly decreased.
- Biochemical: The sugar content of the blood was significantly decreased in males and females at 500 ppm diet. SGOT levels were significantly increased in females at 10 ppm diet. SGPT levels were significantly increased in females at 10 ppm diet and in males at 500 ppm diet. SAP levels were significantly increased at 100 and 500 ppm diet for both sexes.
- The water content of the brain was significantly decreased at 500 ppm diet.

- Organ weights: The following statistically significant changes were observed:
 - o Terminal body weight: decreased in females at 100 ppm diet, and in males and females at 500 ppm diet;
 - o Relative heart weight: increased in females at 500 ppm diet;
 - o Relative kidney weight: increased in males and females at 500 ppm diet;
 - o Relative liver weight: increased in males at 10 ppm diet and in females at 500 ppm diet;
 - o Relative spleen weight: increased in females at 500 ppm diet;
 - o Relative brain weight: increased in males and females at 500 ppm diet;
 - o Relative gonads weight: increased in males at 500 ppm diet;
 - o Relative thymus weight: decreased in males and females at 100 and 500 ppm diet
- Histopathology: 2/5 females at 100 ppm diet, and 5/5 males and 5/5 females at 500 ppm diet had almost complete depletion of lymphocytes resulting in a very small thymus with a uniform picture of the remaining reticula parenchyma, which hardly permitted a distinction between cortex and medulla. This damage of the thymus was occasionally accompanied with little active lymph nodes and a slight reduction of splenic lymphoid cells. In the kidney, 3/5 males and 2/5 females exhibited swollen tubular epithelial cells containing a granular or finely vacuolated cytoplasm.

The NOAEL was determined to be 10 ppm diet (equivalent to 0.5 mg/kg bw/day), on the basis of reduced thymus weight at 25 ppm diet. The LOAEL was determined to be 25 ppm diet (calculated as 1.07-1.24 mg/kg bw/day in males and 1.46-1.51 mg/kg bw/day in females). Calculation of dosage was performed using body weights of 340 g (males) and 200 g (females), and average food consumption of 14.6-16.8 g/rat/day (males) and 11.7-12.1 g/rat/day (females).

2/ In the second subchronic non GLP study (Anonymous, 1974), rats were given Dioctyltin (2-EHMA)/Monoctyltin (2-EHMA) (70/30 %) at 25, 50 and 100 ppm in diet (equivalent to an average daily intake of 0, 1.6, 3.3 and 6.6 mg/kg bw/day) during 90 days. The following relevant effects were observed:

Significant dose-related reduction in absolute and relative thymus weights in the 50 ppm (3.3 mg/kg bw/day) and 100 ppm (6.6 mg/kg bw/day) dose groups. It was considered a toxicologically-relevant change in the thymus (i.e. thymotoxicity) which was in accordance with the known toxicity profile of the test substance and related octyltin substances.

The overall NOAEL was determined to be 25 ppm in the diet (calculated as 1.25 mg/kg bw/day and a food factor of 0.05) and was based on the 50 ppm threshold for thymus toxicity.

There were two recovery assessments associated with this study, a 15-day and a 30-day recovery. The specific target organ, the thymus, recovered completely within 15 days of the cessation of treatment for both the 50 and 100 ppm doses. There were no gross findings related to treatment or adverse histopathological findings in the thymus for either sex at either 15 or 30 days post-exposure.

The two new developmental toxicity studies with DOTE have been regarded for hazard classification as supporting evidence taking into account shorter exposure periods and pregnancy of the animals.

3/ In the new mice developmental toxicity study (Anonymous, 2014b), dams were given DOTE 96.1 % purity at 0, 15, 30, and 60 mg test material/kg bw/day (corresponding to 14.4, 28.8, and 57.7 mg DOTE/kg bw/day) during day 5 to 17 of pregnancy.

Please refer to section 'effects on development' for details.

Maternal toxic effects

Reduction in thymus weight from 30 mg/kg bw/day

MATERNAL DATA

- Gross Pathology [Maternal]

There was a treatment-related macroscopic finding of reduced maternal thymus weight.

The mean maternal thymus weight was statistically significantly reduced (-23 %) in the 30 mg/kg bw/day [mid] and 35 % at the 60 mg/kg bw/day [high] dose groups. The mean maternal thymus weight in the low dose mice was reduced relative to controls, but was not statistically significant. These observations are indicative of a treatment-related specific target organ toxicity resulting from exposure to the test material. No other gross pathological findings were noted in any dose group.

4/ In the new rabbit developmental toxicity study (Anonymous, 2014a), dams were given DOTE 96.1 % purity during day 6-28 of pregnancy at 0, 4, 20 and 80 mg/kg bw/day (corresponding to 3.8, 14.4, and 76.9 mg DOTE/kg bw/day).

Please refer to section 'effects on development' for details.

Maternal toxic effects

Biologically relevant depression [$> 10\%$] in thymus weight at 80 mg/kg bw/day

MATERNAL DATA

- Gross Pathology: There was a treatment-related macroscopic finding of reduced maternal thymus weight. The mean maternal thymus weights were -5.1, -9.6, and -12.8 % in the 4 mg/kg bw/day [low], 20 mg/kg bw/day [mid], and 80 mg/kg bw/day [high] dose groups, respectively when compared to controls. While these decreases were not statistically significant, the results were dose-dependent relative to controls and are consistent with data from other species [rat and mouse] which demonstrate the thymus is a target organ for octyltin compounds. These observations are indicative of a treatment-related specific target organ toxicity resulting from exposure to the test material. No other gross pathological findings were noted in any dose group.

10.12.2 Comparison with the CLP criteria

Category 1: Substances that, on the basis of evidence from studies in experimental animals, can be presumed to have the potential to produce significant toxicity in humans following repeated exposure. Substances are classified in Category 1 for target organ toxicity (repeat exposure) on the basis of:

- Observations from appropriate studies in experimental animals in which significant toxic effects, of relevance to human health, were produced at generally low exposure concentrations.

Classification in Category 1 is applicable, when significant toxic effects observed in a 90-day repeated-dose study conducted in experimental animals are seen to occur at or below the guidance values: oral (rat) $C \leq 10$ mg/kg body weight/day.

Significant toxic effects on thymus are observed in both oral 90-day studies in rats well below the guidance value for Category 1 classification.

These observations are supported by significant toxic effects on thymus in mice and treatment-related as well as dose-dependent effects on thymus in rabbits demonstrating specific target organ toxicity on thymus in several species.

10.12.3 Conclusion on classification and labelling for STOT RE

The evaluation of the repeated dose toxicity was based on four studies:

- Two subchronic oral toxicity tests (rat) with DOTE (70 and 97 % purity) - no guideline studies;

No data on repeated dose toxicity via the dermal or inhalation route are available.

- Two developmental toxicity tests (mouse and rabbit) with DOTE (96 % purity) – OECD TG 414

Based on the four studies, DOTE is proposed to be classified as STOT RE 1, H372: Causes damage to thymus through prolonged or repeated exposure. Furthermore, the tests illustrate that no adverse effects on the reproductive organs were recorded in either of these studies. These observations remain relevant for the broader assessment on the reproductive toxicity endpoint.

10.13 Aspiration hazard

Not evaluated in this dossier.

11 EVALUATION OF ENVIRONMENTAL HAZARDS

11.1 Rapid degradability of organic substances

Table 18: Summary of relevant information on rapid degradability

Method	Results	Remarks	Reference
OECD Guideline 111 (Hydrolysis as a function of pH)	25°C pH 4: half-life > 1 year pH 7: half-life > 1 year pH 9: half-life > 1 year 37°C, 0.1 N HCl pH 1.2: half-life < 1 min degradation product: DOTE (monochloride ester)	Rel. 2 no GLP-study	Anonymous (2015)
Electrospray ionization mass spectrometry	Rapid hydrolysis within minutes to hours	Rel. 3 (major methodological deficiencies)	Anonymous (2003)
OECD Guideline 301 F (Manometric Respirometry Test)	22±2°C 29-43% degradation (BOD) after 28 days Abiotic control: 0 – 10 % degradation (BOD) after 74 days Reference substance: 87 % degradation (BOD) after 13 days Inhibition control: 40-50 % degradation (BOD) after 14 days	Rel. 1 GLP-study	Anonymous (2000)
OECD Guideline 301B (CO ₂ Evolution)	22±2°C 23 % ThCO ₂ evolution after 28 days Reference substance: 75 % ThCO ₂ evolution after 13 days	Rel. 2 (inoculum concentration not mentioned) GLP-study	Anonymous (1992d)
OECD Guideline 301B (CO ₂ Evolution)	22±2°C 19 % ThCO ₂ evolution after 28 days (10.2 mg/L initial concentration) 11 % ThCO ₂ evolution after 28 days (21.4 mg/L initial concentration) Reference substance: 75 % ThCO ₂ evolution after 13 days	Rel. 2 (inoculum concentration not mentioned) GLP-study	Anonymous (1992e)

11.1.1 Ready biodegradability

Ready biodegradation of DOTE was investigated in a study conducted according to OECD Guideline 301 F using 30 mg solids/L inoculum (activated sludge, domestic, non-adapted) and 50 mg/L test substance. After 28 days 29-43 % degradation was observed. As the biodegradation of DOTE was shown to be slow, the test was prolonged to 74 days (36-68% degradation, validity criteria not fulfilled as difference of replicate values is > 20 %). Degradation of the reference substance (aniline) was 87 % after 13 days. 40-50 % degradation at day 14 was shown in the toxicity control (Anonymous, 2000).

In addition, two ready biodegradation tests according to OECD Guideline 301 B are available. In the first study inoculum from activated sludge of a wastewater treatment plant was used (adaption and concentration not specified). The initial concentration of the test substance was 11.9 mg/L and 23.8 mg/L. For both concentration 23 % ThCO₂ evolution was observed after 28 days. ThCO₂ evolution of the reference

substance (aniline) was 75 % after 13 days (Anonymous, 1992d). The second OECD Guideline 301 B study was carried out with 10.2 mg/L and 21.4 mg/L test substance. The concentration of the inoculum (activated sludge, adaption not specified) was also not mentioned. After 28 days 19 % ThCO₂ was observed for the lower initial concentration and evolution and 11% ThCO₂ for the higher initial concentration. The reference substance was degraded to 75 % after 13 days (Anonymous, 1992e).

In conclusion, DOTE is not readily biodegradable.

11.1.2 BOD₅/COD

No data available.

11.1.3 Hydrolysis

The hydrolysis of DOTE was tested according to OECD Guideline 111. After 5 days less than 10 % DOTE was hydrolyzed at 50°C (pH 4, 7, 9). The half-life for 25°C was calculated by the registrant to be > 1 year (Anonymous, 2015)). At simulated gastric conditions (0.1 M HCl, pH 1.2, 37°C) the half-life was < 1 min and 75% DOTE was hydrolysed to dioctyltin chloro 2-ethylhexylmecamptacetate (DOTE_C).

11.1.4 Other convincing scientific evidence

No data available.

11.1.4.1 Field investigations and monitoring data (if relevant for C&L)

Not relevant for this dossier.

11.1.4.2 Inherent and enhanced ready biodegradability tests

No data available.

11.1.4.3 Water, water-sediment and soil degradation data (including simulation studies)

No data available.

11.1.4.4 Photochemical degradation

Not relevant for this dossier.

11.2 Environmental transformation of metals or inorganic metals compounds

Not relevant for this dossier.

11.2.1 Summary of data/information on environmental transformation

Not relevant for this dossier.

11.3 Environmental fate and other relevant information

Based on the estimated high log Kow value (15.35) adsorption to sediment and soil is expected.

No further data available.

11.4 Bioaccumulation

Table 19: Summary of relevant information on bioaccumulation

Method	Results	Remarks	Reference
OECD Guideline 305	0.25 µg/L: BCF = 1294 L/kg (steady state) BCF = 1078.3 L/kg (steady state, lipid normalized) 2.5 µg/L BCF = 99 L/kg (steady state) BCF = 82.5 L/kg (steady state, lipid normalized)	Rel. 3 GLP-study	Anonymous (2010)

11.4.1 Estimated bioaccumulation

A Log Kow value of 15.35 was estimated (KOWWIN v 1.68, see chapter 7).

11.4.2 Measured partition coefficient and bioaccumulation test data

The bioconcentration factor (BCF) of DOTE was measured for *Onchorhynchus mykiss* using OECD Guideline 305. The study was carried out in a flow through system using the solvent acetone (100 mg/L) and two exposure concentrations (0.25 and 2.5 µg/L). After an uptake phase of 30 days the BCF values were assumed to be 1294 L/kg for the lower exposure level and 99 L/kg for the higher exposure level. The BCF values are based on the limit of quantification, as DOTE was not detected above the limit of quantification in the fish tissues. The used treatment concentrations are far above the calculated solubility of DOTE and therefore the study is regarded as invalid according to the validity criteria defined in Guideline OECD 305. Further shortcomings of the study are the high limit of quantification, no proof of a steady state concentration in fish and the use of a solvent, which should generally be avoided. It should be noted that DOTE is technically difficult to test in a BCF study based on the high calculated K_{ow} , the low calculated water solubility and the analytical challenges. However, based on the conducted study, neither a reliable BCF value, nor an estimation regarding the bioaccumulation criterion can be derived.

11.5 Acute aquatic hazard

Table 20: Summary of relevant information on acute aquatic toxicity

Method	Species	Test material	Results	Remarks	Reference
OECD 203/ EU C.1 Limit-test	<i>Danio rerio</i>	CAS 15571-58-1	96h-LC ₅₀ > 24.8 mg/L (measured)	Rel. 1 GLP-study	Anonymous (2004a)
EU Method C.1 GLP study	<i>Danio rerio</i>	CAS 15571-58-1	96h-LC ₅₀ > 5.8 mg/L (measured end conc.)	Rel. 3 (purity TS only 70%; analytical approach not specified, vehicle used, results based on end conc.)	Anonymous (1993c)
OECD 202 (1984) No vehicle used	<i>Daphnia magna</i>	D 27-231 TK 10974 “DOT bis thioglykester”	24h-EC ₅₀ > 0.11 mg/L (measured)	Rel. 3 No details on analytical method; test conc. not as high as the maximum of water solubility, test duration only 24h	Anonymous (1988a)
OECD 202 Vehicle DMSO	<i>Daphnia magna</i>	DOTE CAS 15571-58-1	48h-EC ₅₀ = 24.12 mg/L (nominal) 48h-NOEC = 0.05 mg/L (nominal)	Rel. 1 GLP-study	Anonymous (2016a)

EU Method C.2 vehicle	<i>Daphnia magna</i>	CAS 15571-58-1 TK 10315/A Irgastab 17 MOK-A	48h- EC ₅₀ = 0.17 mg/L (measured) 24h-EC ₅₀ = 0.32 mg/L (measured)	Rel.3 GLP study According to the updated registration dossier (2017-01), the test item contained 6-12% Ethylhexylthioglycolate (EHTG CAS 7659-86-1), which is more toxic than DOTE and makes the result unrepresentative. The study report itself contains no details on the content of EHTG but the registrant submitted details on the specifications of the test substance used in the test. According to these product specifications “TK 10315/A” and “Irgastab 17 MOK-A” contained 6 to 12% EHTG.	Anonymous (1993d)
OECD 201 Vehicle DMSO	<i>Pseudokirchneriella subcapitata</i>	CAS 15571-58-1 DOTE	72h-E _r C ₅₀ > 100 mg/L (nominal)	Rel. 1 GLP-study	Anonymous (2016b)
EU Method C.3 vehicle	<i>Desmodesmus subspicatus</i>	CAS 15571-58-1 TK 10315/A Irgastab 17 MOK-A	72h- E _r C ₅₀ = 0.17 mg/L (mean measured)	Rel.3 GLP study According to the updated registration dossier (2017-01), the test item contained 6-12% Ethylhexylthioglycolate (EHTG CAS 7659-86-1), which is more toxic than DOTE and makes the result unrepresentative. The study report itself contains no details on the content of EHTG but the registrant submitted details on the specifications of the test substance used in the test. According to these product specifications “TK 10315/A” and “Irgastab 17 MOK-A” contained 6-12% EHTG.	Anonymous (1993e)

11.5.1 Acute (short-term) toxicity to fish

A limit-test from the year 2004 using a WAF of 100 mg/L was conducted according to OECD Guideline 203 with *Danio rerio*. The concentration was analytically confirmed by measuring the total Sn and by using a conversion factor to conclude on the concentration of the parent substance DOT(2-EHMA). No vehicle was used. The semi-static test resulted in no effect at the limit concentration of 24.8 mg/L. The validity criteria were met.

There is also another acute toxicity test available. It is conducted in 1993 according to EU Method C.1 with *Danio rerio* under static conditions. The concentration was analytically confirmed but the method was not specified. In addition, the purity of the test substance was only 70 %. Therefore, the study is evaluated with reliability 3. No effects occurred up to a concentration of 5.8 mg/L (measured at the end of the test).

11.5.2 Acute (short-term) toxicity to aquatic invertebrates

The test conducted with DOTE from the year 2016 according to OECD Guideline 202 on *Daphnia magna*, used the vehicle DMSO (100 µL/L) and respectively a solvent control under static conditions. Four replicates per group consisting five Daphnids per replicate were used. The concentrations were analytically confirmed and remained within the acceptable range of $\pm 20\%$ of the nominal concentrations. The EC₅₀ after 48 hours was 24.12 mg/L. The 48h-NOEC was 0.05 mg/L. At 0.1 mg/L immobility occurred (LOEC). The validity criteria were met.

Another acute toxicity test with *Daphnia magna* from the year 1993 used DOTE as test item. In the update of the registration dossier (2017-01), the registrant describes that the test item contained 6-12 % ethylhexylthioglycolate (EHTG CAS 7659-86-1) which shows an acute (48 hours) *Daphnia*-toxicity (EC₅₀) of 0.38 mg/L. Therefore, he assessed the study with reliability 3. From the study report itself it is not obvious that EHTG was a constituent of the test item as the test item description was “TK 10315/A” and “Irgastab 17 MOK-A”. The registrant submitted product specifications from which show that the test item contained $9 \pm 3\%$ EHTG. The test was conducted with a vehicle and the test concentrations were confirmed analytically. The test was conducted under static conditions. The validity criteria were met. The test resulted in an EC₅₀ of 0.17 mg/L (measured concentration). Because of the test item composition the test has to be evaluated with reliability 3.

The study conducted in 1988 with *Daphnia magna* had deficiencies in the documentation. Therefore the reliability was assessed with 3 and the study cannot be used for classification.

11.5.3 Acute (short-term) toxicity to algae or other aquatic plants

One study from the year 2016 with the algae *Pseudokirchneriella subcapitata* (72h exposure) was conducted according to OECD Guideline 201 and GLP with a vehicle (0.1 % DMSO) and the test substance DOTE. It was a limit test under static conditions. Up to a concentration of nominal 100 mg/L no effects occurred. The validity criteria were met.

Another study conducted in the year 1993 with the algae *Desmodesmus subspicatus* and a vehicle according to the EU Method C.3 under static conditions resulted in an E_rC₅₀ of 0.17 and a NOE_rC of 0.04 mg/L (measured end concentrations). Similar to the acute toxicity study with *Daphnia magna* the registrant assessed this study with reliability 3 as the test item contained 6-12 % ethylhexylthioglycolate (EHTG CAS 7659-86-1) as EHTG shows an acute (72 hours) algae toxicity (EC₅₀) of 0.41 mg/L, according to the registrant. This test substance composition is not obvious from the study report itself as there was only the description “TK 10315/A” and “Irgastab 17 MOK-A” for the test substance. The registrant submitted product specifications which show that the test item contained $9 \pm 3\%$ EHTG. The validity criteria were met but the results have to be related to mean measured concentrations instead of measured end concentration. The E_rC₅₀ based on mean measured concentrations (0.03 – 0.17 – 0.69 – 3.29 – 11.17 mg/L) is 2.16 mg/L. Because of the test item composition the test has to be evaluated with reliability 3.

The study conducted in 1988 with *Desmodesmus subspicatus* had deficiencies in the documentation. Therefore, the reliability was assessed with 3 and the study cannot be used for classification.

11.5.4 Acute (short-term) toxicity to other aquatic organisms

Not available.

11.6 Long-term aquatic hazard

Table 21: Summary of relevant information on chronic aquatic toxicity

Method	Species	Test material	Results	Remarks	Reference
OECD 211	<i>Daphnia magna</i>	CAS 15571-58-1 DOT(EHMA)	21d-NOEC _{reproduction} = 1.448 mg/L (mean measured) 21d-NOEC _{parental survival+mobile offspring+body length} = 0.286 mg/L (mean measured)	Rel. 1 GLP-study	Anonymous (2004b)
EU Method C.3 vehicle	<i>Desmodesmus subspicatus</i>	CAS 15571-58-1 TK 10315/A Irgastab 17 MOK-A	72h- NOE _{rC} = 0.04 mg/L (mean measured)	Rel. 3 GLP study According to the updated registration dossier (2017-01), the test item contained 6-12% Ethylhexylthioglycolate (EHTG CAS 7659-86-1), which is more toxic than DOTE and makes the result unrepresentative. The study report itself contains no details on the content of EHTG but the registrant submitted details on the specifications of the substance used in the test. According to these product specifications “TK 10315/A” and “Irgastab 17 MOK-A” contained 6-12% EHTG.	Anonymous (1993e)
OECD 201	<i>Desmodesmus subspicatus</i>	TK 10974 (Sn-content is 23.68%)	72h- NOE _{rC} = 0.12 mg/L (initial)/ 0.057 mg/L (mean measured)	Rel. 3 Composition and purity of TS not reported	Anonymous (1988b)
OECD 201	<i>Pseudokirchneriella subcapitata</i>	CAS 1557-58-1 DOTE	72h-NOE _{rC} > 100 mg/L (nominal)	Rel. 1 GLP-study	Anonymous (2016b)

11.6.1 Chronic toxicity to fish

Not available.

11.6.2 Chronic toxicity to aquatic invertebrates

The test from the year 2004 according to OECD Guideline 211 with *Daphnia magna* was conducted under semi-static conditions. Water Accommodated Fractions (WAF) from DOTE were used but the test concentrations were analytically confirmed nonetheless. No vehicle was used. The validity criteria were met. After 21 days the test resulted in a NOEC for parental survival, reproduction of mobile offspring and body length of 0.286 mg/L based on the arithmetic mean of the measured concentrations. The test fulfils the validity criteria.

11.6.3 Chronic toxicity to algae or other aquatic plants

One study from the year 2016 over 72 hours with the algae *Pseudokirchneriella subcapitata* was conducted according to OECD Guideline 201 and GLP with a vehicle (0.1 % DMSO) and the test substance DOTE. It was a limit test under static conditions. Up to a concentration of nominal 100 mg/L no effects occurred.

Another study conducted in 1993 with the algae *Desmodesmus subspicatus* and a vehicle according to the EU Method C3 under static conditions resulted in an E_{rC}50 of 0.17 and a NOE_{rC} of 0.04 mg/L (measured end concentrations). Similar to the acute toxicity study with *Daphnia magna* the registrant assessed this study with reliability 3 as the test item contained 6-12 % ethylhexylthioglycolate (EHTG CAS 7659-86-1) as

EHTG shows an acute (72 hours) algae toxicity (EC_{50}) of 0.41 mg/L, according to the registrant. This test substance composition is not obvious from the study report as there was only the description “TK 10315/A” and “Irgastab 17 MOK-A” for the test substance. The registrant submitted product specifications from which show that the test item contained $9 \pm 3\%$ EHTG. The validity criteria were met but the results have to be related to mean measured concentrations. Therefore the NOE_rC based on mean measured concentrations (0.03 – 0.17 – 0.69 – 3.29 – 11.17 mg/L) is 0.17 mg/L. Because of the test item composition the test has to be evaluated with reliability 3.

The study conducted in 1988 with *Desmodesmus subspicatus* had deficiencies in the documentation. Therefore, the reliability was assessed with 3 and the study cannot be used for classification.

11.6.4 Chronic toxicity to other aquatic organisms

Not available.

11.7 Comparison with the CLP criteria

11.7.1 Acute aquatic hazard

Table 22: Comparison with criteria for acute aquatic hazards

	Criteria for environmental hazards	DOTE	Conclusion
Acute Aquatic Toxicity	Cat. 1: $LC_{50}/EC_{50}/ErC_{50} \leq 1$ mg/L	<u>Fish</u> : 96h- $LC_{50} > 24.8$ mg/L (measured) <u>Invertebrates</u> : 48h- $EC_{50} = 24.12$ mg/L (measured) <u>Algae</u> : 72h- $ErC_{50} > 100$ mg/L (nominal)	Not Aquatic acute 1

11.7.2 Long-term aquatic hazard (including bioaccumulation potential and degradation)

Table 23: Comparison with criteria for long-term aquatic hazards

	Criteria for environmental hazards	DOTE	Conclusion
Rapid Degradation	Half-life hydrolysis < 16 days Readily biodegradable in a 28-day test for ready biodegradability (> 70 % DOC removal or > 60 % ThCO ₂ , ThOD)	Half-life hydrolysis > 1 year 11-43 % after 28 days (ThCO ₂ and BOD) => not readily biodegradable	Not rapidly degradable
Bioaccumulation	BCF > 500 or log Kow ≥ 4	Log Kow = 15.35	Low potential for bioaccumulation (based on Weight-of-Evidence: high molecular weight, very high log Kow, very low water solubility)
Aquatic Toxicity	Table 4.1.0 (b)(iii): As there is no long-term fish toxicity test available based on acute data: Cat. 1: E(L)C ₅₀ ≤ 1 mg/L Cat. 2: E(L)C ₅₀ >1 to ≤ 10 mg/L Cat. 3: E(L)C ₅₀ >10 to ≤ 100 mg/L Table 4.1.0 (b)(i): Cat. 1: NOEC ≤ 0.1 mg/L Cat. 2: NOEC ≤ 1 mg/L	<u>Fish</u> : 96h-LC ₅₀ > 24.8 mg/L (measured) <u>Fish</u> : no long-term toxicity test available <u>Invertebrates</u> : 21d-NOEC = 0.286 mg/L (mean measured) <u>Algae</u> : 72h- NOE _r C > 100 mg/L (nominal)	Based on Table 4.1.0 (b)(iii): No aquatic chronic classification Based on Table 4.1.0 (b)(i): Aquatic chronic 2, H411 (based on 21d-NOEC _{invertebrates} = 0.286 mg/L)

11.8 CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS

DOTE is not rapidly degradable and therefore, with NOEC of 0.286 mg/L fulfils the classification criteria for hazardous to the aquatic environment “Aquatic Chronic 2”. The hazard statement code is H411.

12 EVALUATION OF ADDITIONAL HAZARDS

Not evaluated in this dossier.

13 REFERENCES

Anonymous (1970). Subchronic (90-day) toxicity studies with two organic tin compounds (Advastab 17 MOK 034 and Advastab 17 MOK 028) in albino rats. Testing laboratory: CIBA-GEIGY Corporation. Report no.: 88-920001834. Owner company: Crompton GmbH. Report date: 1970-01-01.

Anonymous (1974). 90-day dietary study in rats with compound TK 10 315. Testing laboratory: CIBA-GEIGY Ltd. Owner company: Crompton GmbH. Report date: 1974-11-06.

Anonymous (1978a). Mutagenicity Evaluation of Irgastab 17 MOK in the Ames Salmonella/Microsome Plate Test. Testing laboratory: Litton Bionetics, Inc. Report no.: LBI Project No. 20998. Owner company: Crompton GmbH. Report date: 1978-12-01.

Anonymous (1978b). Mutagenicity Evaluation of Irgastab 17 MOK in the Ames Salmonella/Microsome Plate Test. Report no.: LBI Project No. 20998. Owner company: Crompton GmbH. Report date: 1978-12-01.

Anonymous (1979). Salmonella/Mammalian-Microsome Mutagenicity Test with TK10 315. Testing laboratory: CIBA-GEIGY Ltd. Report no.: 78/2657. Owner company: Crompton GmbH. Report date: 1979-08-28.

Anonymous (1983). Salmonella/Mammalian-Microsome Mutagenicity Test. Test Material: TK 10 315 (Irgastab 17 MOK). Testing laboratory: CIBA-GEIGY Ltd. Owner company: Crompton GmbH. Study number: Project No. 830541. Report date: 1983-09-15.

Anonymous (1988a). Report on the Test for Acute Toxicity of TK 10974 to *Daphnia magna*. Testing laboratory: CIBA-GEIGY Ltd. Report no. 884103. Owner company: CIBA-GEIGY Ltd. Report date: 1988-03-17.

Anonymous (1988b). Report on the Alga, Growth Inhibition Test with TK 10974. Testing laboratory: CIBA-GEIGY, Ltd. Report no. 884104. Owner company: CIBA-GEIGY, Ltd. Report date: 1988-03-17.

Anonymous (1992a). Acute Oral Toxicity in the Rat. TK 10315 (Irgastab 17 MOK). Testing laboratory: CIBA-GEIGY Ltd. Report no.: 924128. Owner company: Crompton GmbH. Report date: 1992-11-02.

Anonymous (1992b). Acute Dermal Toxicity in the Rat. TK 10315 (Irgastab 17 MOK). Testing laboratory: CIBA-GEIGY Ltd. Owner company: Crompton GmbH. Study number: Test No. 924129. Report date: 1992-10-14.

Anonymous (1992c). Acute Dermal Irritation/Corrosion Study in the Rabbit. TK 10315/A (Irgastab 17 MOK-A). Testing laboratory: CIBA-GEIGY Ltd. Owner company: Crompton GmbH. Study number: Test No. 924131. Report date: 1992-11-10.

Anonymous (1992d). Report on the test for ready biodegradability in the Modified Sturm Test of Irgastab 17 MOK. Testing laboratory: CIBA-GEIGY Ltd. Owner company: CIBA-GEIGY Ltd.. Study number: Test No.: 928325. Report date: 1992-12-11.

Anonymous (1992e). Report on the test for ready biodegradability in the Modified Sturm Test of Irgastab 17 MOK-A. Testing laboratory: CIBA-GEIGY Ltd. Owner company: CIBA-GEIGY Ltd. Study number: Test No.: 928320. Report date: 1992-12-11.

Anonymous (1993a). Skin Sensitization Test in the Guinea Pig Maximization Test. TK 10315/A (Irgastab 17 MOK-A). Testing laboratory: Ciba-Geigy Ltd. Report no.: 924133. Owner company: Crompton GmbH. Report date: 1993-01-28.

Anonymous (1993b). Skin sensitisation test in the guinea pig Maximisation Test. TK 10315 (Irgastab 17 MOK). Testing laboratory: CIBA-GEIGY Ltd. Report no.: 924130. Owner company: Crompton GmbH. Report date: 1993-02-02.

Anonymous (1993c). Report on the acute toxicity test of Irgastab 17 MOK to zebra-fish (*Brachydanio rerio*). Testing laboratory: CIBA-GEIGY Ltd. Report no. 928323. Owner company: CIBA-GEIGY Ltd. Report date: 1993-01-14.

Anonymous (1993d). Report on the acute toxicity test of Irgastab 17 MOK-A on *Daphnia* (*Daphnia magna* Straus 1820). Testing laboratory: CIBA-GEIGY Ltd. Report no. 928319. Owner company: CIBA-GEIGY Ltd. Report date: 1993-04-21.

Anonymous (1993e). Report on the growth inhibition test of Irgastab 17 MOK-A to green algae (*Scenedesmus subspicatus*). Testing laboratory: CIBA-GEIGY Ltd. Report no. 928318. Owner company: CIBA-GEIGY Ltd. Report date: 1993-04-21.

Anonymous (1997). Two-generation reproduction toxicity study of MOTTG/DOTTG in rats by administration in the diet. Testing laboratory: LPT Laboratory of Pharmacology and Toxicology, Redderweg 8, D-21147 Hamburg. Report no.: LPT Report No. 6247/1/91. Owner company: Crompton GmbH. Report date: 1997-03-05.

Anonymous (2000). Determination of the Biodegradability of TZ 03499 in the Manometric Respirometry Test. Testing laboratory: BASF Aktiengesellschaft. Project No. 99/0607/26/1. Owner company: Chemtura Organometallics Germany. Report date: 2000-11-17.

Anonymous (2003). Electrospray Ionization Mass Spectrometric Study of Dioctyltin Compounds in Solution. Testing laboratory: not reported. Report no. not reported. Owner company: Arkema. Report date: 2003-05-01.

Anonymous (2004a). 96-Hour acute toxicity in zebra-fish with DOT(EHMA) (semi-static). Testing laboratory: NOTOX B.V. Project no. NOTOX Project 374996. TNO study number 5312/01. Owner company: Parametrix, Inc. Report date: 2004-09-03.

Anonymous (2004b). 8-Oxa-3,5-dithia-4-stannatetradecanoic acid, 10-ethyl-4,4-dioctyl-7-oxo-, 2-ethylhexyl ester [Dioctyltin bis(2-ethylhexylmercaptoacetate), CAS. No. 15571-58-1]: *Daphnia magna*, reproduction test (semi-static). Testing laboratory: NOTOX B.V. Report no. NOTOX Project 375097. TNO Study No. 5312/02. Owner company: Parametrix, Inc.

Anonymous (2010). Bioconcentration test in Rainbow Trout with Dioctyltin Bis(2-ethylhexyl Thioglycolate) (Flow-through). Testing laboratory: NOTOX B.V. Report no. 488003. Owner company: ETINSA, European Tin Stabilizers Association. Report date: 2010-03-12.

Anonymous (2014a). Prenatal developmental toxicity study of CD54-0028 administered orally in New Zealand white rabbits. Testing laboratory: Bionees India Private Limited. Report no. BIO-ET 812. Owner Company: Galata Chemicals, LLC. Report date: 2014-11-11.

Anonymous (2014b). Prenatal developmental toxicity study of CD54-0028 administered orally in Swiss albino mice. Testing laboratory: Bionees India Private Limited. Report no. BIO-ET 844. Owner Company: Galata Chemicals, LLC. Report date: 2014-11-12.

Anonymous (2015). DOTE- Hydrolysis as a Function of pH. Testing laboratory: Galata Chemicals GmbH. Report no. not reported. Owner company: Galata Chemicals GmbH. Report date: not reported.

Anonymous (2016a). Acute Immobilisation Test of Dioctyltin Bis-Ethylhexylthioglycolatge [DOTE] on *Daphnia magna* under static conditions. Testing laboratory: Bionees India Private Limited. Report no. BIO-ET 181 (M). Owner company: Galata Chemicals LLC. Report date: 2016-09-10.

Anonymous (2016b). Growth Inhibition Test of Dioctyltin bis-ethylhexylthioglycolate [DOTE] on *Pseudokirchneriella subcapitata*. Testing laboratory: Bionees India Private Limited. Report no. BIO-ET 180 (M). Owner company: Galata Chemicals LLC. Report date: 2016-09-10.

Appel, M. J. and D. H. Waalkens-Berendsen. (2004). Dichlorodioctylstannane [CASRN # 3542-36-7]: Sub-chronic (13 week) oral toxicity study in rats, including a reproduction/developmental screening study. Testing laboratory: TNO Nutrition and Food Research. Report no.: V3964, April 2004. Owner company: ORTEP ASSOCIATION. Report date: 2004-05-27.

Auletta, C. S. and Daly, I. W. (1984). Acute oral toxicity study in rats. Testing laboratory: Biodynamics, Inc. Report no.: 5108-84. Owner company: M&T Chemicals, Inc., P. O. Box 1004, Rahway, New Jersey, 07065, USA. Report date: 1984-11-08.

Baltussen E (2010). Determination of physico-chemical properties of dioctyltin bis(2-ethylhexylmercaptoacetate). Testing laboratory: NOTOX B. V., Hambakenwetering 7, 5231 DD 'S-Hertogenbosch, The Netherlands. Report no.: 492799. Owner company: ReachCentrum SPRL; Organo Tin REACH Consortium, Avenue Edmond van Nieuwenhuysse 6, 1160 BRUSSELS, Belgium. Report date: 2010-05-20.

Battenfeld, R. (1991). Embryotoxicity including teratogenicity study in the rat after daily intragastric administration from day 6 to day 15 of gestation. Testing laboratory: Schering AG. Report no.: IC 18/90, ZK 30.434. Owner company: Crompton GmbH. Report date: 1991-07-25.

Battenfeld, R. (1992). Embryotoxicity including teratogenicity study in the rabbit after daily intragastric administration from day 6 to day 18 of gestation. Testing laboratory: Schering AG. Report no.: IC 14/90, ZK 30.434. Owner company: Crompton GmbH. Report date: 1992-03-02.

CIBA-GEIGY Ltd. (1992). Acute Dermal Toxicity in the Rat. TK 10315 (Irgastab 17 MOK). Testing laboratory: CIBA-GEIGY Ltd. Owner company: Crompton GmbH. Study number: Test No. 924129. Report date: 1992-10-14.

Clarke (1994). The thymus in pregnancy: the interplay of neural, endocrine and immune influences. *Immunology today*. Vol 15. No. 11 1994.

Faqi, A. S., H. Schweinfurth, and I. Chahoud (2001). Developmental toxicity of an octyltin stabilizer in NMRI mice. *Reproductive Toxicology*. 15:117-122.

Hossack D. J. N, Richold, M. and Richardson, J. C. (1980). Micronucleus test on ZK 30 434. Testing laboratory: Huntingdon Research Centre, Huntingdon, Cambridgeshire, UK. Report no.: SHG 182/79184. Owner company: Schering A. G., D-1 Berlin 65, Postfach 65 03 11, Germany. Report date: 1980-01-22.

Kim J (2004). Benchmark dose analysis for dioctyltin dichloride (CAS 3542-36-7). Testing laboratory: Sciences International Inc. Report no.: Project Number 1502. Report date: 2004-04-21.

Krul, C. A. M. (2003). Dichlorodioctylstannane [CAS# 3542-36-7]: Micronucleus test in rat bone marrow cells. Testing laboratory: TNO Chemistry, Department of Biomolecular Sciences, 3700 AJ Zeist, The Netherlands. Report no.: V3404/14. Owner company: ORTEP. Report date: 2003-03-04.

ORTEP Association Stabilizer Task Force (2000). Summary Report - The Simulated Gastric Hydrolysis of Tin Mercaptide Stabilizers. Owner company: ORTEP Association Stabilizer Task Force. Report date: 2000-05-01.

Pelikan, Z. and E. Cerny (1970). The toxic effects of some di- and mono-n-octyl-tin compounds on white mice. *Arch. Toxikol*. 26:196-202.

Steenwinkel MJST, and Van der Horst-Groeneveld JML. (2010). Gene mutation test at the TK-locus of L5178Y cells with Dioctyltin bis(2-ethylhexylmercaptoacetate). Testing laboratory: TNO Quality of Life, Utrechtseweg 48, PO Box 360, 3700 AJ Zeist, The Netherlands. Report no.: V8403/12 (TNO study code: 8403/12). Owner company: ReachCentrum SPRL for account of the members of the OrganoTin reach consortium. Report date: 2010-04-12.

Varsho B. J. (1996). Primary dermal irritation study of dioctyltin bis (2-ethyl-hexylmercaptoacetate) in albino rabbits. WIL Research. Testing laboratory: WIL Research Laboratories, Inc. Report no.: WIL-160060. Owner company: Elf Atochem North America, Inc. Study number: WIL-160060.

Varsho, B. J. (1996). Primary Eye Irritation Study of Dioctyltin bis(2-ethyl-hexylmercaptoacetate) in Albino Rabbits. Testing laboratory: WIL Research Laboratories Inc., 1407 George Road, Ashland, Ohio, 44805-9281, USA. Report no.: WIL-160061. Owner company: Elf Atochem North America, Inc., 2000 Market Street, Philadelphia, Pennsylvania, 19103, USA. Report date: 1996-10-09.

Ward, R. J. (2003). Dioctyltin bis(2-ethylhexylmercaptoacetate): In vitro absorption through human and rat epidermis. Testing laboratory: Central Toxicology Laboratory. Report no.: CTL/JV1701/Regulatory/Report. Owner company: Tin Stabilizer Association. Report date: 2003-01-08.

Yoder, R. (2003). Electrospray Ionization Mass Spectrometric Study of Dioctyltin Compounds in Solution. Testing laboratory: not reported. Report no.: not reported. Owner company: Arkema. Report date: 2003-05-01.

Yoder R. (2000). Measurement of the hydrolysis of various organotin stabilizers under simulated gastric conditions. Ortep report. May 15.

14 ANNEXES

conf. Annex I