

# CLH report

## Proposal for Harmonised Classification and Labelling

Based on Regulation (EC) No 1272/2008 (CLP Regulation),  
Annex VI, Part 2

### International Chemical Identification:

**Dimethomorph (ISO) (E,Z) 4-(3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl)morpholine**

**EC Number:** 404-200-2  
**CAS Number:** 110488-70-5  
**Index Number:** 613-102-00-0

#### Contact details for dossier submitter:

RIVM, Bureau REACH  
PO Box 1, 3720 BA Bilthoven. The Netherlands  
[bureau-reach@rivm.nl](mailto:bureau-reach@rivm.nl)

**Version number: 3**

**Date: November 2018**

# CONTENTS


<b>1</b>	<b>IDENTITY OF THE SUBSTANCE</b> .....	<b>1</b>
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE.....	1
1.2	COMPOSITION OF THE SUBSTANCE.....	2
<b>2</b>	<b>PROPOSED HARMONISED CLASSIFICATION AND LABELLING</b> .....	<b>3</b>
2.1	PROPOSED HARMONISED CLASSIFICATION AND LABELLING ACCORDING TO THE CLP CRITERIA.....	3
<b>3</b>	<b>HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING</b> .....	<b>5</b>
<b>4</b>	<b>JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL</b> .....	<b>5</b>
<b>5</b>	<b>IDENTIFIED USES</b> .....	<b>5</b>
<b>6</b>	<b>DATA SOURCES</b> .....	<b>5</b>
<b>7</b>	<b>PHYSICOCHEMICAL PROPERTIES</b> .....	<b>5</b>
<b>8</b>	<b>EVALUATION OF PHYSICAL HAZARDS</b> .....	<b>6</b>
8.1	EXPLOSIVES.....	6
8.2	FLAMMABLE GASES (INCLUDING CHEMICALLY UNSTABLE GASES).....	6
8.3	OXIDISING GASES.....	6
8.4	GASES UNDER PRESSURE.....	6
8.5	FLAMMABLE LIQUIDS.....	6
8.6	FLAMMABLE SOLIDS.....	7
8.7	SELF-REACTIVE SUBSTANCES.....	7
8.8	PYROPHORIC LIQUIDS.....	7
8.9	PYROPHORIC SOLIDS.....	7
8.10	SELF-HEATING SUBSTANCES.....	7
8.11	SUBSTANCES WHICH IN CONTACT WITH WATER EMIT FLAMMABLE GASES.....	7
8.12	OXIDISING LIQUIDS.....	7
8.13	OXIDISING SOLIDS.....	7
8.14	ORGANIC PEROXIDES.....	7
8.15	CORROSIVE TO METALS.....	7
<b>9</b>	<b>TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)</b> .....	<b>7</b>
9.1	SHORT SUMMARY AND OVERALL RELEVANCE OF THE PROVIDED TOXICOKINETIC INFORMATION ON THE PROPOSED CLASSIFICATION(S).....	9
<b>10</b>	<b>EVALUATION OF HEALTH HAZARDS</b> .....	<b>10</b>
10.1	ACUTE TOXICITY - ORAL ROUTE.....	10
10.2	SKIN CORROSION/IRRITATION.....	11
10.3	SERIOUS EYE DAMAGE/EYE IRRITATION.....	11
10.4	RESPIRATORY SENSITISATION.....	11
10.5	SKIN SENSITISATION.....	11
10.6	GERM CELL MUTAGENICITY.....	11
10.7	CARCINOGENICITY.....	11
10.8	REPRODUCTIVE TOXICITY.....	11
10.8.1	<i>Adverse effects on sexual function and fertility</i> .....	11
10.8.2	<i>Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility</i> .....	12
10.8.3	<i>Comparison with the CLP criteria</i> .....	13
10.8.4	<i>Adverse effects on development</i> .....	14
10.8.5	<i>Short summary and overall relevance of the provided information on adverse effects on development</i> <i>16</i>	16
10.8.6	<i>Comparison with the CLP criteria</i> .....	18
10.8.7	<i>Adverse effects on or via lactation</i> .....	19
10.8.8	<i>Short summary and overall relevance of the provided information on effects on or via lactation</i> .....	19

10.8.9	Comparison with the CLP criteria .....	19
10.8.10	Conclusion on classification and labelling for reproductive toxicity.....	19
10.9	SPECIFIC TARGET ORGAN TOXICITY-SINGLE EXPOSURE .....	19
10.10	SPECIFIC TARGET ORGAN TOXICITY-REPEATED EXPOSURE .....	20
10.10.1	Short summary and overall relevance of the provided information on specific target organ toxicity – repeated exposure.....	27
10.10.2	Comparison with the CLP criteria .....	28
10.10.3	Conclusion on classification and labelling for STOT RE.....	28
10.11	ASPIRATION HAZARD.....	28
<b>11</b>	<b>EVALUATION OF ENVIRONMENTAL HAZARDS.....</b>	<b>28</b>
11.1	RAPID DEGRADABILITY OF ORGANIC SUBSTANCES .....	28
11.1.1	Ready biodegradability.....	29
11.1.2	BOD <sub>5</sub> /COD.....	29
11.1.3	Hydrolysis.....	30
11.1.4	Other convincing scientific evidence.....	30
11.1.4.1	Field investigations and monitoring data (if relevant for C&L) .....	30
11.1.4.2	Inherent and enhanced ready biodegradability tests.....	30
11.1.4.3	Water, water-sediment and soil degradation data (including simulation studies) .....	30
11.1.4.4	Photochemical degradation.....	32
11.2	ENVIRONMENTAL TRANSFORMATION OF METALS OR INORGANIC METALS COMPOUNDS .....	32
11.3	ENVIRONMENTAL FATE AND OTHER RELEVANT INFORMATION .....	32
11.4	BIOACCUMULATION .....	33
11.4.1	Estimated bioaccumulation.....	33
11.4.2	Measured partition coefficient and bioaccumulation test data .....	33
11.5	ACUTE AQUATIC HAZARD.....	33
	* Reliability according to Klimisch et al. (1997), since assessment is based on summaries in the DAR, Ri=1 is not given.....	34
11.5.1	Acute (short-term) toxicity to fish.....	35
11.5.2	Acute (short-term) toxicity to aquatic invertebrates.....	36
11.5.3	Acute (short-term) toxicity to algae and other aquatic plants.....	36
11.6	LONG-TERM AQUATIC HAZARD .....	37
	* Reliability according to Klimisch et al. (1997), since assessment is based on summaries in the DAR, Ri=1 is not given.....	39
11.6.1	Chronic toxicity to fish.....	39
11.6.2	Chronic toxicity to aquatic invertebrates.....	40
11.6.3	Chronic toxicity to algae and other aquatic plants.....	41
11.7	COMPARISON WITH THE CLP CRITERIA.....	42
11.7.1	Acute aquatic hazard .....	42
11.7.2	Long-term aquatic hazard (including bioaccumulation potential and degradation).....	42
11.8	CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS .....	43
<b>12</b>	<b>EVALUATION OF ADDITIONAL HAZARDS .....</b>	<b>43</b>
12.1	HAZARDOUS TO THE OZONE LAYER.....	43
<b>13</b>	<b>ADDITIONAL LABELLING .....</b>	<b>43</b>
<b>14</b>	<b>REFERENCES .....</b>	<b>43</b>
<b>15</b>	<b>ANNEXES .....</b>	<b>48</b>

## 1 IDENTITY OF THE SUBSTANCE

### 1.1 Name and other identifiers of the substance

**Table 1: Substance identity and information related to molecular and structural formula of the substance**

<b>Name(s) in the IUPAC nomenclature or other international chemical name(s)</b>	(E,Z) 4-[3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl]morpholine
<b>Other names (usual name, trade name, abbreviation)</b>	(E,Z) 4-[3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)-1-oxo-2-propenyl]-morpholine
<b>ISO common name (if available and appropriate)</b>	Dimethomorph (ISO)
<b>EC number (if available and appropriate)</b>	404-200-2
<b>EC name (if available and appropriate)</b>	(E,Z) 4-[3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl]morpholine
<b>CAS number (if available)</b>	110488-70-5
<b>Other identity code (if available)</b>	EINECS-No: 404-200-2 CIPAC-No: 483
<b>Molecular formula</b>	C <sub>21</sub> H <sub>22</sub> ClNO <sub>4</sub>
<b>Structural formula</b>	 <p style="text-align: center;">E-Isomer                      Z-Isomer</p>
<b>SMILES notation (if available)</b>	-
<b>Molecular weight or molecular weight range</b>	387.86
<b>Information on optical activity and typical ratio of (stereo) isomers (if applicable and appropriate)</b>	E/Z isomer ratio: ranges from 40/60 to 50/50 % w/w
<b>Description of the manufacturing process and identity of the source (for UVCB substances only)</b>	Not applicable
<b>Degree of purity (%) (if relevant for the entry in Annex VI)</b>	≥ 96.5%

## 1.2 Composition of the substance

**Table 2: Constituents (non-confidential information)**

Constituent (Name and numerical identifier)	Concentration range (% w/w minimum and maximum in multi-constituent substances)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)
Dimethomorph, CAS no 110488-70-5	96.5%	Aquatic Chronic 2; H411	Aquatic Chronic 2; H411

Besides the self-classification using EC number 404-200-2, there are also self-classifications notified under EC number 600-969-5. These notifiers mostly classify as Aquatic Chronic 2; H411 but some as Aquatic Chronic 1; H410.

**Table 3: Impurities (non-confidential information) if relevant for the classification of the substance**

Impurity (Name and numerical identifier)	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)	The impurity contributes to the classification and labelling
No relevant impurities have been identified for dimethomorph				

**Table 4: Additives (non-confidential information) if relevant for the classification of the substance**

Additive (Name and numerical identifier)	Function	Concentration range (% w/w minimum and maximum)	Current CLH in Annex VI Table 3.1 (CLP)	Current self-classification and labelling (CLP)	The additive contributes to the classification and labelling
Confidential information					

**Table 5: Test substances (non-confidential information)**

Identification of test substance	Purity	Impurities and additives (identity, %, classification if available)	Other information	The study(ies) in which the test substance is used
Dimethomorph, CAS no 110488-70-5			E/Z isomer ratio ranges from 40/60 to 50/50 % w/w In addition, acute oral and 28-day oral studies are available with the individual stereoisomers.	

## 2 PROPOSED HARMONISED CLASSIFICATION AND LABELLING

### 2.1 Proposed harmonised classification and labelling according to the CLP criteria

**Table 6: Proposed harmonised classification and labelling of dimethomorph according to the CLP criteria**

	Index No	International Chemical Identification	EC No	CAS No	Classification		Labelling			Specific Conc. Limits, M-factors	Notes
					Hazard Class and Category Code(s)	Hazard statement Code(s)	Pictogram, Signal Word Code(s)	Hazard statement Code(s)	Suppl. Hazard statement Code(s)		
Current Annex VI entry	613-102-00-0	Dimethomorph (ISO) 4-(3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl)morpholine	404-200-2	110488-70-5	Aquatic Chronic 2	H411	GHS09	H411			
Dossier submitters proposal	613-102-00-0	Dimethomorph (ISO) 4-(3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl)morpholine	404-200-2	110488-70-5	<b>Add</b> Repr. 1B <b>Retain</b> Aquatic Chronic 2	<b>Add</b> H360FD <b>Retain</b> H411	<b>Add</b> GHS08 Dgr <b>Retain</b> GHS09	<b>Add</b> H360FD <b>Retain</b> H411			
Resulting Annex VI entry if agreed by RAC and COM	613-102-00-0	Dimethomorph (ISO) 4-(3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloyl)morpholine	404-200-2	110488-70-5	Repr. 1B Aquatic Chronic 2	H360FD H411	GHS08 GHS09 Dgr	H360FD H411			

**Table 7: Reason for not proposing harmonised classification and status under public consultation**

<b>Hazard class</b>	<b>Reason for no classification</b>	<b>Within the scope of public consultation</b>
<b>Explosives</b>	hazard class not assessed in this dossier	No
<b>Flammable gases (including chemically unstable gases)</b>	hazard class not assessed in this dossier	No
<b>Oxidising gases</b>	hazard class not assessed in this dossier	No
<b>Gases under pressure</b>	hazard class not assessed in this dossier	No
<b>Flammable liquids</b>	hazard class not assessed in this dossier	No
<b>Flammable solids</b>	hazard class not assessed in this dossier	No
<b>Self-reactive substances</b>	hazard class not assessed in this dossier	No
<b>Pyrophoric liquids</b>	hazard class not assessed in this dossier	No
<b>Pyrophoric solids</b>	hazard class not assessed in this dossier	No
<b>Self-heating substances</b>	hazard class not assessed in this dossier	No
<b>Substances which in contact with water emit flammable gases</b>	hazard class not assessed in this dossier	No
<b>Oxidising liquids</b>	hazard class not assessed in this dossier	No
<b>Oxidising solids</b>	hazard class not assessed in this dossier	No
<b>Organic peroxides</b>	hazard class not assessed in this dossier	No
<b>Corrosive to metals</b>	hazard class not assessed in this dossier	No
<b>Acute toxicity via oral route</b>	hazard class not assessed in this dossier	No
<b>Acute toxicity via dermal route</b>	hazard class not assessed in this dossier	No
<b>Acute toxicity via inhalation route</b>	hazard class not assessed in this dossier	No
<b>Skin corrosion/irritation</b>	hazard class not assessed in this dossier	No
<b>Serious eye damage/eye irritation</b>	hazard class not assessed in this dossier	No
<b>Respiratory sensitisation</b>	hazard class not assessed in this dossier	No
<b>Skin sensitisation</b>	hazard class not assessed in this dossier	No
<b>Germ cell mutagenicity</b>	hazard class not assessed in this dossier	No
<b>Carcinogenicity</b>	hazard class not assessed in this dossier	No
<b>Reproductive toxicity</b>	harmonised classification proposed	Yes
<b>Specific target organ toxicity-single exposure</b>	hazard class not assessed in this dossier	No
<b>Specific target organ toxicity-repeated exposure</b>	hazard class not assessed in this dossier	No
<b>Aspiration hazard</b>	hazard class not assessed in this dossier	No
<b>Hazardous to the aquatic environment</b>	harmonised classification proposed	Yes
<b>Hazardous to the ozone layer</b>	hazard class not assessed in this dossier	No

### 3 HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING

Dimethomorph has previously been assessed for harmonised classification by TC C&L. Dimethomorph has an Annex VI entry as Aquatic Chronic 2 (H411).

Dimethomorph has previously been evaluated and authorized within the plant protection products framework. Dimethomorph is part of the AIR3 renewal programme for active substances (Commission Implementing Regulation (EU) No 844/2012). For a renewal-application under Regulation (EC) 1107/2009, the compound is currently being re-evaluated with member state The Netherlands as rapporteur member state (RMS). The RAR was peer reviewed by the Co-Rapporteur Member State Germany. This process is currently ongoing. A draft assessment report (including a proposed decision) of the Netherlands has been published by EFSA for public consultation April 10<sup>th</sup> 2018.

### 4 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

[B.] Justification that action is needed at Community level is required.

Reason for a need for action at Community level:

- *Change in existing entry due to new data*

Further detail on need of action at Community level

New data is available for reproduction toxicity: an extended one-generation reproduction toxicity study in wistar rats following administration via the diet. Based on the effects observed in this study: delayed puberty (effect on fertility or development), shortened gestation length (effect on fertility) and reduced pup weight (effect on development) it is proposed to classify dimetomorph as reproductive toxicant category 1B; H360FD.

Dimethomorph is currently classified as Aquatic Chronic 2; H411. However, classification in category 1 was proposed in the RAR. This proposal was not taken over in this CLH proposal for the reasons described in Section 11.7.2. However, seen the discussion, inclusion of the environmental hazard class within the scope of the proposal is required.

### 5 IDENTIFIED USES

Dimethomorph is an active substance of a plant protection product and is used as a fungicide.

### 6 DATA SOURCES

*This CLH report was based on the available data from the revised DAR as prepared during the renewal application of dimethomorph under Regulation (EC) 1107/2009 (published by EFSA for public consultation April 10<sup>th</sup> 2018). In addition, the data available in the registration dossier were considered. However, the level of detail of the summaries in the registration dossier (NONS) was limited.*

### 7 PHYSICOCHEMICAL PROPERTIES

**Table 8: Summary of physicochemical properties**

Property	Value	Reference	Comment (e.g. measured or estimated)
Physical state at 20°C and 101,3 kPa	white crystalline solid	RAR 2017	
Melting/freezing point	125.2-149.2°C	Registration dossier	E/Z mixture
Boiling point			Not applicable due to decomposition at 280 °C



Property	Value	Reference	Comment (e.g. measured or estimated)
Relative density	1.32 at 20°C	Registration dossier	
Vapour pressure	9.7 x 10 <sup>-7</sup> Pa for the E-isomer and 1.0 x 10 <sup>-6</sup> Pa for the Z-isomer	RAR 2017	
Surface tension	60.4 mN/m at 20 °C	Registration dossier	
Water solubility	49.2 mg/L at 20°C and pH 7 for the sum of isomers	RAR 2017	
Partition coefficient n-octanol/water	2.63 (E-isomer) and 2.73 (Z-isomer)	RAR 2017	measured, HPLC-method
Henry's law constant	5.4 x 10 <sup>-6</sup> Pa m <sup>3</sup> /mol (E-isomer) and 2.5 x 10 <sup>-5</sup> Pa m <sup>3</sup> /mol (Z-isomer)	RAR 2017	
Flash point			
Flammability	Negative	RAR 2017	
Explosive properties	Negative	RAR 2017	
Self-ignition temperature			
Oxidising properties	Negative	RAR 2017	
Granulometry			
Stability in organic solvents and identity of relevant degradation products			
Dissociation constant	-1.3	RAR 2017	calculated
Viscosity			

## 8 EVALUATION OF PHYSICAL HAZARDS

### 8.1 Explosives

*Not evaluated in this dossier*

### 8.2 Flammable gases (including chemically unstable gases)

*Not evaluated in this dossier*

### 8.3 Oxidising gases

*Not evaluated in this dossier*

### 8.4 Gases under pressure

*Not evaluated in this dossier*

### 8.5 Flammable liquids

*Not evaluated in this dossier*

**8.6 Flammable solids***Not evaluated in this dossier***8.7 Self-reactive substances***Not evaluated in this dossier***8.8 Pyrophoric liquids***Not evaluated in this dossier***8.9 Pyrophoric solids***Not evaluated in this dossier***8.10 Self-heating substances***Not evaluated in this dossier***8.11 Substances which in contact with water emit flammable gases***Not evaluated in this dossier***8.12 Oxidising liquids***Not evaluated in this dossier***8.13 Oxidising solids***Not evaluated in this dossier***8.14 Organic peroxides***Not evaluated in this dossier***8.15 Corrosive to metals***Not evaluated in this dossier***9 TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)****Table 9: Summary table of toxicokinetic studies**

Method	Results	Remarks	Reference
ADME by oral route Directive 96/54/EC B 36; OECD Guideline 417 (EPA Pesticide Assessment Guidelines, subdivision F: Hazard Evaluation: Human and Domestic Animal, § 85-1 (October 1982) claimed by the author)	Dimethomorph is rapidly absorbed in the gastrointestinal tract following oral administration to rats. The amount absorbed is limited at high dose levels. Absorbed dimethomorph is efficiently metabolised and rapidly excreted mainly via the feces. Accumulation of dimethomorph in organs and tissues did not		B.6.1.1.1a/b

CLH REPORT FOR DIMETHOMORPH

Method	Results	Remarks	Reference
	occur. The main degradation pathway was found to be the demethylation of the dimethoxyphenyl ring. To a smaller extent, degradation also occurred by oxidation of the morpholine ring.		
ADME by oral route  Directive 96/54/EC B 36; OECD Guideline 417 (EPA Guidelines, 40 CFR, Part 158,85-1 (October 1982) claimed by the author)	Dimethomorph is efficiently absorbed and metabolised in the rat. Dimethomorph is mainly excreted via the bile after conjugation to glucuronides. The main aglycone was Z67 and/or Z69.		B.6.1.1.2
ADME by oral route  Directive 96/54/EC B 36; OECD Guideline 417 (USEPA Pesticide Assessment Guideline 85-1 claimed by the author)	The rate and route of degradation is similar for male and female rat with about 95 % and > 80 % excretion of administered dose in 48 hours, respectively. Feces was the major route of excretion in both sexes and both labelled compounds (chlorophenyl-ring labelled and morpholine-ring labeled). There was no difference between metabolic profile for rats treated with the 14C-chlorophenyl and 14C-morpholine ring		B.6.1.1.3
ADME by oral route  Directive 96/54/EC B 36; OECD Guideline 417 (EPA Pesticide Assessment Guideline 85-1 (October 1982) claimed by the author).  Mass spectroscopic investigations were not conducted in compliance with GLP.	Similar urine and feces excretion patterns were found in male and female rats after single oral administration of dimethomorph at 50 mg/kg bw. The demethylation of one of the methoxy groups of the dimethoxyphenyl ring is the major metabolism pathway of dimethomorph. The presence of Z98 and Z93 confirmed also the fact that there is a second minor metabolism pathway resulting in a degradation of the morpholine ring.		B.6.1.1.4
ADME by oral route  Directive 96/54/EC B 36; OECD Guideline 417 (JMAFF Testing Guidelines for Toxicology Studies: Metabolism Study (59 NohSan No. 4200, Jan. 28, 1985) claimed by the author)  Feed and water analyses were not conducted under GLP. 0.1 % Tween 80 served as a control substance and was not characterised under GLP	At low dose absorption is quicker than at high dose. At both dose levels, the elimination occurred within 72 hours post-dosing.		B.6.1.1.5
ADME by oral route	Dimethomorph is rapidly absorbed in the gastrointestinal		B.6.1.1.6

## CLH REPORT FOR DIMETHOMORPH

Method	Results	Remarks	Reference
JMAFF Testing Guideline for Toxicity Studies: Metabolism Study (59 NohSan No. 4200, Jan. 28, 1985) claimed by the author  Minor alterations were described in the protocol.	tract following oral administration to rats. Absorbed dimethomorph is rapidly excreted. Accumulation of dimethomorph in organs and tissues did not occur.		
ADME by oral route  OECD 417, 2004/10/EC of 11 February 2004, EPA 870.7485	The metabolism of Dimethomorph in rat is very extensive. The main metabolic steps were identified as: * hydroxylation of either the dimethoxy or chlorophenyl ring and subsequent glucuronidation * demethylation of the dimethoxy ring and subsequent glucuronidation * hydroxylation and oxidative opening of the morpholine ring and subsequent conjugation * cleavage and release of the intact morpholine ring		B.6.1.1.7
In vitro metabolism study  No guideline available	All the components identified in human hepatocytes were also detected in rat and dog hepatocyte samples		Birks V. 2015

### 9.1 Short summary and overall relevance of the provided toxicokinetic information on the proposed classification(s)

Short summary, as presented in the RAR:

Dimethomorph following a single oral dose to both sexes at the low dose level is rapidly and almost completely absorbed (urine, bile and residual carcass). The excretion balance of dimethomorph demonstrates that approximately 3-17% of the applied radioactivity is excreted via the urine at 10, 50 and 500 mg/kg bw (B.6.1.1.1a, B.6.1.1.3, B.6.1.1.4, B.6.1.1.7). Excretion via the feces accounts for 80 – 90 % of the dose for non-cannulated rats (B.6.1.1.1a, B.6.1.1.2, B.6.1.1.3, B.6.1.1.4, B.6.1.1.7).

In a study with bile cannulated rats, more than 90 % of the administered dose at 10 mg/kg bw was eliminated via the bile within 24 hours with a short half-life of about 3 hours (B.6.1.1.2). At higher dose levels (500 mg/kg bw), absorbed dimethomorph in both sexes was limited as indicated by lower amounts eliminated via the bile (31-49%).

In a new study biliary excretion amounted to 61-88% of the administered dose at 10 mg/kg bw and 46-60% for the 250 mg/kg bw dose group (B.6.1.1.7). Urinary excretion in this study amounted to 3-5 % dose in males and 10-23 % of the dose in females. Based on the this study the total absorbed dose is considered to be 80-90% for the low dose group and approximately 60% for the high dose group.

Dimethomorph was found to be primarily metabolised by demethylation of one of the methoxy groups. Further conjugation of the resulting metabolites could be identified as the major radioactive residues in the urine as well as in the bile of rats. Oxidation of the morpholine ring was found as a

second minor degradation pathway in the rat. This step was considered to be the first in the pathway resulting in the cleavage and further degradation of the morpholine ring. This assumption was further substantiated by the identification of metabolite CUR 7117, and characterisation of traces of CUR 7586, CUR 7216 as well as Z43.

In addition to the previously peer-reviewed data, a new rat study focussing on the identification and quantification of metabolites in urine, faeces, bile, plasma and tissues after administration of dimethomorph (in two radiolabeled forms: chlorophenyl and morpholine ring label) was submitted for the purpose of the renewal, which confirms and extends the knowledge obtained from previous investigations.

The new study shows an extensive metabolism of dimethomorph with over 100 identified structures. The observed metabolic pathway is in good agreement with the information coming from the already peer reviewed studies. Metabolite CUR 7586 (=M550F010), depicted in the pathway based on previous data, was found in its other isomeric form as M550F051. Metabolite Z43 (=M550F005) from the previous pathway was observed in its single or double demethylated form as M550F039, M550F040 and M550F037, respectively. All other previously identified metabolites from the rat were detected in the new study as well. This confirms that all metabolic steps observed in the previous studies (with the chlorophenyl radiolabel) are observed in the new study (with chlorophenyl and morpholine label) as well.

The following main transformation steps were observed in rats:

- o hydroxylation of either the dimethoxy or chlorophenyl ring and subsequent glucuronidation (Metabolic steps 1 and 2)
- o demethylation of the dimethoxy ring and subsequent hydroxylation and/or glucuronidation (Metabolic step 3)
- o hydroxylation of the morpholine ring and subsequent modification (further hydroxylation, ring opening, degradation, conjugation) (Metabolic step 5)
- o cleavage and release of the intact morpholine ring (Metabolic step 4)

The combination of these reactions followed by conjugation steps results in a huge number of metabolites.

The proposed metabolic pathway is provided in Annex 1 (Figure 1).

In a comparative *in vitro* metabolism study of dimethomorph, incubation of dimethomorph with liver hepatocytes of humans, rats and dogs showed similar metabolite patterns and MS analysis demonstrated that all components formed by human hepatocytes were also observed with rat and dog hepatocytes. Thus, there have not been found any human specific metabolites which were not also observed in the tested animal species.

## 10 EVALUATION OF HEALTH HAZARDS

### Acute toxicity

#### 10.1 Acute toxicity - oral route

*Not evaluated in this dossier*

**10.2 Skin corrosion/irritation***Not evaluated in this dossier***10.3 Serious eye damage/eye irritation***Not evaluated in this dossier***10.4 Respiratory sensitisation***Not evaluated in this dossier***10.5 Skin sensitisation***Not evaluated in this dossier***10.6 Germ cell mutagenicity***Not evaluated in this dossier***10.7 Carcinogenicity***Not evaluated in this dossier***10.8 Reproductive toxicity****10.8.1 Adverse effects on sexual function and fertility****Table 10: Summary table of animal studies on adverse effects on sexual function and fertility**

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels, duration of exposure	Results	Reference
Dietary two generation reproductive study Directive 96/54/EC B34, OECD Guideline 416; (US EPA Guideline 83-4; Japanese MAFF 59 NohSan 4200 claimed by the author) Sprague Dawley rats, P1:	Dimethomorph (SAG 151); Batch No. DW 11/86; purity 96.6 % 0, 100, 300 and 1000 ppm (Males: 6.9, 21 and 69 mg/kg bw/day; Females: 8.0, 24 and 79 mg/kg bw/day) 100 days pre-mating-sacrifice	<u>Parental toxicity:</u> 1000 ppm: reductions in pre-mating body weight gains for P1 and F1 females.  <u>Sexual function/fertility:</u> 1000 ppm: shortened pregnancy duration for P1 and F1 females (not statistically significant)  NOAEL for parental toxicity: 300 ppm (equivalent to 21 mg/kg bw/day) NOAEL for fertility/sexual function: 1000 ppm (equivalent to 69/79 mg/kg bw/day)	B.6.6.1.1 Reliability 1

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
30/sex/dose; F1: 25/sex/dose			
Dietary extended one-generation reproduction toxicity study OECD 443 CrI:WI(Han) Wistar rats, 25/sex/dose	Dimethomorph (BAS 550 F; Batch: COD-001646; Purity 99.7%) 0, 300, 800 and 1600 ppm (26, 70, 144 mg/kg bw/day) 75 days pre-mating-sacrifice	<u>Parental toxicity:</u> ≥ 800 ppm: decreased food consumption and body weight/body weight gain, decreased seminal vesicle weight, clinical-chemical changes and pathological evidence of liver toxicity  <u>Sexual function/fertility:</u> 1600 ppm: reduced gestational length  NOAEL for parental toxicity: 300 ppm (equivalent to 26 mg/kg bw/day) NOAEL for fertility/sexual function: 800 ppm (equivalent to 70 mg/kg bw/day)	B.6.6.1.2 Reliability 1

### 10.8.2 Short summary and overall relevance of the provided information on adverse effects on sexual function and fertility

In a two-generation reproduction toxicity study conducted with Sprague-Dawley rats, dimethomorph technical was administered via the diet at doses of 100, 300 and 1000 ppm (6.9, 21 and 69 mg/kg bw/day for males; 8.0, 24 and 79 mg/kg bw/day for females). No treatment related effects on mortality or clinical signs of toxicity were observed. Body weight gains during the pre-mating treatment period for P1 females in the 1000 ppm group were reduced 14.7 % as compared to controls. Although not statistically significant, a decrease in body weight gain of 6.8 % was observed during the pre-mating treatment period for F1 females in the 1000 ppm group as compared to controls. Dimethomorph did shorten the gestational length at the top dose in P0 as well as F1 dams for the generation of the F2a offspring (statistically significant using ANOVA). However, this effect was only slight and, based on a re-evaluation using the Dunnett test, not statistically significant. No other effects on fertility parameters were observed. The NOAEL for parental toxicity was 300 ppm (equivalent to 21 mg/kg bw/day), based on reductions in pre-mating body weight gains for P1 and F1 females in the 1000 ppm group. The NOAEL for fertility and reproductive function, was 1000 ppm (highest dose tested).

In an extended one generation study, the NOAEL (no observed adverse effect level) for general systemic toxicity is 300 ppm (approx. 26 mg/kg bw/day), based on decreased food consumption and body weight/body weight gain, decreased seminal vesicle weight, as well as clinical-chemical changes and pathological evidence of liver toxicity at 800 and/or 1600 ppm (approx. 70 and 144 mg/kg bw/day), in the F0 parental animals and adult F1 offspring. The NOAEL for fertility and reproductive performance for the parental rats is 800 ppm (approx. 70 mg/kg bw/day), based on a significantly reduced duration of gestation (outside historical control range) (Table 11).

**Table 11. Gestational length in the 2 generation and extended one generation study in rats**

Gestation	Dose level (ppm)	HC
-----------	------------------	----

length	0	100	300	800	1000	1600	
2 gen F1	22.0 +/- 0.3	22.1 +/- 0.3	21.9 +/- 0.4		21.8 +/- 0.4**		
2 gen F2a	21.9 +/- 0.4	21.9 +/- 0.3	22.1 +/- 0.3		21.7 +/- 0.5**		
2 gen F2b	21.9 +/- 0.2	22.2 +/- 0.5	21.9 +/- 0.3		21.8 +/- 0.4		
EOGRT	22.3		22.2	22.0		21.4**	21.5- 22.3

\*p<0.05 (ANOVA)

\*\*p<0.01 (Dunnett test)

# upon re-evaluation using Dunnett test, statistical significance was not reached.

In addition, in the 90-day and 1-year dog repeated dose toxicity studies (see 10.10), an increase in prostate weight and fibrosis was observed. In addition, there was an increase in testes weight in the 1-year dog study. This could be considered as indicative for possible effects on the fertility and reproductive function possibly requiring classification. These effects were observed in presence of other general toxicity and it is unclear whether the effects on the male reproductive organs were a direct effect of dimethomorph or secondary to the general toxicity. No such effects were observed in the available repeated dose studies with rats and mice.

### 10.8.3 Comparison with the CLP criteria

Since no human studies are available for effects on fertility, classification in Repr. 1A is not appropriate.

The classification of a substance in Category 1B is largely based on data from animal studies. Such data shall provide 'clear evidence of an adverse effect on sexual function and fertility in the absence of other toxic effects, or if occurring together with other toxic effects the adverse effect on reproduction is considered not to be a secondary non-specific consequence of other toxic effects. However, when there is mechanistic information that raises doubt about the relevance of the effect for humans, classification in Category 2 may be more appropriate'.

According to the guidance on the application of the CLP criteria 'Effects on sexual function and fertility include, but are not limited to, **alterations to the female and male reproductive system**, adverse effects on onset of puberty, gamete production and transport, reproductive cycle normality, sexual behaviour, fertility, **parturition**, pregnancy outcomes, premature reproductive senescence, or modifications in other functions that are dependent on the integrity of the reproductive systems'.

In the dog 90-day and 1-year study increased prostate weight combined with prostatic interstitial fibrosis was observed. Since these effects were observed in presence of other general toxicity, but it cannot be excluded that they are primary effects on the reproductive system, they would warrant classification in Category 2. This is further justified as no such effects were observed in the repeated dose studies with rats and mice.

In the reproductive toxicity studies no effect on the mating index, fertility index, gestation index, live birth index and sperm parameters was observed. However, in the 2 generation study, dimethomorph did shorten the gestational length at the top dose in P0 (21.8 days vs 22.0 days in control group) as well as F1 dams for the generation of the F2a offspring (21.7 days vs 21.9 days in control group) (statistically significant using ANOVA). It is noted that this effect was only slight and, based on a re-evaluation using the Dunnett test, not statistically significant. A statistically significant (Dunnett test) reduction in gestation length was however observed in the high dose group of the extended one generation study (21.4 days vs 22.3 days in control group) which was below the lower range of the historical control data (21.5 days). A shortened gestation length can result in adverse effects such as



decreased fetal weight (as shown in the extended generation study). In addition, the effect is also relevant for humans. Since there is evidence for a statistically significantly reduced gestation length in the extended one generation study (supported by a slight, though not significantly, reduced gestation length in the 2-generation study), we conclude that there is clear evidence for an adverse effect on sexual function or fertility, which is relevant for humans. The reduced gestation length was observed in the presence of maternal toxicity. The maternal effects in the EOGRT at the highest dose consisted of decreased food consumption and body weight/body weight gain, as well as clinical-chemical changes and pathological evidence of liver toxicity. The reduced gestation length is unlikely to be secondary to the reductions in food consumption, body weight and body weight gain as food restriction in rats results in an increase in the gestation length (Chernoff, 2009 table 2). As the reduced gestation length could have resulted in the observed reduced pup weight and development, this is considered an adverse and severe effect as in humans it is known that reduced birth weight is related to other adverse effects. Therefore, dimethomorph should be classified as **Repr 1B for effects on sexual function and fertility.**

#### 10.8.4 Adverse effects on development

**Table 12: Summary table of animal studies on adverse effects on development**

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
Dietary two generation reproductive study Directive 96/54/EC B 34, OECD Guideline 416; (US EPA Guideline 83-4; Japanese MAFF 59 NohSan 4200 claimed by the author) Sprague Dawley rats, P1: 30/sex/dose; F1: 25/sex/dose	Dimethomorph (SAG 151); Batch No. DW 11/86; purity 96.6 % 0, 100, 300 and 1000 ppm (Males: 6.9, 21 and 69 mg/kg bw/day; Females: 8.0, 24 and 79 mg/kg bw/day) 100 days pre-mating-sacrifice	<u>Parental toxicity:</u> 1000 ppm: reductions in pre-mating body weight gains for P1 and F1 females.  <u>Development:</u> No adverse effects observed  NOAEL for parental toxicity: 300 ppm (equivalent to 21 mg/kg bw/day) NOAEL for development: 1000 ppm (equivalent to 69 mg/kg bw/day)	B.6.6.1.1 Reliability 1
Dietary extended one-generation reproduction toxicity study OECD 443 CrI:WI(Han) Wistar rats,	Dimethomorph (BAS 550 F; Batch: COD-001646; Purity 99.7%) 0, 300, 800 and 1600 ppm (26, 70, 144 mg/kg bw/day)	<u>Parental toxicity:</u> ≥ 800 ppm: decreased food consumption and body weight/body weight gain, decreased seminal vesicle weight, clinical-chemical changes and pathological evidence of liver toxicity  <u>Development:</u> ≥ 800 ppm: decreased anogenital distance/index in males (pup-	B.6.6.1.2 Reliability 1

CLH REPORT FOR DIMETHOMORPH

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels, duration of exposure	Results	Reference
25/sex/dose	75 days pre-mating-sacrifice	based data), reduced seminal vesicle and prostate weight 1600 ppm: decrease in preweaning pup body weight, delay of puberty in males  NOAEL for parental toxicity: 300 ppm (equivalent to 26 mg/kg bw/day) NOAEL for development: 300 ppm (equivalent to 26 mg/kg bw/day)	
Preliminary Oral Developmental toxicity Study  Guideline not specified  female Sprague-Dawley rats, 8/dose group	Dimethomorph; ZTH 236Z50; Batch No. T2/85; purity 98.7 %  50, 120 or 300 mg/kg bw/day  GD 6 to 15	No treatment-related effects observed  NOAEL for maternal toxicity: 300 mg/kg bw/day NOAEL for development: 300 mg/kg bw/day	B.6.6.2.1  Reliability 3
Preliminary Oral Developmental toxicity Study  Guideline not specified  female Sprague-Dawley rats, 4/dose group	Dimethomorph; CME151, Batch No. not specified; purity not specified  150 or 300 mg/kg bw/day  GD 6 to 15	≥ 150 mg/kg bw: Intra-uterine death, early resorptions, 300 mg/kg bw: reduced fetal weight  NOAEL for maternal toxicity: not determined NOAEL for development: < 150 mg/kg bw/day	B.6.6.2.2  Reliability 3
Oral teratogenicity study  Directive 96/54/EC B 31; OECD Guideline 414; (US EPA Guideline 83-3 claimed by the author)  Female Sprague-Dawley rats, 30 /group	Dimethomorph (SAG 151); Batch No. DW 11/86; purity 96.6 %  0, 20, 60 and 160 mg/kg bw/day  GD 6 to 15	<u>Maternal toxicity:</u>  160 mg/kg bw/day: reduced food consumption, reduced body weight and body weight gain  <u>Development:</u>  160 mg/kg bw/day: total litter loss in two animals  NOAEL for maternal toxicity: 60 mg/kg bw/day NOAEL for development: 60 mg/kg bw/day NOAEL for teratogenicity: > 160 mg/kg bw/day	B.6.6.2.3  Reliability 1
Preliminary	Dimethomorph;	<u>Maternal toxicity:</u>	B.6.6.2.4

Method, guideline, deviations if any, species, strain, sex, no/group	Test substance, dose levels duration of exposure	Results	Reference
<p>Oral Developmental toxicity Study</p> <p>The author claimed that the study was conducted according to guideline No. 83-3 of the "U.S. EPA Pesticide Assessment Guidelines, Subdivision F (November 1982)"</p> <p>Female New Zealand White rabbits, 8 or 9/group</p>	<p>CME 151; ZTH 236Z50; purity 98.7 %</p> <p>0, 300, 600 or 1000 mg/kg bw/day</p> <p>GD6-18</p>	<p>≥ 300 mg/kg bw/day: reduced body weight gain</p> <p>≥ 600 mg/kg bw/day: reduced food consumption</p> <p><u>Development:</u></p> <p>≥ 600 mg/kg bw/day: reduced fetal weight</p> <p>1000 mg/kg bw/day: high rate of abortions and increased number of intra-uterine deaths</p> <p>NOAEL for maternal toxicity: &lt; 300 mg/kg bw/day</p> <p>NOAEL for development: 300 mg/kg bw/day</p> <p>NOAEL for teratogenicity: &gt; 600 mg/kg bw/day</p>	<p>Reliability 3</p>
<p>Oral Developmental toxicity Study</p> <p>Directive 96/54/EC B31, OECD Guideline 414, (US EPA Guideline 83-3 claimed by the author)</p> <p>Female New Zealand White rabbits, 22/dose</p>	<p>Dimethomorph (SAG 151); Batch No. DW 11/86; purity 96.6 %</p> <p>0, 135, 300 or 650 mg/kg bw/day</p> <p>GD6-18</p>	<p><u>Maternal toxicity:</u></p> <p>650 mg/kg bw/day: reduced food consumption, reduced body weight gain</p> <p><u>Development:</u></p> <p>650 mg/kg bw/day: slightly increased abortion rate</p> <p>NOAEL for maternal toxicity: 300 mg/kg bw/day</p> <p>NOAEL for development: 300 mg/kg bw/day</p> <p>NOAEL for teratogenicity: &gt; 650 mg/kg bw/day</p>	<p>B.6.6.2.5</p> <p>Reliability 1</p>

### 10.8.5 Short summary and overall relevance of the provided information on adverse effects on development

In a two-generation reproduction toxicity study conducted with Sprague-Dawley rats, dimethomorph technical was administered via the diet at doses of 100, 300 and 1000 ppm (6.9, 21 and 69 mg/kg bw/day for males; 8.0, 24 and 79 mg/kg bw/day for females). No treatment related effects on mortality or clinical signs of toxicity were observed. Body weight gains during the pre-mating treatment period for P1 females in the 1000 ppm group were reduced 14.7 % as compared to controls. Although not statistically significant, a decrease in body weight gain of 6.8 % was observed during the pre-mating treatment period for F1 females in the 1000 ppm group as compared to controls. No effects were

observed on pup survival or mean pup weight. In the 1000 ppm group, the percentage of pups in the F1, F2a and F2b generations which achieved incisor eruption was reduced on one or more days from postnatal days 9 - 11 when compared to controls and the differences were statistically significant. However, the delay in incisor eruption did not interfere with the development of feeding ability and is therefore not considered an adverse finding. There were no other treatment-related developmental effects observed. The NOAEL for parental toxicity was 300 ppm (equivalent to 21 mg/kg bw/day), based on reductions in pre-mating body weight gains for P1 and F1 females in the 1000 ppm group. The NOAEL for development was 1000 ppm (equivalent to 69 mg/kg bw/day), the highest dose administered.

In an extended one generation study, the NOAEL (no observed adverse effect level) for general systemic toxicity is 300 ppm (approx. 26 mg/kg bw/day), based on decreased food consumption and body weight/body weight gain, decreased seminal vesicle weight, as well as clinical-chemical changes and pathological evidence of liver toxicity at 800 and/or 1600 ppm (approx. 70 and 144 mg/kg bw/day), in the F0 parental animals and adult F1 offspring. No effects were observed on pup survival. At 1600 ppm, mean pup body weight was 13% below control at PND 1 and still 9% below controls at PND 21. Anogenital (AG) distance of male pups (pup-based data) was statistically significantly reduced at all dose levels; 2, 6 and 10% below control, respectively (below historical control values in mid and high dose). When corrected for body weight (AG index) the reduction was 2, 4 and 5%, respectively (below historical control values in high dose only). AG distance of all female treated pups was also statistically significantly decreased, by 2, 3 and 6%, respectively. When corrected for body weight (AG index) only the low and mid-dose pups were significantly below control, and without dose response. An additional evaluation of these data, based on litter data (as provided by the applicant upon a request by EFSA during the renewal application), showed that for males only the changes at the top dose are statistically significant and slightly outside the historical controls. For females, the analysis based on litter data confirmed that no treatment-related effects are noticed. Furthermore, a delay of puberty onset at 800 ppm (males) and 1600 ppm (both sexes) was observed. However, the delay in vaginal opening was concluded to be due to a decrease in body weight. The same was concluded for the delay in preputial separation in mid dose males. However, for the high dose group a specific effect on preputial separation could not be excluded. In addition, reductions in absolute and relative seminal vesicle and prostate weight were observed in the adult F1 at 800 and 1600 ppm without histopathological changes. The NOAEL for developmental toxicity in the F1 progeny is 300 ppm (approx. 26 mg/kg bw/day), due to the decreased anogenital distance/index at 800 ppm (approx. 70 mg/kg bw/day) based on pup-based data.

Developmental toxicity studies with dimethomorph, conducted in Sprague-Dawley rats (day 6 to 15 of gestation) and in New Zealand White rabbits (day 6 to 18 of gestation), showed no evidence of teratogenic effects for fetuses, and no evidence of developmental toxicity in the absence of maternal toxicity. In the rat developmental toxicity study, the NOAEL for maternal toxicity was 60 mg/kg bw/day based on decreased body weights, body weight gains, and food consumption at 160 mg/kg bw/day (highest dose tested). The NOAEL for developmental toxicity was also 60 mg/kg bw/day, based on a slightly increased number of total litter losses at 160 mg/kg bw/day. In the rabbit developmental toxicity study, the NOAEL for maternal toxicity was 300 mg/kg bw/day based on decreased body weight gains and food consumption at 300 mg/kg bw/d and a slightly increased abortion rate at 650 mg/kg bw/day. The NOAEL for developmental toxicity was 300 mg/kg bw/day based on a slightly increased embryoletality presenting as abortion. As important parts of the development of the reproductive organs happens after gestation day 15, these effects were not fully studied in these tests.

Dimethomorph was anti-androgenic in the Yeast Androgen Screening (YAS) assay using the hAR yeast strain (Woitkowiak, 2011d). No androgenic effects were observed. In a Yeast Estrogenic Screening (YES) assay using the hER $\alpha$  yeast strain no estrogenic nor anti-estrogenic effects were observed (Woitkowiak, 2011e). These results are supported by public literature studies where dimethomorph did not insert an estrogenic effect in the E-screen Assay in MCF-7 human breast cancer cells (Bitsch, 2002) but did result in anti-androgenic effects in MDA-kB2 reporter gene assays and the YAS assay (Orton, 2011; Orton, 2014; Archer, 2015). In a public literature study reporting on the Phase I of the ToxCast program dimethomorph was indicated to have no effects on estrogen or thyroid pathways but was

reported to have a potential effect on the androgen pathway. A summary of the *in vitro* studies is included in Table 13: Summary of the *in vitro* on endocrine disruption.

**Table 13: Summary of the *in vitro* on endocrine disruption**

Test system	Result	Reference
Yeast androgen screening (YAS-assay) with hAR yeast strain	Anti-androgenic: positive Androgenic: negative	Woitkowiak, 2011d
Yeast estrogen screening (YES-assay) with hER $\alpha$ yeast strain	Anti-estrogenic: negative Estrogenic: negative	Woitkowiak, 2011e
E-Screen assay with MCF-7 breast cancer cells	Estrogenic: negative	Bitsch, 2002
YAS assay and MDA-kb2 cell line	Anti-androgenic: positive Androgenic: negative	Orton, 2011
Battery of 467 <i>in vitro</i> high-throughput screening assays (Phase I of ToxCast)	No effect on estrogen pathway or thyroid identified. Potential effect on androgen pathway	Reif, 2010
MDA-kb2 cells	Potential anti-androgen	Orton, 2014
YAS assay	Weak anti-androgen	Archer, 2015

### 10.8.6 Comparison with the CLP criteria

Since no human studies are available for effects on development on the offspring, classification in Repr. 1A is not appropriate.

The classification of a substance in this Category 1B is largely based on data from animal studies. Such data shall provide ‘clear evidence of an adverse effect on development in the absence of other toxic effects, or if occurring together with other toxic effects the adverse effect on reproduction is considered not to be a secondary non-specific consequence of other toxic effects. However, when there is mechanistic information that raises doubt about the relevance of the effect for humans, classification in Category 2 may be more appropriate’.

According to the CLP criteria ‘Developmental toxicity includes, in its widest sense, any effect which interferes with normal development of the conceptus, either before or after birth, and resulting from exposure of either parent prior to conception, or exposure of the developing offspring during prenatal development, or postnatally, to the time of sexual maturation’. It should be noted that the adverse effects must be induced during pregnancy, or as a result of parental exposure.

In the extended one-generation a delay in preputial separation was observed at the high dose level of 1600 ppm (5.9 days). In addition, decreased anogenital length was observed in males (F1) at the 800 ppm and 1600 ppm (-6.3% and -9.5%; pup-based data). Both effects were outside of the historical control range.

A delay in preputial separation in days secondary to reduced maternal body weight is normally observed as a reduction in body weight for reaching preputial separation (Carney et al, 2004, Toxicological sciences 82: 237-249 table 3). However, with dimethomorph there is also an increase in body weight for reaching preputial separation. The same applies to anogenital distance.

A decrease in absolute and relative seminal vesicle and prostate weight was observed in the adult F1 at 800 and 1600 ppm without histopathological changes.

The reduced anogenital distance and delayed sexual maturation appear to reflect the anti-androgenic effects of dimethomorph. Dimethomorph was anti-androgenic in the Yeast Androgen Screening (YAS) assay using the hAR yeast strain and in the MDA-kB2 reporter gene assay. In the extended-one generation a reduction in absolute and relative seminal vesicle weight was observed which also gives an indication of the anti-androgenic properties of dimethomorph. However, some other typical effects of anti-androgenic substances such as nipple retention were not observed. However, as there was 95% nipple retention on day 12 and 0% on day 20, this may have been difficult to detect.

In the same study, at 1600 ppm, mean pup body weight was 13% below control at PND 1 and still 9% below controls at PND 21.

In the 2-generation study, AGD and the time until puberty were not determined.

Where a causal relationship between maternal and developmental toxicity has been established, the level of concerns for the developmental effects is reduced. Therefore, the observed effects should be considered in relation to possible maternal toxicity. At the high dose level (1600 ppm) maternal toxicity consisted of slightly reduced maternal body weight (-5.7%), reduced body weight gain (-10.7%). Food consumption was also reduced in females (max -7.6%) but only during the lactation period. Increased liver weight (+14%) was observed in males and females. In females minimal or slight centrilobular hepatocellular hypertrophy was noted as well as apoptotic hepatocytes and an increase in the severity of lymphoid infiltration. In short, while some maternal toxicity was observed the effects are not considered to be severe. Based on the individual pup data it does not appear that the observed effect on preputial separation and anogenital distance only occurs in litters with higher maternal toxicity. Moreover, based on the *in vitro* information a non-maternally mediated mode of action through the anti-androgenic properties of dimethomorph seems likely. The observed effects are therefore considered to be related to the anti-androgenic effect of dimethomorph and not secondary to maternal toxicity. This mode of action is considered to be relevant to humans and therefore classification for developmental toxicity is considered necessary.

The severity of the observed effects can be questioned as they could be considered a delay in normal development. However, the type of effects and the *in vitro* results indicate that the observed effects are due to an anti-androgenic mechanism although not all effects typically observed with anti-androgens were observed in this study. It is considered likely that the anti-androgenic mechanism will also induce other adverse effects not observed in standard studies. This includes effects on the brain by *in utero* exposure to anti-androgens such as phthalates (Miodovnik, 2014).

Based on reduced anogenital distance, delayed puberty, reduced pup weight and reduced seminal vesicle and prostate weight not secondary to maternal toxicity, we conclude that there is clear evidence for an adverse effect on development, which is relevant for humans. Therefore, dimethomorph should be classified as **Repr 1B for effects on development**.

#### **10.8.7 Adverse effects on or via lactation**

There are no effects meeting the CLP criteria for effects on or via lactation

#### **10.8.8 Short summary and overall relevance of the provided information on effects on or via lactation**

There are no effects meeting the CLP criteria for effects on or via lactation

#### **10.8.9 Comparison with the CLP criteria**

There are no effects meeting the CLP criteria for effects on or via lactation

#### **10.8.10 Conclusion on classification and labelling for reproductive toxicity**

Based on the reduced gestation length, delayed puberty and reduced pup weight observed in rat studies, we conclude that dimethomorph should be classified as **Repr. 1B; H360FD**

### **10.9 Specific target organ toxicity-single exposure**

*Not evaluated in this dossier*

**10.10 Specific target organ toxicity-repeated exposure**

The information in this chapter is provided to support the assessment for reproductive toxicity and is not compared to the STOT RE criteria.

**Table 14: Summary table of animal studies on STOT RE**

<b>Method, guideline, deviations if any, species, strain, sex, no/group</b>	<b>Test substance, route of exposure, dose levels, duration of exposure</b>	<b>Results</b>	<b>Reference</b>
<p>Oral 28 day study</p> <p>Method not specified in the report; however, the conduct of this study corresponds to EU Testing Method B7 and OECD 407.</p> <p>Deviations from current OECD 407:</p> <ul style="list-style-type: none"> <li>- reticulocytes and a measure of blood clotting time/potential not included.</li> <li>- epididymides, prostate and thymus were not weighed.</li> <li>- no histopathology was carried out</li> </ul> <p>Sprague-Dawley rats, 5/sex/group</p>	<p>Dimethomorph (ZTH 236 Z50); Batch No. L 5000; purity 99 %</p> <p>Diet: 0, 200, 1000 and 5000 ppm (males: 15.8, 80.9 and 305.9 mg/kg bw; females: 17.5, 81.1 and 283.3 mg/kg bw/day)</p> <p>28 days</p>	<p>5000 ppm: soft feces, swollen abdomen, hunched posture, piloerection, emaciation, lethargy and unsteady gait, reduced body weight, reduced food consumption, increases in platelet and neutrophil counts, increased blood urea nitrogen. Moribund animals were sacrificed.</p> <p>5000 ppm, females only: decreased plasma albumin, increased plasma globulin, increased urine volume, increased relative liver weight</p> <p>≥1000 ppm: increased blood urea nitrogen (females), not considered adverse</p> <p>NOAEL: 81 mg/kg bw</p>	<p>B.6.3.1.1</p> <p>Reliability 3</p>

CLH REPORT FOR DIMETHOMORPH

<p>Oral 28 day study</p> <p>Method not specified in the report; however, the conduct of this study corresponds to EU Testing Method B7 and OECD 407.</p> <p>Deviations from current OECD 407 guideline:</p> <ul style="list-style-type: none"> <li>- No haematological or clinical chemistry evaluation.</li> <li>- Adrenals, testes, epididymides, prostate, thymus were not weighed.</li> <li>- Histopathology was only carried out on the liver.</li> </ul> <p>Sprague-Dawley rats, 10/sex/group</p>	<p>Dimethomorph (CME 151); Batch No. DW 11/86; purity 96.6 ± 0.8 %</p> <p>Diet: 0, 2000, 3000 and 4000 ppm (males: 175, 300 and 400 mg/kg bw; females: 200, 300 and 400 mg/kg bw/day)</p> <p>28 days</p>	<p>4000 ppm: piloerection, swollen abdomen and emaciation, reduced body weight and body weight gain</p> <p>≥3000 ppm: increased relative liver weight, hepatocellular hypertrophy</p> <p>≥2000 ppm, females only: reduced body weight and body weight gain, increased relative liver weight, hepatocellular hypertrophy</p> <p>NOAEL: &lt;200 mg/kg bw</p>	<p>B.6.3.1.2</p> <p>Reliability 3</p>
---	--	--	---------------------------------------



CLH REPORT FOR DIMETHOMORPH

<p>Oral 28 day study</p> <p>Method not specified in the report; however, the conduct of this study corresponds to EU Testing Method B7 and OECD 407.</p> <p>Deviations from current OECD 407:</p> <ul style="list-style-type: none"> <li>- A measure of blood clotting time/potential was not included in the haematological evaluation.</li> <li>- Epididymides, prostate and thymus were not weighed</li> <li>-</li> <li>- Histopathology was only carried out on gross macroscopic lesions, adrenals, heart, intestines, kidneys, liver, pituitary, spleen and stomach.</li> </ul> <p>Fisher 344 rats, 7/sex/dose</p>	<p>Dimethomorph (SAG 151 - E isomer); Batch No. L4785; purity: E isomer 99.5 - 99.7 %, Z-isomer 0.3 - &lt; 0.5 %</p> <p>Gavage: 0, 10, 100 or 750 mg/kg bw/day.</p> <p>28 days</p>	<p>750 mg/kg bw: increased food intake (males), decreases in total blood haemoglobin, increased platelet counts, increased mean platelet volume (males), increases in protein, bilirubin, gamma glutamyl transpeptidase, cholesterol, calcium and triglyceride, increases in serum urea and creatinine, decreased splenic weight (males), increased relative liver weight, slight to moderate hepatic enlargement, dark discoloration of the liver, slight to moderate caecal enlargement and fluid caecal contents, patchy mid zonal hepatocellular cytoplasmic lipid vacuolation</p> <p>≥100 mg/kg bw, males only: increased relative liver weight, increased absolute and relative adrenal weight, slight to moderate hepatic enlargement, dark discoloration of the liver</p> <p>NOAEL: 10 mg/kg bw</p>	<p>B.6.3.1.3</p> <p>Reliability 2</p>
--	--	---	---------------------------------------

CLH REPORT FOR DIMETHOMORPH

<p>Oral 28 day study</p> <p>Method not specified in the report; however, the conduct of this study corresponds to EU Testing Method B7 and OECD 407.</p> <p>Deviations from current OECD 407 guideline: - A measure of blood clotting time/potential was not included in the haematological evaluation.</p> <p>- Epididymides, prostate and thymus were not weighed</p> <p>- Histopathology was only carried out on gross macroscopic lesions, adrenals, heart, brain, intestines, kidneys, liver, pituitary, testes, spleen and stomach.</p> <p>Fischer 344 rats, 7/sex/dose</p>	<p>SAG 151 Z isomer; Batch No. Th H296, ST90/106; purity 95.6 %</p> <p>Diet: 0, 10, 100 and 750 mg/kg bw/day.</p> <p>28 days</p>	<p>750 mg/kg bw/day: increases in serum bilirubin and protein, decreased albumin to globulin ration, slight caecal enlargement, fluid caecal contents (males)</p> <p>≥100 mg/kg bw: increased relative liver weight, patchy midzonal cytoplasmic lipid vacuolation</p> <p>NOAEL: 10 mg/kg bw</p>	<p>B.6.3.1.4</p> <p>Reliability 2</p>
<p>Oral, 28 day, immunotoxicity study.</p> <p>EPA Guideline 870.7800/ GLP</p> <p>Wistar rats, 8 male/dose</p>	<p>Dimetomorph, batch no AC9978-131, purity 97.5%</p> <p>0, 300, 800 and 2400 ppm in diet (23, 61 and 184 mg/kg bw/d)</p> <p>28 days</p>	<p>2400 ppm: reduced bw gain</p> <p>No immunotoxic effects observed.</p> <p>NOAEL ( immunotoxicity): 2400 ppm (184 mg/kg bw/d)</p> <p>NOAEL (general toxicity): 800 ppm (61 mg/kg bw/day)</p>	<p>B.6.8.2.15</p>

CLH REPORT FOR DIMETHOMORPH

<p>Oral 6 week study</p> <p>No guideline specified</p> <p>Only limited number of endpoints evaluated</p> <p>CD-1 mice, 10/sex/group</p>	<p>Dimethomorph (CME 151); Batch No. DW 11/86; purity not specified</p> <p>Diet: males: 0, 300/10000, 800 and 2000/5000 ppm; females: 0, 300/8000, 800 and 2000/4000 ppm</p> <p>6 weeks</p>	<p>All dose groups (females) and 300/10000 and 2000/5000 males showed significant increases in absolute and relative liver weight</p>	<p>B.6.3.1.5</p> <p>Reliability 3</p>
<p>Oral 14 day study</p> <p>No guideline specified</p> <p>Beagle dogs, 1/sex/dose</p>	<p>Dimethomorph CME 151; Batch No. DW 11/86, purity not specified</p> <p>Part A: Diet: 1000, 750, 900 and 1200 ppm for 7 days</p> <p>Part B: Diet: 1200 ppm for 14 days</p>	<p>≥900 ppm: body weight loss (males)</p> <p>1000 ppm: reduced food intake</p>	<p>B.6.3.1.6</p> <p>Reliability 3</p>
<p>Oral 90 day study plus 4 weeks recovery</p> <p>(US EPA Guideline 82-1 and OECD Guideline 408 claimed by the author); EU Testing Method B 26</p> <p>Deviations from current OECD guideline 408:</p> <ul style="list-style-type: none"> <li>- Epididymides and thymus were not weighed</li> </ul> <p>Sprague-Dawley rats, 20/sex/dose</p>	<p>Dimethomorph (CME 151); Batch No. T2/85; purity 98.7 ± 1.5 %</p> <p>Diet: 0, 40, 200 and 1000 ppm (males: 2.9, 14.2 and 73 mg/kg bw/day; females: 3.2, 15.8 and 82 mg/kg bw/day) for 13 weeks</p>	<p>1000 ppm males: decreased PCV, RBC, MCHC, total white blood cell count and lymphocyte counts, decreased urinary pH</p> <p>1000 ppm females: decreased heart weight, increased liver weight, decreased urinary pH</p> <p>≥200 ppm males only: decreased relative liver weight</p> <p>≥40 ppm females only: decreased kidney weight (not dose related)</p> <p>NOAEL: 200 ppm (16 mg/kg bw)</p>	<p>B.6.3.2.1</p> <p>Reliability 2</p>

CLH REPORT FOR DIMETHOMORPH

<p>Oral 90 day neurotoxicity study (US EPA Guideline 870.6200 and OECD Guideline 424; GLP) Wistar rats, 10/sex/dose</p>	<p>Dimetomorph (Batch No: AC9978-131; purity 98.3%) Diet: 0, 300, 800 and 2400 ppm (males: 0, 21.7, 58.7, and 177.9 mg/kg bw/day, females: 0, 25.7, 69.6, and 204.0 mg/kg bw/day) 90 days</p>	<p>2400 ppm: reduced food consumption, impaired body weight gain and food efficiency. No neurotoxic effect observed. NOAEL (general toxicity): 800 ppm NOAEL (neurotoxicity): 2400 ppm</p>	<p>B.6.7.1.2 Reliability 1</p>
<p>Oral 90 day study EU Testing Method B 27 (OECD Guideline 409 and US EPA Guideline 82-1 claimed by the author) Beagle dogs, 4/sex/group</p>	<p>Dimethomorph (CME 151; Batch No. DW 11/86; purity 96.6 ± 0.8 %) Diet: 0, 150, 450 and 1350 ppm (males: 5, 15 and 43 mg/kg bw/day; females: 6, 15 and 44 mg/kg bw/day) for 13 weeks</p>	<p>1350 ppm: lip-licking, occasional subdued behaviour, body tremors, increased serum alkaline phosphatase activity (males), increased absolute and relative thymus weight (males), increased relative liver weight (females), decreased absolute prostate weight (males), prostate fibrosis increased (males)  NOAEL: 450 ppm (15 mg/kg bw)</p>	<p>B.6.3.2.2 Reliability 1</p>
<p>Oral 52 week study Directive 96/54/EC B 30 (OECD Guideline 409 and US EPA Guideline 83-1 claimed by the author) Beagle dogs, 4/sex/dose</p>	<p>Dimethomorph (SAG 151; CME 151); Batch No. DW 11/86; purity 96.6 % Diet: 0, 150, 450 and 1350 ppm (males: 4.9, 14.7 and 44.6 mg/kg bw/day; females: 5.0, 15.7 and 47 mg/kg bw/day) for about 12 months</p>	<p>1350 ppm: increased serum alkaline phosphatase, increased absolute (males only) and relative liver weight, decreased absolute prostate weight together with interstitial fibrosis, ≥450 ppm males: increased relative testes weight ≥450 ppm females and ≥150 ppm males: reduced body weight (150 ppm females: increased body weight) ≥450 ppm females: increased hepatic lipid (in males only at 1350 ppm)  NOAEL: 150 ppm (4.9 mg/kg bw/day)</p>	<p>B.6.3.2.3 Reliability 1</p>
<p>Dermal 28 day study OECD 410, EPA 870.3200 Crl:WI(Han) rats, 10/sex/dose</p>	<p>AS 550 F (Dimethomorph). Lot/Batch #: COD-001244. Purity: 99.8% Dermal: 0, 100, 300, and 1000 mg/kg bw/day for 28 days</p>	<p>No treatment-related findings observed NOAEL: 1000 mg/kg bw</p>	<p>B.6.3.3.1 Reliability 1</p>

CLH REPORT FOR DIMETHOMORPH

<p>Long term toxicity and carcinogenicity study</p> <p>EU Testing Method B 30 (US EPA Guideline 83-5; Guidelines of OECD and Japanese MAFF, claimed by the author)</p> <p>Deviations from current OECD guideline:</p> <ul style="list-style-type: none"> <li>- Cholesterol not measured.</li> <li>- Thyroid and uterus were not weighed</li> </ul> <p>Sprague-Dawley rats, 20/sex/dose</p>	<p>Dimethomorph (SAG 151; CME 151); Batch No. DW 11/86; purity 96.6 %</p> <p>Diet: 0, 200, 750, and 2000 ppm (males: 9.4, 36.2 and 99.9 mg/kg bw/day; females: 11.9, 57.7 and 157.8 mg/kg bw/day) for 104 weeks.</p>	<p>2000 ppm: reduced body weight and body weight gain, decreased red blood cell count, decreased hemoglobin (females), decreased hematocrit (females), increased bone marrow cellularity (females), increased incidence of hepatocellular hypertrophy and/or increased amount of pigment in hepatocytes (females), increased incidences of dilated mesenteric blood vessels, arteritis and testicular interstitial cell proliferation (males)</p> <p>≥750 ppm: increased incidence of "ground-glass" foci of cellular alteration in the liver (males only at 2000 ppm)</p> <p>NOAEL: 200 ppm (9 mg/kg bw)</p>	<p>B.6.5.1.1 – study 1</p> <p>Reliability 2</p>
<p>Long term toxicity and carcinogenicity study</p> <p>EU Testing Method B 32 (US EPA Guideline 83-5; Guidelines of OECD and Japanese MAFF claimed by the author)</p> <p>Sprague-Dawley rats, 50/sex/dose</p>	<p>Dimethomorph (SAG 151; CME 151); Batch No. DW 11/86; purity 96.6 %</p> <p>Diet: 0, 200, 750, and 2000 ppm (males: 8.8, 33.9 and 94.6 mg/kg bw/day; females: 11.3, 46.3 and 132.5 mg/kg bw/day) for 104 weeks</p>	<p>2000 ppm: decreased food intake (females), increased incidence of dilated mesenteric blood vessels and arteritis in the abdominal vessels, predominantly in the pancreas (males), increase in "ground-glass" foci of cellular alteration in the liver, increased incidence in hepatocellular pigmentation and hypertrophy and increased severity of bone marrow cellularity (females)</p> <p>≥750 ppm: reduced body weight gain (males only at 2000 ppm)</p> <p>NOAEL: 200 ppm (9 mg/kg bw/day)</p>	<p>B.6.5.1.1 – study 2</p> <p>Reliability 1</p>

<p>Long term toxicity and carcinogenicity study</p> <p>EU Testing Method B.32 (US EPA Guideline 83-2; Guidelines of OECD and Japanese MAFF, claimed by the author)</p> <p>Charles River CD-1 mice, 50/sex/dose</p>	<p>Dimethomorph (SAG 151; CME 151); Batch No. DW 11/86; purity 96.6 %</p> <p>Diet nominal dose: 0, 10, 100, and 1000 mg/kg bw/day for 104 weeks</p>	<p>1000 mg/kg bw/day: reduced body weight (males) and body weight gain (females also at 100 mg/kg bw/day), increased alkaline phosphatase activity (males), increased aspartate aminotransferase activity (females), increased absolute and relative (females) liver weight</p> <p>NOAEL: 100 mg/kg bw/day</p>	<p>B.6.5.2</p> <p>Reliability 1</p>
--	---	--	-------------------------------------

#### 10.10.1 Short summary and overall relevance of the provided information on specific target organ toxicity – repeated exposure

The short-term toxicity of dimethomorph technical (E/Z racemat) was investigated in 2 dietary 28-day studies in Sprague-Dawley rats. In addition 2 dietary 28-day studies in Fisher 344 rats with E isomer and with Z isomer respectively were performed. Furthermore, dimethomorph technical (E/Z racemat) was investigated in a dietary 90-day study in Sprague-Dawley rats and Beagle dogs each, a 6 week dietary dose range finding study in CD-1 mice and a one-year dietary study in Beagle dogs.

Short-term (28-day) exposure of Sprague-Dawley rats to dimethomorph at dietary concentrations of 200, 1000, and 5000 ppm resulted in a NOAEL of 1000 ppm (equal to 80 mg/kg bw/day based on food consumption data). The highest dietary concentration tested (5000 ppm, equal to 280-300 mg/kg bw) induced increased morbidity and clinical signs of toxicity, reductions in body weight gains and in food consumption, increased liver to body weight ratios and increased blood urea nitrogen for both sexes. No microscopic examinations were performed.

In a second short-term (28-day) toxicity study in Sprague-Dawley rats, using dietary concentrations of 0, 2000, 3000, and 4000 ppm, the NOAEL was less than 2000 ppm based on decreased body weight gains for both sexes at all dietary concentrations, and dose-related increases in liver-to-body weight ratios and hepatocellular hypertrophy for both sexes at 3000 and 4000 ppm and for females at 2000 ppm. Based on food consumption data, the 2000 ppm concentration is equal to 150 mg/kg bw/day.

A 28-day exposure of Fisher 344 rats to E isomer dimethomorph at dietary concentrations of 10, 100 or 750 mg/kg bw/day resulted in a NOAEL of 10 mg/kg bw/day, based on a dark discoloration and enlargement of the liver, an increase in adjusted liver weight in the males and a mid-zonal hepatocellular cytoplasmic lipid vacuolation in male and female rats at 100 mg/kg bw/day.

A 28-day exposure of Fisher 344 rats to Z isomer dimethomorph at dietary concentrations of 0, 10, 100 and 750 mg/kg bw/day resulted in a NOAEL of 10 mg/kg bw/day, based on a slight mid-zonal lipid vacuolation at 100 or 750 mg/kg bw/day in both sexes.

In a 28-day oral immunotoxicity study in Wistar rats, no immunotoxic effects were noticed.

In the subchronic (90-day) feeding study in Sprague-Dawley rats with dimethomorph, the NOAEL of 200 ppm was based on increased liver weights for females at 1000 ppm. Based on food consumption data, the NOAEL of 200 ppm is equal to approximately 16 mg/kg bw/day.

In a 6 week dietary dose range finding study in mice a liver weight increase in males (5000 ppm and above) and females (4000 ppm and above) was observed.

In the subchronic (90-day) feeding study in Beagle dogs with dimethomorph, increases in alkaline phosphatase activity (both sexes), increases in relative liver weight in females, reductions in absolute and relative prostate weights, and an increased incidence of prostatic interstitial fibrosis in males were observed at 1350 ppm (highest concentration tested). The NOAEL for this study is 450 ppm, equal to approximately 15 mg/kg bw/day as based on food consumption data.

In a 90-day oral neurotoxicity study in Wistar rats, no neurotoxic effects were noticed.

In the 1-year feeding study in beagle dogs with dimethomorph technical, increased relative liver weights in females and increased relative testes weights in males occurred at 450 ppm. The NOAEL for this study is 150 ppm, equal to an approximate daily intake of 5 mg/kg bw/day as based on food consumption data.

A 28-dermal study was performed in Wistar rats, no systemic or local treatment-related effects were observed.

Long-term dietary toxicity studies were conducted in rats and mice. In both species liver effects were observed at high doses.

In the 2-year dietary toxicity study in Sprague-Dawley rats with dimethomorph, effects on the liver occurred in females at 750 ppm. The NOAEL for chronic toxicity in this study was 200 ppm, equal to approximately 9 mg/kg bw/day, based on food consumption data.

Similarly, for the 2-year carcinogenicity study in Sprague-Dawley rats, the NOAEL for chronic toxicity was 200 ppm (equal to approximately 9 mg/kg bw/day), as based on a decrease in overall body weight gain for females at 750 ppm. In both 104 week dietary toxicity studies in rats, there was an increased incidence of testicular tumors. However, the differences are not considered to be an oncogenic effect of dimethomorph (see Annex I for explanation).

In the 104-week carcinogenicity study with dimethomorph in CD-1 mice, the data support a systemic toxicity NOAEL of about 97 mg/kg bw/day, based on reduced body weight in males and reduced body weight gains in males and females at about 1000 mg/kg bw/day.

### 10.10.2 Comparison with the CLP criteria

*Not discussed in this dossier*

### 10.10.3 Conclusion on classification and labelling for STOT RE

*Not applicable.*

### 10.11 Aspiration hazard

*Not evaluated in this dossier*

## 11 EVALUATION OF ENVIRONMENTAL HAZARDS

### 11.1 Rapid degradability of organic substances

**Table 15: Summary of relevant information on rapid degradability**

Method	Results	Remarks	Reference
OECD Guideline 301B and D	No degradation observed in both studies	Not readily biodegradable (Ri 2)*	Turner (1988)
Ready			

Method	Results	Remarks	Reference
biodegradability			
OECD Guideline 111 Hydrolysis	stable at environmentally relevant conditions	(Ri 2)*	Ochsenbein (1989)
Photochemical degradation in water (US-EPA, OECD and BBA guidelines)	Half-life: 25 to 107 days	Degradation product detected at <10% of the applied dose (Ri 2)*	Knoch and Holman (1998), Panek et al. (2001), Van Dijk (1990)
OECD 302C Inherent biodegradability	maximum biodegradation of 27 % ThOD	Some evidence of inherent, primary biodegradability (Ri 2)*	Battersby (1993)
OECD Guideline 309 Surface water simulation	Half-life: 621 days	Result from one trial, other trials showed no significant degradation (Ri 2)*	Yeomans (2015)
Aquatic-sediment simulation study	Dissipation half-lives for the whole system: 2.6-58.4 days	Bound residues were observed in the range of 47 to 75% of the applied radiation (Ri 2)*	Ebert (2002), Knoch (1993), Knoch (1994b), Knoch (1994a), Maleri (2015)

\* Reliability according to Klimisch et al. (1997), since assessment is based on summaries in the DAR, Ri=1 is not given.

### 11.1.1 Ready biodegradability

The following study was available in the original registration dossier, the description below is based on the summary in the RAR.

Ready biodegradability of dimethomorph (94% purity, E/Z isomer ratio 45/55) has been assessed by Turner (1988) in a GLP study following the OECD 301B and 301D guidelines. Three tests were performed: a Closed Bottle Test; a Modified Sturm Test; and a Microbial Inhibition Test. For the Closed Bottle Test and Modified Sturm Test dimethomorph was emulsified with a non-biodegradable detergent (Dobane PT sulphonate) and this solution was added to the inoculum. For the closed bottle test a nominal concentration of dimethomorph of 3 mg/L was applied and for the Modified Sturm Test the concentration of dimethomorph in the inoculum was 20 mg/L. A reference control system used sodium benzoate to monitor the viability of the microbial population. A test system containing both the test and reference substances as the sole sources of carbon was used to evaluate the toxicity of the test compound. The amount of oxygen consumed was measured over time. In the Microbial Inhibition Test, the inhibition of the growth of a pure culture of *Pseudomonas fluorescens* by dimethomorph was studied. An acetone solution of dimethomorph was added to a mixture of *P. fluorescens* in a growth medium at a concentration of 50 mg a.s./L. A known microbial growth inhibitor, sodium pentachlorophenate, was also evaluated. The growth of the organism was measured over time. In the Closed Bottle Test, no oxygen was consumed in 28 days. Therefore it was concluded that dimethomorph was not degraded. There was no inhibition of microbial activity under the test conditions. In the Modified Sturm Test, there was no evolution of carbon dioxide from dimethomorph over 28 days. Therefore it was concluded that dimethomorph was not degraded. There was no inhibition of microbial activity under the test conditions. It was concluded that Dimethomorph is not “readily biodegradable” and does not inhibit the growth or activity of microorganisms. The results are considered reliable (Ri 2) and can be used for classification purposes.

### 11.1.2 BOD<sub>5</sub>/COD

*Not evaluated in this dossier*



### 11.1.3 Hydrolysis

In the original registration dossier one study on hydrolytic degradation of dimethomorph was available. No additional studies were added to the dossier for the renewal. The description below is based on the summary in the RAR.

Hydrolysis of radiolabelled and unlabelled dimethomorph (E/Z ratio 49.5/50.5) was tested in a GLP study by Ochsenbein (1989) in three buffer solutions of pH 4, 7 and 9 and at temperatures of 70 and 90 °C. The incubation was performed under a constant stream of nitrogen. The incubation flasks were connected to two traps to collect <sup>14</sup>CO<sub>2</sub> and organic volatiles. Samples were taken at start and after 2, 3, 4, 6, 8 and 10 weeks. At pH 4, 7 and 9 the substance was stable at both test temperatures. On the basis of this, it can be concluded that the substance is stable at pH 4 to 9 which are environmental relevant pHs. The results are considered reliable (Ri 2) and can be used for classification purposes.

### 11.1.4 Other convincing scientific evidence

*Not evaluated in this dossier*

#### 11.1.4.1 Field investigations and monitoring data (if relevant for C&L)

In the RAR some information from public literature on field investigations and monitoring data is reported. This included environmental monitoring of various pesticides in surface water, groundwater and/or sediment outside EU (e.g. China, South Africa). Most studies lacked standard methods or quantitative information. Frequency of detections were rare and concentrations not indicated or very low. It was reported that DT50 values of 10.3 to 31.5, 9.5 to 9.7 days or 14.6 days in soil were stated in several studies. It is unclear if these half-lives are for degradation or dissipation.

#### 11.1.4.2 Inherent and enhanced ready biodegradability tests

Inherent biodegradability has been assessed by Battersby (1993) in a GLP study following the OECD 302c guideline. The inherent biodegradability of dimethomorph was assessed by the modified MITI (II) test using fresh activated sludge as the inoculum, and 97.5 % pure dimethomorph (E/Z isomer 46/54). The MITI (II) inherent biodegradability test is similar to the modified MITI (I) ready biodegradability test except that the ratio of biomass to test substance is greater, therefore providing greater potential for biodegradation. A buffered salts medium was inoculated with activated sludge to get a mixed liquor suspended solids (MLSS) concentration of 100 mg/L. Duplicate respirometer flasks were set up containing 500 mL of inoculated medium amended with the test substance at 30 mg/L. Dimethomorph was added as an emulsion. A biodegradable reference with sodium benzoate was also performed. The flasks were equilibrated to the test temperature (20 °C) for one hour. Logging of oxygen uptake was then started and monitored every two hours over 28 days. At the end of this period, logging was stopped and each flask analysed for pH and N<sub>ox</sub>- (NO<sub>2</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>) concentrations. The following parameters were calculated: net oxygen uptake, biochemical oxygen demand (BOD), nitrogenous oxygen demand (NOD) and percent biodegradation. Biodegradation was expressed as a percentage of the theoretical maximum (ThOD). In one of the two replicate flasks the amount of oxygen uptake was 27 % ThOD. In the other replicate there was slight inhibition of oxygen uptake. Over 60 % of the biodegradable reference substance was degraded after 7 days of incubation, and the test was therefore considered valid. It was concluded that the maximum biodegradation of dimethomorph of 27 % ThOD in one of the replicate flasks was some evidence of “inherent, primary biodegradability”. The results are considered reliable (Ri 2) and can be used for classification purposes.

#### 11.1.4.3 Water, water-sediment and soil degradation data (including simulation studies)

In the dossier for the original registration two water/sediment simulation studies were available, these studies were reassessed for the RAR. In the renewal dossier, a new study on the aerobic mineralisation in surface water is also available, the descriptions below are based on the summaries in the RAR.

Knoch (1993, 1994b, 1994a) examined the degradation of radiolabelled dimethomorph (E/Z isomer ratio: 48.9/51.1) in two water-sediment systems. The labelling was performed on the chlorophenyl-moiety of the molecule, isomers were not determined separately. Two systems with water/sediment from different locations were used in the test. Dimethomorph was added directly to the surface of the water layer to give an initial concentration of 0.128 mg a.s./L. The samples were incubated at 20°C with a steady stream of CO<sub>2</sub>-free air passing over the water/sediment system and through traps to collect <sup>14</sup>CO<sub>2</sub> and organic volatiles. Total recovery was in the range of 93-106% for both systems. In both systems the dimethomorph moved quickly to the sediment and was fixed as bound residues. After 105 days of exposure, the complete mineralisation was 14- 22% of the applied radiation. No metabolites >10% were detected. 57 to 69% of the applied radiation was reported as non-extractable radiation in the sediment after 105 days. In a re-evaluation of the data by Maleri (2015), half-lives for the total systems determined according to the FOCUS degradation kinetics were 3.6 and 2.6 days, in this the fraction non-extractable from sediment was considered as degraded. These half-lives should be considered as dissipation half-lives since it is unclear if the fraction non-extractable from sediment has actually been degraded or is still the parent compound. The results are considered reliable (Ri 2) and can be used for classification purposes.

Ebert (2002) examined the degradation of two forms of radiolabelled dimethomorph (E/Z ratio: 42/58) in two water-sediment systems. The labelling was performed on the chlorophenyl- or the morpholine-moiety of the molecule. Two systems with water/sediment from different locations were used in the test. Dimethomorph was added directly to the surface of the water layer at a rate of 70 µg per test vessel and 210 µg per test vessel for isolation and determination of potential degradation products. The samples were incubated for 100 days (or 139 days for the degradation products) at 20°C. Total recovery was in the range of 93-106% for both systems. In both systems the dimethomorph moved quickly to the sediment and was fixed as bound residues in levels of 46 to 75% of the applied radiation. After 100 days of exposure, the complete mineralisation was 3.2-8.6% of the applied radiation. No metabolites >10% were detected. In a re-evaluation of the data by Maleri (2015), dissipation half-lives for the total systems were determined to be 15.4 and 58.4 days, in this the fraction non-extractable from sediment was considered as dissipated. The results are considered reliable (Ri 2) and can be used for classification purposes.

Yeomans (2015) examined the mineralization and degradation rates of dimethomorph (BAS 550F) in an aquatic system under dark conditions. The study was performed according to OECD guideline 309. The test was performed at two concentrations (10 and 100 µg a.s./L) using two differently <sup>14</sup>C labelled test items (morpholine and chlorophenyl labels), respectively. Sterile samples were tested for each label at the higher concentration. The test vessels were attached to a flow-through system for continuous aeration and incubated at a temperature of 20 ± 2°C in the dark. Samples for the experiment were taken at days 0, 3, 7, 14, 22, 36 and 59. The amount and nature of radioactivity in the water samples was determined by liquid scintillation counting (LSC) and radio-HPLC. Volatiles were trapped in 2 M sodium hydroxide and also analyzed by LSC. Parent substance identification was done by co chromatography with the corresponding reference items of the E and Z isomer on HPLC. From the obtained results it was concluded that dimethomorph was not significantly degraded in the test system. After 59 days, at least 86.2 and 91.6% of the total applied radioactivity was recovered as the unchanged active substance for the morpholine and chlorophenyl label, respectively. Several minor metabolites were observed during the study in small amounts of up to 4.0% applied radiation, In one single sample a component with 7.6% of the applied radiation was detected. During the test, no systematic change to the isomer ratio was observed for either the morpholine or chlorophenyl label. At the end of the study, the radioactivity in the water accounted for 92.4 to 96.5% of the applied radiation for the viable test vessels and for 95.3 to 96.0% of the applied radiation for the sterilized vessels. Radioactivity in the volatile traps did not exceed 2.1 or 0.7% of the applied radiation for the morpholine and chlorophenyl label respectively, indicating a low rate of mineralization. Overall, the compound was considered to be stable in the test systems. Degradation kinetics were not reported as no significant degradation was observed. In the RAR, the RMS has confirmed that for 5 out of 6 of the experiments with dimethomorph, no degradation can be demonstrated because the rate of degradation does not significantly differ from 0 and/or the t-test is > 0.05. The results are considered reliable (Ri 2) and can be used for classification purposes.

#### 11.1.4.4 Photochemical degradation

Three GLP studies on direct photochemical degradation in water were present in the original registration dossier, no additional studies were added to the dossier for the renewal. The descriptions below are based on the summary in the RAR.

Van Dijk (1990) examined photodegradation of radiolabelled dimethomorph in water in a GLP study according to a US-EPA guideline (US EPA Pesticide Assessment Guidelines, Subdivision N, Chemistry: Environmental Fate, Section 161-2: Photodegradation Studies in Water). Dimethomorph was dissolved in a buffer at pH 5 at a concentration of 6 µg/L. The solutions were continuously exposed at 25°C for 15 days. The light intensity was 90-92 KLux. Samples were taken at start and days 1, 2, 4, 7 and 15. A steady stream of air passed through the system, and passed over traps to collect <sup>14</sup>CO<sub>2</sub> and organic volatiles. A recovery of the radiolabels was achieved of 95 to 99.6% and the recovery in the control (dark) samples was 99.7 to 102.9%. After 15 days, only 0.5% of the applied radioactivity was retrieved the volatile traps and 70% of the parent compound was still present in the test solutions. Individual metabolites did not exceed 10%. The half-life of the parent was calculated to be 25-28 days. The results are considered reliable (Ri 2) and can be used for classification purposes.

Panek et al. (2001) examined photodegradation of radiolabelled dimethomorph (E/Z ratio: 42/58) in water in a GLP study according to a US-EPA guideline (US EPA Pesticide Assessment Guidelines, Subdivision N, Chemistry: Environmental Fate, Section 161-2: Photodegradation Studies in Water). Two different labels of dimethomorph were tested separately (<sup>14</sup>C(U)-chlorophenyl]-dimethomorph, > 98 % and 2,3,5,6-<sup>14</sup>C-morpholine]-dimethomorph, > 99 %). The substances were dissolved in a buffer at pH 5 at a concentration of 5.3 µg/L. The solutions were continuously exposed at 25°C for 21 days. The light intensity was 489-490 W/m<sup>2</sup> and wavelengths <290 nm were filtered out. Samples were taken at start and days 4, 8, 15 and 21. A steady stream of air passed through the system, and passed over traps to collect <sup>14</sup>CO<sub>2</sub> and organic volatiles. A recovery of the radiolabels was achieved of 100-103% and the recovery in the control (dark) samples was 99.5-106.2%. After 21 days, at most 2.2% of the applied radioactivity was retrieved from the volatile traps and at least 97.9% of the radioactivity was still present in the test solutions. 80.9-87.2% of the applied radioactivity was present as the parent. Individual metabolites did not exceed 10%. The half-life of the parent was calculated to be 107 and 86 days for the chlorophenyl and morpholine label, respectively. The results are considered reliable (Ri 2) and can be used for classification purposes.

Knoch and Holman (1998) examined photodegradation of unlabelled dimethomorph (E/Z isomer ratio: 44/56) in water in a GLP study according to a BBA and draft OECD guideline (Richtlinien für die Prüfung von Pflanzenschutzmitteln im Zulassungsverfahren Teil IV, 6-1; Biologische Bundesanstalt für Land- und Forstwirtschaft (BBA), D-38104 Braunschweig”, July 1990; Organisation for Economic Co-operation and Development (OECD), OECD Draft Test Guideline: “Phototransformation of Chemicals in Water”, December 1992). Dimethomorph with a purity of 97.6% was dissolved in a buffer at pH 7 at a concentration of 3.9 µg/L. The solutions were continuously exposed at 25°C for 72 hours. Wavelengths <290 nm were filtered out and the relative intensity of the light was 2.3 sun hours per instrument hour. Degradation was observed and the half live under test conditions was determined to be 303 hours. Normalised to normal sunlight the half live is estimated to be 29.2 hours. Metabolites were not reported. The results are considered reliable (Ri 2) and can be used for classification purposes.

### 11.2 Environmental transformation of metals or inorganic metals compounds

Not relevant

### 11.3 Environmental fate and other relevant information

In the RAR data on photochemical oxidation in the atmosphere is presented, considering the low vapour pressure ( $9.7 \times 10^{-7}$  and  $1.0 \times 10^{-6}$  Pa at 50°C) and low Henrys law constant of ( $5.4 \times 10^{-6}$  and  $2.5 \times 10^{-5}$  Pa m<sup>3</sup>/mol), this data is considered not relevant for classification purposes.

## 11.4 Bioaccumulation

**Table 16: Summary of relevant information on bioaccumulation**

Method	Results	Remarks	Reference
log Pow	E-isomer: 2.63 Z-isomer: 2.73	Measured (HPLC-method)	RAR
OECD 305  <i>Lepomis macrochirus</i>	1.4 and 2.0	based on the parent concentration and normalised to 5% fat (Ri=2)	B.9.2.3.1 (1999)

### 11.4.1 Estimated bioaccumulation

-None

### 11.4.2 Measured partition coefficient and bioaccumulation test data

Reference B.9.2.3.1 (1999) examined the bioconcentration of dimethomorph (BAS 550F) in *Lepomis macrochirus*. The fish were exposed for 28 days to radiolabelled dimethomorph at nominal concentrations of 0.020 mg/L and 0.200 mg/L (corresponding to mean measured concentrations of 0.021 mg/L and 0.210 mg/l). Steady state was achieved by day 14 of the uptake period. The exposure was followed by a 14 day depuration period. In each treatment 120 fish were exposed and fish were sampled for analysis at days 7, 14, 21 and 28 of the uptake period and days 1 and 3 of the depuration period. The lipid content of the fish was 8.41 % and 8.07% for whole fish in the low and high treatment groups respectively. The kinetic BCFs calculated from total radioactive residues are 16 and 22 for the low and high exposure respectively. Normalised to 5% fat, the BCF values are 10 and 13 respectively. Kinetic BCFs based on the concentration of dimethomorph are 2.4 and 3.2 for the low and high exposure respectively. Normalised to 5% fat, the BCF values are 1.4 and 2.0 respectively. The endpoints are considered reliable (Ri=2) and can be used for classification purposes.

The highest BCF reported for dimethomorph is 2.0 (normalised to 5% fat) and the measured LogK<sub>ow</sub> values range from 2.63-2.73. Therefore it is considered to have a low potential for bioaccumulation.

## 11.5 Acute aquatic hazard

**Table 17: Summary of relevant information on acute aquatic toxicity**

The dimethomorph used in the aquatic toxicity studies was of technical grade, the E/Z ratio was not specified in the RAR.

Method	Species	Test material	Results	Remarks	Reference
<b>Fish</b>					
Static toxicity study according to OECD guideline 203	<i>Oncorhynchus mykiss</i>	Dimethomorph Purity: 94.8%	LC50 = 6.1 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.1 (1986), B.9.2.1.4 (2010)
Flow through toxicity study according to OECD guideline 203	<i>Oncorhynchus mykiss</i>	Dimethomorph Purity: 98%	LC50 = 6.8 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.7 (2001)
Flow through toxicity study	<i>Cyprinodon variegatus</i>	Dimethomorph Purity: 98.0%	LC50 = 11.3 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.8 (1997)

CLH REPORT FOR DIMETHOMORPH

Static GLP toxicity study according to OECD guideline 203	<i>Cyprinus carpio</i>	Dimethomorph Purity: 94.8%	LC50 = 16.6 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.3 (1986), B.9.2.1.2 (2010)
Static GLP toxicity study according to OECD guideline 203	<i>Lepomis macrochirus</i>	Dimethomorph Purity: 96.6%	LC50 >13.7 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.5 (1988), B.9.2.1.6 (2010)
Flow through toxicity study according to OECD guideline 203	<i>Lepomis macrochirus</i>	Dimethomorph Purity: 98%	LC50 >9.5 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.9 (2001)
Flow through toxicity study	<i>Pimephales promelas</i>	Dimethomorph Purity: 99.7%	LC50 >8.4 mg a.s./L (mortality; mean measured)	Study is considered reliable (Ri=2)*	B.9.2.1.10 (2014)
<b>Invertebrates</b>					
Static GLP study according to OECD guideline 202	<i>Daphnia magna</i>	Dimethomorph Purity: 94.8%	EC50 = 20.1 a.s. mg/L (mobility; mean measured)	Study is considered reliable (Ri=2)*	Ellgehausen (1986a), Habekost (2010)
Static GLP study according to OECD guideline 202	<i>Daphnia magna</i>	Dimethomorph Purity: 98.3%	EC50 >10.6 a.s. mg/L (mobility; mean measured)	Study is considered reliable (Ri=2)*	Mitchell (2001)
Flow through toxicity study according to EPA guideline 72-3	<i>Americamysis bahia</i>	Dimethomorph Purity: 98.0%	EC50 = 7.92 mg a.s./L (mobility; mean measured)	Study is considered reliable (Ri=2)*	Mitchell (1997a)
Flow through toxicity study according to EPA guideline 72-3	<i>Crassostrea virginica</i>	Dimethomorph Purity: 98.0%	EC50 = 4.42 mg a.s./L (shell growth; mean measured)	Endpoint for shell growth. Study is considered reliable (Ri=2)*	Mitchell (1997b)
<b>Algae/Aquatic Plants</b>					
Static GLP study according to OECD guideline 201	<i>Scenedesmus subspicatus</i>	Dimethomorph Purity: 94.8%	E <sub>0</sub> C50 = 29.2 mg a.s./L (biomass; nominal) E <sub>r</sub> C50 = not available (growth rate)	Study is considered unreliable for classification purposes (Ri=3)*	Ellgehausen (1986b)
Static GLP study according to OECD guideline 201	<i>Pseudokirchneriella subcapitata</i>	Dimethomorph Purity: 99.57%	EC50 = 82.2 mg a.s./L (growth rate; nominal) EC50 = 41.1 mg a.s./L (biomass; nominal)	Study is considered unreliable for classification purposes (Ri=3)*	Jatzek (2001)
Static study	<i>Lenma minor</i>	Dimethomorph	EC50 = not reported	Study is considered unreliable (Ri=3)*	Megateli and al. (2009), Megateli and al. (2013)

\* Reliability according to Klimisch et al. (1997), since assessment is based on summaries in the DAR, Ri=1 is not given.

### 11.5.1 Acute (short-term) toxicity to fish

One additional study on acute toxicity to fish was included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were reevaluated for the renewal. The descriptions below are based on the summaries in the RAR.

Reference B.9.2.1.1 (1986) has tested *Oncorhynchus mykiss* at five exposure concentrations for 96 hours in a batch exposure according to OECD guideline 203. The dimethomorph was technical grade with a purity of 94.87% and the five nominal test concentrations were 1, 2, 5, 7 and 10 mg/L. Mortality was observed in concentrations of 2 mg/L and higher. In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 2.4, 4.0, 8.0, 10.3 and 14.1 mg/L. Sub-lethal effects were observed at the lowest exposure concentration and higher. In a reassessment (B.9.2.1.4, 2010), the LC50 based on mean measured concentrations was determined to be 6.1 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.3 (1986) has tested *Cyprinus carpio* at four exposure concentrations for 96 hours in a batch exposure according to OECD guideline 203 at 21-23°C. The dimethomorph was technical grade with a purity of 94.87% and the four nominal test concentrations were 12, 14, 16 and 20 mg/L. Mortality was observed in concentrations of 14 mg/L and higher. In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 115.6 – 129.2% of nominal. Sub-lethal effects were observed at the lowest exposure concentrations and higher. In a reassessment (B.9.2.1.2, 2010), the LC50 based on mean measured concentrations was determined to be 18.1 mg a.s./L but in the RAR, the RMS has recalculated the LC50 to 16.6 mg/L. The latter endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.5 (1988) has tested *Lepomis macrochirus* at five exposure concentrations for 96 hours in a batch exposure according to OECD guideline 203 at 20-21°C. The dimethomorph was technical grade with a purity of 96.67% and the five nominal test concentrations were 2.5, 4.5, 8.0, 14 and 25 mg/L. Mortality was observed in concentrations of 8.0 mg/L and higher, the mortality in the highest test concentration was 50%. In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 1.9, 3.3, 5.3, 8.1 and 13.7 mg/L. Sub-lethal effects were observed at the 4.5 mg/L nominal exposure concentration and higher. In a reassessment (B.9.2.1.6, 2010), the LC50 based on mean measured concentrations was determined to be >13.7 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.7 (2001) has tested *Oncorhynchus mykiss* at five exposure concentrations for 96 hours in a flow-through exposure according to OECD guideline 203. The dimethomorph was technical grade with a purity of 98% and the five nominal test concentrations were 2.6, 4.4, 7.2, 12 and 20 mg/L. Mortality was observed in concentrations of 4.4 mg/L and higher. In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 1.75, 3.13, 4.71, 7.84 and 12.0 mg/L. Sub-lethal effects were observed at the nominal exposure concentration of 4.4 mg/L and higher. The LC50 was determined to be 6.79 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.8 (1997) has tested *Cyprinodon variegatus* at five exposure concentrations for 96 hours in a flow-through exposure according to OECD guideline 203. The dimethomorph was technical grade with a purity of 98.0% and the five nominal test concentrations were 2.6, 4.4, 7.2, 12 and 20 mg/L. Mortality was observed in concentrations of 4.4 mg/L and higher. In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 2.00, 3.29, 5.27, 8.82 and 14.6 mg/L. Sub-lethal effects were observed at the nominal exposure concentration of 4.4 mg/L and higher. The LC50 was determined to be 11.3 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.9 (2001) has tested *Lepomis macrochirus* at five exposure concentrations for 96 hours in a flow-through exposure according to OECD guideline 203. The dimethomorph was technical grade with a purity of 98% and the five nominal test concentrations were 2.6, 4.4, 7.2, 12 and 20 mg/L. Mortality was observed in the highest test concentration (5%). In the control and solvent control no mortalities were observed. Test concentrations were verified by analysis and were 1.62, 2.78, 4.19, 6.75 and 9.53 mg/L. Sub-

lethal effects were not observed. The LC50 was determined to be >9.53 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Reference B.9.2.1.10 (2014) has tested *Pimephales promelas* at five exposure concentrations for 96 hours in a flow-through exposure according to OECD guideline 203. The dimethomorph had a purity of 99.7% and the five nominal test concentrations were 0.63, 1.3, 2.5, 5.0 and 10 mg/L. Mortalities were not observed in any test concentration, control or solvent control. Test concentrations were verified by analysis and were 0.60, 1.3, 2.5, 4.9 and 8.4 mg/L. Sub-lethal effects were not observed. The LC50 was determined to be >8.4 mg/L. The endpoint is considered reliable and can be used for classification purposes.

### **11.5.2 Acute (short-term) toxicity to aquatic invertebrates**

No additional studies on acute toxicity to aquatic invertebrates were included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were reevaluated for the renewal. The descriptions below are based on the summaries in the RAR.

Acute toxicity of dimethomorph to *Daphnia magna* was tested in a static test by Ellgehausen (1986a) according to OECD guideline 202. The test compound was technical grade with a purity of 94.8% and eight concentrations were tested of 7.81, 15.6, 31.3, 62.5, 125, 250, 500, and 1000 mg/L. Test concentrations were verified by analysis and were 3.6, 11.8, 19.0, 27.3, 35.9, 42.7, 49.9 and 58.0 mg/L. After 48 hours immobilisation was observed in the solutions with nominal concentration of 15.6 mg a.s. and higher. In a reassessment (Habekost, 2010), the EC50 based on mean measured concentrations was determined to be 20.1 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Acute toxicity of dimethomorph to *Daphnia magna* was tested in a static test by Mitchell et al. (2001) according to OECD guideline 202. The test compound of technical grade had a purity of 98.3% and five concentrations were tested of 1.3, 2.5, 5.0, 10 and 20 mg/L. Test concentrations were verified by analysis and ranged from 53 to 97% of nominal. were 1.26, 2.38, 4.66, 6.98 and 10.6 mg/L. After 48 hours immobilisation (5%) was only observed in the highest test concentration. The EC50 based on mean measured concentrations was >10.6 mg/L. The endpoint is considered reliable and can be used for classification purposes.

Acute toxicity of dimethomorph to *Americamysis bahia* was tested in a flow-through test by Mitchell et al. (1997a) according to EPA guideline 72-3. The test compound of technical grade had a purity of 98.0% and five concentrations were tested of 1.3, 2.2, 3.5, 6.0 and 10 mg/L. Test concentrations were verified by analysis and were 1.21, 2.16, 3.45, 5.89 and 9.75 mg/L. After 48 hours immobilisation was observed in the nominal test concentration of 2.2 mg a.s./L and higher. The EC50 based on mean measured concentrations was 7.92 a.s. mg/L. The endpoint is considered reliable and can be used for classification purposes.

Acute toxicity of dimethomorph to *Crassostrea virginica* was tested in a flow-through test by Mitchell et al. (1997b) according to EPA guideline 72-3(b). The test compound of technical grade had a purity of 98.0% and five concentrations were tested of 1.3, 2.2, 3.5, 6.0 and 10 mg/L. Test concentrations were verified by analysis and were 1.33, 2.24, 3.63, 6.15 and 10.1 mg/L. After 96 hours mortality of one oyster was observed in the highest test concentration. Shell growth was statistically significantly inhibited at the three highest test item concentrations compared to the control. The EC50 based on mean measured concentrations for shell growth inhibition was 4.42 a.s. mg/L. The endpoint is considered reliable and can be used for classification purposes.

### **11.5.3 Acute (short-term) toxicity to algae and other aquatic plants**

No additional studies on acute toxicity to algae and other aquatic plants were included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were re-evaluated for the renewal. From a literature search four publications were included in the RAR and assessed on their reliability. The descriptions below are based on the summaries in the RAR.

Aquatic toxicity to algae was tested by Ellgehausen (1986b) on *Scenedesmus subspicatus* in a 72 hours exposure according to OECD test guideline 201. The dimethomorph of technical grade with a purity of 94.8% was tested in nominal concentrations of 10, 20, 40, 80 and 160 mg/L in three replicates. Test

concentrations were verified by analysis and were within 80 to 123% of nominal for the concentration of 20 mg/L but 23.5 to 45.1% for the higher test concentrations. The latter two solutions were turbid because the concentrations exceed the water solubility and only clear solutions were analysed. The actual mean measured concentrations are not available. Over the exposure period inhibition of growth was observed from the lowest test concentration and the EC50 based on nominal concentrations was determined to be 29.2 mg/L. In the RAR for the assessment of the renewal dossier it was requested that the endpoint had to be based on mean measured concentration. In the absence of the raw data, this was not possible. The test is considered unreliable and will not be used for classification purposes.

Aquatic toxicity to algae was tested by Jatzek (2001) on *Pseudokichneriella subcapitata* in a 72 hours exposure according to OECD test guideline 201. The dimethomorph of technical grade with a purity of 98.3% was tested in nominal concentrations of 6.25, 12.5, 25, 50 and 100 mg/L in three replicates. Test concentrations were verified by analysis and the concentrations ranged from 75.8 to 82.5% of the nominal concentrations at 0 hours, to 77.6 to 94.0% of the nominal concentrations at 72 hours. It is also reported that the two highest test concentrations which are at or above the water solubility, were turbid. It is unclear if analysis of the two higher test solution is performed on clear solutions or that undissolved substance is included in the analysis. The actual mean measured concentrations are not presented in the RAR. Over the exposure period inhibition of growth was observed from the lowest test concentration and the EC50 based on nominal concentrations was determined to be 41.1 mg/L for biomass and 82.2 for growth rate. In the RAR comments of the RMS are not provided on this study. However for the study of Ellgehausen (1986b) on *Scenedesmus subspicatus* it was requested that the endpoint had to be based on mean measured concentrations. It is presumed that this also applies to the endpoints of the study of Jatzek (2001), especially since reported measured endpoints are lower than 80% and turbidity is observed in the two higher test solutions. In the absence of an endpoint based on measured data, the test is considered unreliable and will not be used for classification purposes.

Megateli et al. (2009, 2013) studied the effect of dimethomorph, as single substance or in combination with copper sulfat, on the growth of the aquatic plant *Lemna gibba* in a 168 h static toxicity test. The dimethomorph of technical grade with a purity of 97% was tested in nominal concentrations of 0.040, 0.200, 0.400, 0.800 and 1.000 mg/L in three replicates. The tests with copper sulfate are not relevant for classification and therefore not further discussed in this report. Test concentrations were verified by analysis and concentrations lower than 80% of the nominal concentrations were reported (10-40% of nominal). Raw data are not available and endpoint are based on nominal concentrations. EC50 values are not reported in the RAR and it was stated that the study could not be used for the renewal application especially since endpoints are based on nominal concentrations while measured concentrations are well below the nominal concentrations. For the same reasons the study is considered unreliable and will not be used for classification purposes.

## 11.6 Long-term aquatic hazard

**Table 18: Summary of relevant information on chronic aquatic toxicity**

The dimethomorph used in the aquatic toxicity studies was of technical grade, the E/Z ratio was not specified in the RAR.

Method	Species	Test material	Results	Remarks	Reference
<b>Fish</b>					
GLP, flow through, Early Life Stages test according to OECD guideline 210	<i>Oncorhynchus mykiss</i>	Dimethomorph Purity: 97.6%	NOEC (96d) = 0.056 mg a.s./L (weight; mean measured)  EC <sub>10</sub> = 0.116 mg a.s./L (weight; mean measured) EC <sub>10</sub> > 0.897 mg a.s./L (length; mean measure)	The study is considered reliable (Ri = 2)*  Calculation of EC <sub>10</sub> values	B.9.2.2.2 (2015), B.9.2.2.1 (1997)  B.9.2.2.2 (2015)



CLH REPORT FOR DIMETHOMORPH

GLP Static Early Life Stages test according to EPA guideline 850.1400	<i>Cyprinodon variegatus</i>	Dimethomorph Purity: 97.5%	NOEC (40 d) = 0.136 mg a.s./L (hatching, weight and length; mean measured)  EC10 = 0.150 mg a.s./L (weight; mean measured) EC10 = 0.759 mg a.s./L (length; mean measured)	The study is considered reliable (Ri = 2)*  Calculation of EC <sub>10</sub> values	B.9.2.2.2 (2015), B.9.2.2.6 (2010)  B.9.2.2.2 (2015)
GLP Static Early Life Stages test according to OECD guideline 210	<i>Pimephales promelas</i>	Dimethomorph Purity: 98.3%	NOEC (34 d) = 0.107 mg a.s./L (hatching mean measured)  EC10 > 0.92 mg a.s./L (weight; mean measured)	The NOEC for hatching is considered reliable (Ri = 2)*  The EC10 for weight is considered unreliable (Ri=3)*	B.9.2.2.5 (2002)  B.9.2.2.2 (2015)
GLP 21 day Short term reproduction assay flow-through study	<i>Pimephales promelas</i>	Dimethomorph Purity: 99.7%	NOEC (21-d) ≥ 0.488 mg a.s./L (survival, weight, length, behaviour; mean measured)	The study is considered reliable (Ri = 2)*	B.9.2.4.1 (2014)
<b>Invertebrates</b>					
GLP 22 day renewal study according to OECD guideline	<i>Daphnia magna</i>	Dimethomorph Purity: 95.6%	NOEC = 0.10 mg a.s./L (survival and reproduction; nominal)  EC10 = 0.15 mg a.s./L (reproduction; nominal)	The study is considered reliable (Ri = 2)*	Anonymous (1993), Brausch (2015), Memmert and Knoch (1993)
GLP 21 day flow-through study according to OECD guideline	<i>Daphnia magna</i>	Dimethomorph Purity: 97.6%	NOEC = 0.22 mg a.s./L (length; mean measured)  EC10 = 0.42 mg a.s./L (reproduction; mean measured) EC10 > 2.0 mg a.s./L (length; mean measured) EC10 = 1.343 mg a.s./L (weight; mean measured)	The study is considered reliable (Ri = 2)*	Brausch (2015), Murrell (1997)
GLP 28 day flow-through study according to OECD guideline	<i>Americamysis bahia</i>	Dimethomorph Purity: 97.5%	NOEC = 0.24 mg a.s./L (reproduction; mean measured)  EC10 = 0.24 mg a.s./L (reproduction; nominal)	The study is considered reliable (Ri = 2)*	Hicks (2010)
24 day static study according to OECD guideline	<i>Chironomus riparius</i>	<sup>14</sup> C-dimethomorph Purity: 99.3%	NOEC = 4.11 mg a.s./L (emergence and weight; mean measured, initial)  EC10 = 3.02 mg a.s./L (weight; nominal) EC10 > 15.6 mg a.s./L (emergence; nominal)	Water spiked water-sediment system, the study is considered reliable (Ri = 2)*	Brausch (2015), England and al. (1997)
<b>Algae/Aquatic Plants</b>					
Static GLP study according to OECD guideline 201	<i>Scenedesmus subspicatus</i>	Dimethomorph Purity: 94.8%	NOEC/EC10 = not available	The study is considered unreliable (Ri = 3)*	Ellgehausen (1986b)

Static GLP study according to OECD guideline 201	<i>Pseudokirchneriella subcapitata</i>	Dimethomorph Purity: 98.3%	EC10 = 27.3 mg a.s./L (growth rate; nominal) EC10 = 9.23 mg a.s./L (biomass; nominal)	Study is considered unreliable for classification purposes (Ri=3)*	Jatzek (2001)
Static study	<i>Lenma minor</i>	Dimethomorph	NOEC/EC10 = not reported	Study is considered unreliable (Ri=3)*	Megateli and al. (2009), Megateli and al. (2013)

\* Reliability according to Klimisch et al. (1997), since assessment is based on summaries in the DAR, Ri=1 is not given.

### 11.6.1 Chronic toxicity to fish

Three new studies on chronic toxicity to fish were included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were re-evaluated for the renewal. The descriptions below are based on the summaries in the RAR.

Chronic toxicity in fish was determined in *Onchorhynchus mykiss* according to the OECD test guideline 210 by Reference B.9.2.2.1. (1997). The test compound of technical grade with a purity of 97.6% was tested in five nominal concentrations of 0.033, 0.065, 0.13, 0.25, 0.50, and 1.0 mg/L, and mean measured concentrations were 0.0341, 0.0562, 0.120, 0.240, 0.449 and 0.897 mg/L (89.7 to 103% of nominal). The experiment started with fertilised egg and exposure lasted for 96 days. At completion of hatching, 100% survival was observed for the control and 98-100% for the treatments. This was not significantly different. At test termination, survival was not significantly affected in any of the treatments. For growth, mean weight was significantly affected at mean measured concentration of 0.12 mg/L. Therefore, the NOEC (for weight), based on mean measured concentrations, was determined to be 0.056 mg a.s./L. In a reassessment (B.9.2.2.2, 2015), the LC10 for weight, based on mean measured concentrations, was determined to be 0.116 mg a.s./L. The study is considered reliable and the NOEC and EC10 can be used for classification purposes.

Chronic toxicity in fish was determined in *Pimephales promelas* according to the OECD test guideline 210 by Reference B.9.2.2.5 (2002). The test compound of technical grade with a purity of 98.3% was tested in five nominal concentrations of 0.01, 0.033, 0.1, 0.33 and 1 mg a.s./L, and mean measured concentrations were 0.0082, 0.0310, 0.107, 0.347 and 0.92 mg/L (82 to 107% of nominal). The experiment started with fertilised egg and exposure lasted for 34 days. At completion of hatching, 85% survival was observed for the control and 71-87% for the treatments. Only in the nominal concentration of 0.33 mg a.s./L, embryo survival was significantly lower (71%) than the control. Embryo survival in the highest test concentration of 1 mg/L (nominal) was also lower (75%) than the control but this effect was not significant. At test termination, survival of the hatched larvae was not significantly affected in any of the treatments. Growth and mean weight was significantly increased at the nominal concentration of 0.33 mg a.s./L and not in the highest concentration of 1 mg/L (highest). This observation was considered an artifact because of the lower number of in the replicates of this group due to the increased mortality rate until hatch. And an effect of the test substance was considered not plausible since body weight and length in the highest test concentration group were comparable to the control. Because in the highest test concentration no significant effects were observed there was no clear dose-response relation. Altogether only the reduction in embryo survival is considered an effect of the test item since this is confirmed by the (not-significant) reduction in the highest test concentration. This results in a NOEC for hatching based on mean measured concentrations of 0.107 mg a.s./L. An EC10 of >0.92 mg a.s./L is also calculated for weight and length but as the observation is probably due to an artifact and there is no dose-response relation this value is considered unreliable. The NOEC for hatching is considered sufficiently reliable for classification purposes (Ri=2).

Chronic toxicity in fish was determined in *Cyprinodon variegatus* according to the EPA test guideline 850-1400 by Reference B.9.2.2.6 (2010). The test compound of technical grade with a purity of 97.5% was tested in five nominal concentrations of 0.065, 0.13, 0.25, 0.50, 1.0 and 2.0 mg a.s./L, and mean measured concentrations were 0.0630, 0.136, 0.266, 0.536, 1.02 and 2.01 mg/L (94 to 112% of nominal). The experiment started with fertilised egg and exposure lasted for 40 days. At completion of hatching, 84% survival was observed for the control and 84-91% for the treatments. The time to hatch was significantly

longer for the nominal test concentration of 0.25 mg a.s./L and higher. At test termination, survival of the hatched larvae was not significantly affected in any of the treatments. Length and mean weight was significantly affected at the nominal concentration of 0.25 mg a.s./L and higher. The NOEC (for hatching, weight and length), based on mean measured concentrations, was determined to be 0.136 mg a.s./L. EC10 values were determined in a re-assessment (B.9.2.2.2, 2015). The EC10 for mean length is 0.759 mg a.s./L and the EC10 for mean weight is 0.150 mg a.s./L. The study is considered reliable and the NOEC and EC10 can be used for classification purposes.

Potential endocrine activity of dimethomorph to *Pimephales promelas* was determined by Reference B.9.2.4.1 (2014). The test compound of technical grade with a purity of 99.7% was tested in five nominal concentrations of 0.047, 0.15 and 0.48 mg a.s./L, and mean measured concentrations were 0.046, 0.143 and 0.488 mg/L (at test start 80.7 to 92.7% of nominal and at termination 110.7-115.2% of nominal). The fish were approx. 5 months old at test initiation. Exposure lasted for 21 days. The evaluated endpoints were survival, fecundity, fertilization success, nuptial tubercle score, fish weight and length, blood plasma vitellogenin (VTG) concentration, histological examination of gonadal tissues as well as behavior and appearance, including secondary sexual characteristics. The biological results are based on mean measured concentrations. Percent survival of males, females and the survival rates based on combined data sets in the control were 100%. No statistically significant effects on fish wet weight and total length compared to the control were observed at any test concentration. The vitellogenin concentration in the blood plasma of male fish in comparison to the control group was statistically significantly increased in the 0.15 mg a.s./L test group only by using the Wilcoxon test. However, the median vitellogenin values of the males were not dose-dependently altered. The percentage of fertilized eggs in the control and all test item concentrations was 99.8%. The mean number of eggs per female per day was not statistically significantly different in comparison to the control group. Tubercles were not observed in females; therefore, they were not scored. For males, no statistically significant differences in mean tubercle scores compared to the control were determined in any test concentration. Furthermore, no notable abnormalities were observed with regards to behavior, coloration/banding, changes in ovipositor appearance or size of dorsal nape pad. Dimethomorph did not demonstrate any (anti-)estrogenic or (anti-)androgenic potential when tested at the maximum tolerated dose (MTC) in fathead minnow. The overall NOEC (21 d) for dimethomorph was determined to be  $\geq 0.488$  mg a.s./L based on mean measured concentrations. The study is considered reliable and the NOEC can be used for classification purposes.

### 11.6.2 Chronic toxicity to aquatic invertebrates

In addition to the existing studies, one new study on chronic toxicity to invertebrates was included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were re-evaluated for the renewal. The descriptions below are based on the summaries in the RAR.

A GLP 22 day static renewal test on *Daphnia magna* was performed by Memmert and Knoch (1993) with technical grade dimethomorph (95.6%). The test was performed according to OECD guideline 202. The nominal exposure concentrations were 0.10, 0.31, 0.96, 3.1 and 9.6 mg a.s./L. Ten animals were tested at each exposure concentration. The test concentrations were analytically determined (Anonymous, 1993) and the mean measured concentrations were 86.9 to 105.5% of the nominal concentrations. The results are based on the nominal concentrations. Survival of the first generation was 100% for the solvent control and the lowest test concentration. 10% mortality was observed in the control. Significant mortality of the parents was only observed in the highest test concentration. The number of offspring per surviving adult was significantly affected at exposure concentration 0.31 mg a.s./L and higher. The NOEC for survival of the parents, based on nominal concentrations, was determined to be 0.1 mg/L. The NOEC for number of offspring is 0.1 mg a.s./L. In a reassessment (Brausch, 2015), the EC10 for reproduction, based on nominal concentrations, was determined to be 0.152 mg a.s./L. The endpoints are considered reliable and can be used for classification purposes.

A GLP 21 day flow-through study on *Daphnia magna* was performed by Murrell et al. (1997) with technical grade dimethomorph (97.6%). The test was performed according to EPA guideline 72-4. The nominal exposure concentrations were 0.13, 0.25, 0.50, 1.0 and 2.0 mg a.s./L. Four replicates were performed with ten adults each. The test concentrations were analytically determined and the mean measured

concentrations were 85 to 95% of the nominal concentrations. The results are based on mean measured concentrations. Survival of the first generation was 95% in the control. In the exposures, no significant effects were observed on adult survival. The number of offspring per surviving adult was significantly affected at the two highest exposure concentrations. Mean length of surviving adults was significantly affected at concentrations 0.50 mg a.s./L and higher. The overall NOEC, based on mean length, was determined to be 0.22 mg a.s./L (mean measured). In a reassessment (Brausch, 2015), EC10s for off spring per adult, length and body weight were determined to be 0.421, >2.0 and 1.343 mg a.s./L respectively. The endpoints are considered reliable and the EC10 for number of offspring can be used for classification purposes.

A GLP 28 day flow-through study on *Americamysis bahia* was performed by Hicks (2010) with technical grade dimethomorph (97.5%). The test was performed according to EPA guideline 72-4. The nominal exposure concentrations were 0.065, 0.13, 0.25, 0.50 and 1.0 mg a.s./L. Three replicates were performed with 15 adults each. The test concentrations were analytically determined and the measured concentrations ranged from 78 to 122% of the nominal concentrations. The results are based on mean measured concentrations. Survival of the first generation (F0) was 93% in the control. In the exposures, no significant effects were observed on F0 and F1 survival. The number of offspring per female was significantly affected at the two highest exposure concentrations. For mean length of F0, some significant effects were observed but these did not indicate a dose-effect relation. The overall NOEC, based on reproduction, was determined to be 0.24 mg a.s./L (mean measured) and the EC10 was 0.238 mg a.s./L. The endpoints are considered reliable and can be used for classification purposes.

A GLP 24 day water spiked water-sediment study on *Chironomus riparius* was performed by England et al. (1997) with radiolabelled dimethomorph (99.3%). The test was performed according to EPA guideline 72-4. The nominal exposure concentrations were 1.13, 2.25, 4.50, 9.00 and 18.0 mg a.s./L (corresponding to initially mean measured concentrations: 1.08, 2.15, 4.11, 7.97 and 15.6 mg a.s./L). 14 replicates were performed for each treatment of which eight with test organisms and six for analytical sampling. The test concentrations were analytically determined and the initial measured concentrations ranged from 87 to 96% of the nominal concentrations. The concentrations in the water phase declined to one third of the initial concentration during the 24 days of exposure. The concentrations in the sediment phase increased in this period. The results are based on initial mean measured concentrations. Emergence in the control was 94% at day 24. Survival and number emerged were not significantly affected. Time to emerge and weight was significantly affected at the two highest concentration. The overall NOEC was determined to be 4.11 mg a.s./L (initial mean measured). In a reassessment (Brausch, 2015), EC10 for weight was determined to be 3.022 mg a.s./L. For emergence (time and number of males and females) the EC10 values was >15.6 mg a.s./L. The endpoints are considered reliable and can be used for classification purposes.

### 11.6.3 Chronic toxicity to algae and other aquatic plants

One additional studies on chronic toxicity to aquatic plants was included in the renewal dossier for dimethomorph since the previous evaluation. Some studies from the previous evaluation were re-evaluated for the renewal. From a literature search four publications were included in the RAR and assessed on their reliability. The descriptions below are based on the summaries in the RAR.

Aquatic toxicity to algae was tested by Ellgehausen (1986b) on *Scenedesmus subspicatus* in a 72 hours exposure according to OECD test guideline 201. The dimethomorph of technical grade with a purity of 94.8% was test in nominal concentrations of 10, 20, 40, 80 and 160 mg/L in three replicates. Test concentrations were verified by analysis and were within 80 to 123% of nominal for the concentration of 20 mg/L but 23.5 to 45.1% for the higher test concentrations. The latter two solutions were turbid because the concentrations exceed the water solubility and only clear solutions were analysed. The actual mean measured concentrations are not available. Over the exposure period inhibition of growth was observed from the lowest test concentration but significance of the observed inhibition is not reported in the RAR. Therefore a NOEC cannot be determined. Also an EC10 is not reported. In the RAR it was requested that the endpoint had to be based on mean measured concentration. In the absence of the raw data, this was not possible. Therefore it is also not expected that reliable chronic endpoints can be determined from this study. The test is considered unreliable and will not be used for classification purposes.

Aquatic toxicity to algae was tested by Jatzek (2001) on *Pseudokichneriella subcapitata* in a 72 hours exposure according to OECD test guideline 201. The dimethomorph of technical grade with a purity of 98.3% was tested in nominal concentrations of 6.25, 12.5, 25, 50 and 100 mg/L in three replicates. Test concentrations were verified by analysis and the concentrations ranged from 75.8 to 82.5% of the nominal concentrations at 0 hours, to 77.6 to 94.0% of the nominal concentrations at 72 hours. It is also reported that the two highest test concentrations which are at or above the water solubility, were turbid. It is unclear if analysis of the two higher test solutions is performed on clear solutions or that undissolved substance is included in the analysis. The actual mean measured concentrations are not presented in the RAR. Over the exposure period inhibition of growth was observed from the lowest test concentration but significance of the observed inhibition is not reported in the RAR. Therefore a NOEC cannot be determined. EC10 values for this study are reported in the List of end points of the RAR and are 27.3 mg a.s./L for growth rate and 9.23 mg a.s./L for biomass, both based on nominal data. In the RAR comments of the RMS are not provided on this study. However for the study of Ellgehausen (1986b) on *Scenedesmus subspicatus* it was requested that the endpoint had to be based on mean measured concentrations. It is presumed that this also applies to the endpoints of the study of Jatzek (2001), especially since reported measured endpoints are lower than 80% and turbidity is observed in the two higher test solutions. Therefore it is also not expected that reliable chronic endpoints can be determined from this study. The test is considered unreliable and will not be used for classification purposes.

Megateli et al. (2009, 2013) studied the effect of dimethomorph, as single substance or in combination with copper sulfate, on the growth of the aquatic plant *Lemna gibba* in a 168 h static toxicity test. The dimethomorph of technical grade with a purity of 97% was tested in nominal concentrations of 0.040, 0.200, 0.400, 0.800 and 1.000 mg/L in three replicates. The tests with copper sulfate are not relevant for classification and therefore not further discussed in this report. Test concentrations were verified by analysis and concentrations lower than 80% of the nominal concentrations were reported (10-40% of nominal). Raw data are not available and endpoints are based on nominal concentrations. NOECs or EC10 values are not reported in the RAR and it was stated that the study could not be used for the renewal application especially since endpoints are based on nominal concentrations while measured concentrations are well below the nominal concentrations. For the same reasons the study is considered unreliable and will not be used for classification purposes.

## **11.7 Comparison with the CLP criteria**

### **11.7.1 Acute aquatic hazard**

For dimethomorph, there are reliable acute data for fish and invertebrates. The lowest endpoint for fish is the value of 6.1 mg a.s./L for *Oncorhynchus mykiss* and for invertebrates this is 4.42 mg/L *Crassostrea virginica*. The lowest value of 4.42 mg a.s./L is above 1 mg/L, classification of dimethomorph as Aquatic acute is not applicable.

### **11.7.2 Long-term aquatic hazard (including bioaccumulation potential and degradation)**

Dimethomorph is hydrolytically stable at pH 4 to 9 which are environmental relevant pHs. There are two ready biodegradability studies (OECD 301B and 301D) available for dimethomorph. In the 301D test, no oxygen was consumed in 28 days and in the 301B, there was no evolution of carbon dioxide over 28 days. It was concluded that dimethomorph is not readily biodegradable. Inherent biodegradability was also assessed in a GLP study following the OECD 302C test guideline. Some evidence of inherent-primary biodegradability was observed (27% maximum biodegradation over 28 days). Low half-lives for some water-sediment systems are reported but since these are based on irreversible sorption to sediment, these cannot be considered as actual biodegradation and cannot be used for conclusion on rapid degradability.

On this basis, dimethomorph is considered not rapidly degradable and the chronic classification will be based on the criteria for non-rapidly degradable substances.

The highest BCF reported for dimethomorph is 2.0 and the measured LogK<sub>ow</sub> values range from 2.63 - 2.73. Therefore it is considered to have a low potential for bioaccumulation.

Reliable experimental chronic toxicity endpoints are available for fish and invertebrates. Where more than one acceptable toxicity value (NOEC and EC<sub>10</sub>) is available for a study, the EC<sub>10</sub> value is considered for classification. Where EC<sub>10</sub> values are available for a species, they are preferred over NOEC values for the same endpoint (ECHA, 2015, OECD, 2006).

The following toxicity values are available for fish: *O. mykiss* EC<sub>10</sub>= 0.116 mg/L, *C. variegatus* EC<sub>10</sub>=0.150 mg/L and *P. promelas* NOEC = 0.107 mg/L. The most sensitive chronic endpoint for fish is the NOEC of 0.107 mg/L for *Pimephales promelas* therefore it is selected as key study. For invertebrates the lowest values are EC<sub>10</sub> = 0.15 mg/L for *Daphnia magna*, NOEC/EC<sub>10</sub> of 0.24 mg/L for *A. Bahia* and EC<sub>10</sub> = 3.02 mg/L for *C. riparius*. The most sensitive chronic endpoint for invertebrates is the EC<sub>10</sub> of 0.15 mg/L.

There are no adequate acute and chronic data for algae and as a consequence the surrogate method cannot be applied for this substance.

The lowest value is of 0.107 mg/L and the substance is considered non-rapidly biodegradable. Based on the criteria set out in CLP, Annex I, section 4.1, Table 4.1.0(b) (i), dimethomorph fulfils the criteria for classification as Aquatic Chronic 2.

### **11.8 CONCLUSION ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS**

A classification for aquatic acute toxicity is not required.

The proposed classification for aquatic chronic toxicity is Aquatic chronic 2.

## **12 EVALUATION OF ADDITIONAL HAZARDS**

### **12.1 Hazardous to the ozone layer**

Not evaluated in this dossier

## **13 ADDITIONAL LABELLING**

*Not evaluated in this dossier*

## **14 REFERENCES**

- B.6.1.1.1a - Anonymous 1990, The Biokinetics and Metabolism of 14C-Dimethomorph in the Rat, BASF Document No. DK-440-001. As summarized in RAR 2017.
- B.6.1.1.1b – Anonymous 1991, 14C-Dimethomorph: Investigation on the Nature of the Metabolites Occurring in the Rat, BASF Document No. DK-440-006 [complementary data supporting study report DK-440-001] As summarized in RAR 2017.
- B.6.1.1.2 - Anonymous 1990, 14C-Dimethomorph (CME 151): Absorption, Distribution, Metabolism and Excretion after Bile Cannulation and Single Oral Administration to the Rat. BASF Document No. DK-440-002. As summarized in RAR 2017.
- B.6.1.1.3 – Anonymous 1993, 14C-Dimethomorph (CME 151): Investigation of the Metabolite Profiles in the Rat Following Treatment with two Different Labels, BASF Document No. DK-440-011. As summarized in RAR 2017.
- B.6.1.1.4 - Anonymous 1994, 14C-Dimethomorph: Additional Investigation on the Nature of the Metabolites Occurring in Rats, BASF Document No. DK-440-010. As summarized in RAR 2017.

- B.6.1.1.5 - Anonymous 1995, Dimethomorph (CL 336,379): Blood Pharmacokinetics of C-14 CL 336,379 Derived Residues in the Rat, BASF Document No. DK-452-008. As summarized in RAR 2017.
- B.6.1.1.6 - Anonymous, Dimethomorph (CL 336,379): Tissue distribution of C-14 CL 336,379 derived residues in the rat, BASF document No. DK-440-013 unpublished. As summarized in RAR 2017.
- B.6.1.1.7 - Anonymous 2015 BAS 550 F: Excretion and metabolism in the rat after oral administration 2015/1000602. As summarized in RAR 2017.
- B.6.3.1.1 - Anonymous 1985, ZTH 236 Z50 Preliminary assessment of toxicity to rats by dietary admixture for four weeks, unpublished, BASF Document No. DK-420-005. As summarized in RAR 2017.
- B.6.3.1.2 - Anonymous 1986, CME 151: 4 week dietary dose range finding study in rats, unpublished, BASF Document No. DK-420-002. As summarized in RAR 2017.
- B.6.3.1.3 - Anonymous 1990, SAG 151 E isomer: A 28 day oral toxicity study in rats, , unpublished, Shell Agrar Document No. 151AA-432-005. As summarized in RAR 2017.
- B.6.3.1.4 - Anonymous 1991, SAG 151 Z isomer: A 28 day oral toxicity study in rats, BASF document No. DK-470-016. As summarized in RAR 2017.
- B.6.3.1.5 - Anonymous 1986, CME 151 6 week dietary dose range finding study in mice,, unpublished. BASF Document No. DK-420-003. As summarized in RAR 2017.
- B.6.3.1.6 - Anonymous 1986, CME 151 Dietary maximum tolerated dose study in dogs, unpublished. BASF Document No. DK-420-001. As summarized in RAR 2017.
- B.6.3.2.1 - Anonymous 1987, CME 151: Toxicity to rats by dietary admixture for 13 weeks with a 4 week withdrawal period, unpublished, BASF Document No. DK-425-001. As summarized in RAR 2017.
- B.6.3.2.2 - Anonymous 1986, CME 151: 13 week dietary toxicity study in dogs, unpublished. BASF Document No. DK-425-002. As summarized in RAR 2017.
- B.6.3.2.3 - Anonymous 1990, SAG 151: 52 week dietary toxicity study in dogs, unpublished, BASF Document No. DK-427-003. As summarized in RAR 2017.
- B.6.3.3.1 - Anonymous 2010. a BAS 550 F (Dimethomorph) - Repeated-dose 28-day dermal toxicity study in Wistar rats. 2010/1151903. As summarized in RAR 2017.
- B.6.5.1.1-study 1 - Anonymous 1990, SAG 151: 104 week dietary toxicity study in rats, unpublished, BASF Document No, DK-427-006 (BASF Document No. DK-427-004 [Addendum BASF Document No. DK-427-005/1991 (revised figures)]). As summarized in RAR 2017.
- B.6.5.1.1-study 2 - Anonymous 1991, SAG 151: 104 week dietary carcinogenicity study in rats, unpublished, BASF Document No. DK-428-005. As summarized in RAR 2017.
- B.6.5.2 - Anonymous 1991, SAG 151: 104 week dietary carcinogenicity study in mice, unpublished, BASF Document No. DK-428-004. As summarized in RAR 2017.
- B.6.6.1.1. - Anonymous 1990, SAG 151: Two generation oral (dietary administration) reproduction toxicity study in the rat (two litters in the F1 generation), unpublished, BASF Document No. DK-430-001. As summarized in RAR 2017.
- B.6.6.1.2 - Anonymous 2014. Extended one-generation reproduction toxicity study in wistar rats - Administration via the diet. 2014/1181670. As summarized in RAR 2017.
- B.6.6.2.1 - Anonymous 1986: ZHT 236Z50 – Preliminary Oral (Gavage) Embryotoxicity Study in Rat, unpublished BASF Document No. DK-432-001. As summarized in RAR 2017.
- B.6.6.2.2 - Anonymous 1987: CME 151 – Letter Report: Treatment of Pregnant Sprague Dawley rats with CME 151; unpublished BASF Document No. DK-432-006. As summarized in RAR 2017.
- B.6.6.2.3 - Anonymous 1989, SAG 151: Oral (gavage) teratogenicity study in the rat, unpublished, BASF Document No. DK-432-002. As summarized in RAR 2017.

- B.6.6.2.4 - Anonymous 1987: CME 151 – Preliminary Oral (Gavage) Embryotoxicity Study in the Rabbit; unpublished. BASF Document No. DK-432-003. As summarized in RAR 2017.
- B.6.6.2.5 - Anonymous 1989, SAG 151: Oral (gavage) teratogenicity study in the rabbit, unpublished, BASF Document No. DK-432-004. As summarized in RAR 2017.
- B.6.7.1.2 - Anonymous 2004a. BAS 550 F - Subchronic neurotoxicity study in Wistar rats - Administration in the diet for 3 months. 2004/1013810. As summarized in RAR 2017.
- B.6.8.2.15 - Anonymous 2010b. BAS 550 F (Dimethomorph) - Immunotoxicity study in male Wistar rats - Administration via the diet for 4 weeks. 2010/1043718. As summarized in RAR 2017
- Archer. E, et al, 2015. The potential anti-androgenic effect of agricultural pesticides used in the Western Cape: In vitro investigation of mixture effects. *Water SA*, 41 (1), 129-137. As summarized in RAR 2017.
- Birks V., 2015. 14C-BAS 550 F: Comparative in-vitro metabolism studies with rat, dog and human cryopreserved hepatocytes. 2015/1245078. As summarized in RAR 2017.
- Bitsch N. et al, 2002. In vitro screening of the estrogenic activity of active components in pesticides (in German). *Z. Umweltchem. Okotox.* 14 (2) 76-84. As summarized in RAR 2017.
- Carney et al, 2004. The Effects of Feed Restriction during in Utero and Postnatal Development in Rats. *Toxicological sciences* 82: 237-249
- Miodovnik, A. et al, 2014. Developmental neurotoxicity of ortho-phthalate diesters: Review of human and experimental evidence. *NeuroToxicology* 41:112-122.
- Orton F. et al, 2011. Widely used pesticides with previously unknown endocrine activity revealed as in vitro anti-androgens. *Environ. Health Perspect.*, 109 (6), 794-800. As summarized in RAR 2017.
- Orton F. et al, 2014. Mixture effects at very low doses with combination of anti-androgenic pesticides, antioxidants, industrial pollutant and chemicals used in personal care products. *Tox. Appl. Pharmacol.*, 278, 201-208. As summarized in RAR 2017.
- RAR Dimethomorph 2017
- Registration dossier Dimethomorph
- Reif D.M. et al, 2010. Endocrine profiling and prioritization of environmental chemicals using ToxCast data. *Environ. Health Perspect.*, 118 (12), 1714-1720. As summarized in RAR 2017.
- Woitkowiak C., 2011. BAS 550 F (Dimethomorph) - Testing for potential androgenic and antiandrogenic activity using the YAS-assay (AR) (yeast androgen screening). 2011/1140605. As summarized in RAR 2017.
- Woitkowiak C., 2011. BAS 550 F (Dimethomorph) - Testing for potential estrogenic and antiestrogenic activity using the YES-assay (ERalpha) (yeast estrogen screening). 2011/1140606. As summarized in RAR 2017.

### References for ecotoxicity chapter 11

- Anonymous. 1993. Determination of the concentrations of Dimethomorph (SAG 151) in test medium. Rossdorf, Germany: RCC Umweltchemie GmbH & Co. KG. Report no. DK-123-221, As summarized in RAR 2017.
- B.9.2.1.1 – Anonimous. 1986. 96-hour acute toxicity (LC50) with CME 151 in the rainbow trout. Itingen, Switzerland: RCC Umweltchemie AG. Report no. DK-511-002, As summarized in RAR 2017.
- B.9.2.1.2 – Anonimous. 2010. BAS 550 F - Re-calculation of toxicity values for study DK-511-001 (96-hour acute toxicity study (LC50) with CME 151 in the carp). Limburgerhof, Germany: BASF SE. Report no. 2010/1177241, As summarized in RAR 2017.
- B.9.2.1.3 – Anonimous. 1986. 96-hour acute toxicity (LC50) with CME 151 in the carp. Itingen, Switzerland: RCC Umweltchemie AG. Report no. DK-511-001, As summarized in RAR 2017.



- B.9.2.1.4 – Anonymous. 2010. BAS 550 F - Re-calculation of toxicity values for study DK-511-002 (96-hour acute toxicity study (LC50) with CME 151 in the rainbow trout). Limburgerhof, Germany: BASF SE. Report no. 2010/1177242, As summarized in RAR 2017.
- B.9.2.1.5 – Anonymous. 1988. The acute toxicity of CME 151 to bluegill sunfish (*Lepomis macrochirus*). Huntingdon, UK: Huntingdon Research Centre Ltd. Report no. DK-511-003, As summarized in RAR 2017.
- B.9.2.1.6 – Anonymous. 2010. BAS 550 F - Re-calculation of toxicity values for study DK-511-003 (The acute toxicity of CME 151 to bluegill sunfish (*Lepomis macrochirus*)). Limburgerhof, Germany: BASF SE. Report no. 2010/1177243, As summarized in RAR 2017.
- B.9.2.2.6 – Anonymous. 2010. BAS 550 F: Early life-stage toxicity test with the sheepshead minnow, *Cyprinodon variegatus*, under flow-through conditions. Columbia, USA: ABC Laboratories Inc. Report no. 2010/7012691, As summarized in RAR 2017.
- B.9.2.1.7 – Anonymous. 2001. Acute toxicity of BAS 550 F (AC 336379, Dimethomorph) to rainbow trout, *Oncorhynchus mykiss*, under flow-through test conditions. Report no. DK-511-007, As summarized in RAR 2017.
- B.9.2.1.8 – Anonymous. 1997. Acute toxicity of AC 336379 (Dimethomorph) to the sheepshead minnow (*Cyprinodon variegatus*) under flow-through test conditions. Report no. DK-511-005, As summarized in RAR 2017.
- B.9.2.1.9 – Anonymous. 2001. Acute toxicity of BAS 550 F (AC 336379, Dimethomorph) to bluegill sunfish, *Lepomis macrochirus* under flow-through test conditions. Report no. DK-511-006, As summarized in RAR 2017.
- B.9.2.1.10 - Anonymous. 2014. BAS 550 F: A 96-hour flow-through acute toxicity test with the fathead minnow (*Pimephales promelas*). Easton MD, USA: Wildlife International Ltd. Report no. 2014/7002827, As summarized in RAR 2017.
- B.9.2.2.1 - Anonymous. 1997. Toxicity of AC 336,379 (Dimethomorph) technical during the early life-stages of rainbow trout (*Oncorhynchus mykiss*). Princeton, USA: American Cyanamid Co. Report no. DK-512-002, As summarized in RAR 2017.
- B.9.2.2.2 - Anonymous. 2015. Calculation of EC10 values and geomean for chronic fish studies with BAS 550 F (Dimethomorph). Report no. 2014/1224007, As summarized in RAR 2017.
- B.9.2.2.5 - Anonymous. 2002. BAS 550 F - Early life-stage toxicity test on the Fathead minnow (*Pimephales promelas*). Ludwigshafen/Rhein, Germany: BASF AG. Report no. 2002/1011268, As summarized in RAR 2017.
- B.9.2.3.1 - Anonymous. 1999. Dimethomorph (AC 336379): Uptake, depuration, bioconcentration and metabolism of carbon-14 labeled AC 336379 in bluegill sunfish (*Lepomis macrochirus*) under flow-through conditions. Princeton, USA: American Cyanamid Co. Report no. DK-519-001, As summarized in RAR 2017.
- B.9.2.4.1 - Anonymous. 2014. BAS 550 F (Dimethomorph) - Fish short term reproduction assay on the fathead minnow (*Pimephales promelas*). Ludwigshafen/Rhein, Germany: BASF SE. Report no. 2014/1187237, As summarized in RAR 2017.
- Battersby N. 1993. Dimethomorph (CME-151): An assessment of inherent biodegradability in the modified MITI(II) test. Report no. DK-690-005. As summarized in RAR 2017.
- Brausch JM. 2015. Calculation of EC10 values for chronic invertebrate studies with BAS 550 F (Dimethomorph) BASF SE, Limburgerhof, Germany. Report no. 2015/1238096, As summarized in RAR 2017.
- Ebert D. 2002. Dimethomorph degradation and metabolism in aquatic systems. Report no. DK-390-060. As summarized in RAR 2017.
- ECHA. 2015. Guidance on the Application of the CLP Criteria - Guidance to Regulation (EC) No 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures - Version 4.1. Helsinki: ECHA. Report no. ECHA-15-G-05-EN.
- Ellgehausen H. 1986a. 48-hour acute toxicity of CME 151 to *Daphnia magna* (OECD immobilization test). Itingen, Switzerland: RCC Umweltchemie AG. Report no. DK-521-002, As summarized in RAR 2017.
- Ellgehausen H. 1986b. Acute toxicity of CME 151 to *Scenedesmus subspicatus* (OECD algae growth inhibition test). Itingen, Switzerland: RCC Umweltchemie AG. Report no. DK-521-001, As summarized in RAR 2017.

- England DC, al. e. 1997. Effects of AC 336,379 (Dimethomorph) on the development of sediment-dwelling larvae of *Chironomus riparius* in a water-sediment system. Princeton, USA: American Cyanamid Co. Report no. DK-529-002, As summarized in RAR 2017.
- Habekost M. 2010. BAS 550 F - Re-calculation of toxicity values for study DK-521-002. Limburgerhof, Germany: BASF SE. Report no. 2010/1177164, As summarized in RAR 2017.
- Hicks SL. 2010. BAS 550 F: Life-cycle toxicity test of the saltwater mysid, *Americamysis bahia*, conducted under flow-through conditions. Columbia, USA: ABC Laboratories Inc. Report no. 2010/7008279, As summarized in RAR 2017.
- Jatzek H-J. 2001. BAS 550 F - Determination of the inhibitory effect on the cell multiplication of unicellular green algae. Report no. DK-521-007, As summarized in RAR 2017.
- Klimisch H-J, Andreae M, Tillmann U. 1997. A Systematic Approach for Evaluating the Quality of Experimental Toxicological and Ecotoxicological Data. Regulatory Toxicology and Pharmacology. 25: 1-5.
- Knoch E. 1993. Dimethomorph degradation and metabolism in aquatic systems. Report no. DK-630-002. As summarized in RAR 2017.
- Knoch E. 1994a. Humin fraction determination in selected sediments of a water/sediment study of Dimethomorph (SAG 151) (Supplemental data to BASF Document No. DK-630-002). Report no. DK-630-005. As summarized in RAR 2017.
- Knoch E. 1994b. Screening of a not identified metabolite fraction of a water/sediment study of Dimethomorph (SAG 151) (Supplemental data to BASF Document No. DK-630-002). Report no. DK-630-006. As summarized in RAR 2017.
- Knoch E, Holman J. 1998. Dimethomorph (AC 336379): Determination of the direct phototransformation in buffered medium at pH 7. Report no. DK-324-004, As summarized in RAR 2017.
- Maleri M. 2015. Kinetic evaluation of the aerobic soil degradation studies for BAS 550 F - Dimethomorph according to FOCUS. Limburgerhof, Germany: BASF SE. Report no. 2014/1183299, As summarized in RAR 2017.
- Megateli S, al. e. 2009. Toxicity of Copper/Dimethomorph combination for *Lemna minor* and depuration of the fungicides by aquatic plant. Report no. 2009/1131842, As summarized in RAR 2017.
- Megateli S, al. e. 2013. Simultaneous effects of two fungicides (Copper and Dimethomorph) on their phytoremediation using *Lemna minor*. Report no. 2013/1419680, As summarized in RAR 2017.
- Memmert U, Knoch E. 1993. Influence of Dimethomorph on survival and reproduction of *Daphnia magna* in a semistatic test (22 days). Rossdorf, Germany: RCC Umweltchemie GmbH & Co. KG. Report no. DK-524-001, As summarized in RAR 2017.
- Mitchell GC. 1997a. Acute toxicity of AC 336379 (Dimethomorph) to the mysid (*Mysidopsis bahia*) under flow-through test conditions. Report no. DK-521-004, As summarized in RAR 2017.
- Mitchell GC. 1997b. Effect of AC 336379 (Dimethomorph) on new shell growth in the Eastern oyster (*Crassostrea virginica*) under flow-through test conditions. Report no. DK-522-001, As summarized in RAR 2017.
- Mitchell GC. 2001. Acute toxicity of BAS 550 F (AC 336379, Dimethomorph) to *Daphnia magna* under static test conditions. Report no. DK-521-006, As summarized in RAR 2017.
- Murrell HR. 1997. Chronic toxicity of AC 336,379 (Dimethomorph) technical during the complete life-cycle of *Daphnia magna* under flow-through test conditions. Princeton, USA: American Cyanamid Co. Report no. DK-523-001, As summarized in RAR 2017.
- Ochsenein U. 1989. Hydrolysis determination of <sup>14</sup>C-Dimethomorph (CME 151) at different pH values. Report no. DK-322-003. As summarized in RAR 2017.
- OECD. 2006. Series on testing and assessment number 54, Current approaches in the statistical analysis of ecotoxicity data: a guidance to application. Paris: OECD. Report no. ENV/JM/MONO(2006)18.
- Panek MP, Mills C, Bolin D. 2001. Aqueous photolysis of 14C-BAS 550 F. Report no. DK-324-006, As summarized in RAR 2017.
- Turner S. 1988. CMT-151 (WL127294) An assessment of ready biodegradability. Report no. DK-690-001. As summarized in RAR 2017.
- Van Dijk A. 1990. Photodegradation study of 14-C Dimethomorph (CME 151) in water. Report no. DK-630-001, As summarized in RAR 2017.
- Yeomans P. 2015. 14C-Dimethomorph (BAS 550 F): Aerobic mineralisation in surface water. Harrogate, UK: Smithers Viscient (ESG) Ltd. Report no. 2014/1184744, As summarized in RAR 2017.

**15 ANNEXES**

*An Annex I dossier without confidential information is available.*

*An Annex II with a confidential reference list is provided*