

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

L'APPAT CEREALES DES RONGEURS

Product type(s)

PT14: Rodenticides

Authorisation number: FR-2012-0505

R4BP asset number: FR-0001002-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

| | |
|---------------|--|
| Trade name(s) | L'APPAT CEREALES DES RONGEURS BLE MEUNIER SOLUTION RAKAO GRAIN DF 25 KA TUE RATICIDE 25 KA TUE SOURICIDE 25 RACAN BLE 25 SUPRA GRAIN VULCANO GRAIN 25 |
|---------------|--|

1.2. Authorisation holder

| | | |
|--|---------|---|
| Name and address of the authorisation holder | Name | LODI S.A.S. |
| | Address | Parc d'Activités des Quatre Routes 35390 Grand Fougeray France |
| Authorisation number | | FR-2012-0505 |
| <i>R4BP asset number</i> | | FR-0001002-0000 |
| Date of the authorisation | | 28/12/2012 |
| Expiry date of the authorisation | | 31/12/2026 |

1.3. Manufacturer(s) of the product

| | |
|---------------------------------|--|
| Name of manufacturer | LODI SAS |
| Address of manufacturer | PA des Quatre Routes 35390 Grand-Fougeray France |
| Location of manufacturing sites | PA des Quatre Routes 35390 Grand Fougeray France |

1.4. Manufacturer(s) of the active substance(s)

| | |
|---------------------------------|---|
| Active substance | Difenacoum |
| Name of manufacturer | Pelgar International Ltd |
| Address of manufacturer | Unit 13, Newman Lane Industrial Estate, Newman Lane, Alton Hampshire GU34 2 QR Alton Hampshire United Kingdom of Great Britain and Northern Ireland (the) |
| Location of manufacturing sites | Prqazska 54, 280 02 Kolin Czechia |

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

| Common name | IUPAC name | Function | CAS number | EC number | Content (%) |
|-------------|---|------------------|------------|-----------|-------------|
| Difenacoum | 3-(3-biphenyl-4-yl-1,2,3,4-tetrahydro-1-naphthyl)-4-hydroxycoumarin | active substance | 56073-07-5 | 259-978-4 | 0,0025 |

2.2. Type(s) of formulation

RB-(bait ready for use) grain

3. HAZARD AND PRECAUTIONARY STATEMENTS

| | |
|--------------------------|--|
| Hazard statements | H373: May cause damage to organs (blood) through prolonged or repeated exposure {1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard:}. |
| Precautionary statements | P260: Do not breathe dust/fume/gas/mist/vapours/spray. P314: Get medical advice/attention if you feel unwell. P501: Dispose of {0:....:contents container} to {1:....:}. |

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice and/or rats – trained professionals – indoor

| | |
|--|--|
| Product type | PT14: Rodenticides |
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Indoor |
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mice Development stage: Juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: black or roof rat Development stage: Juveniles and adults |
| Field(s) of use | indoor use Indoor |
| Application method(s) | Method: bait application Detailed description: Bait formulations:- Ready-to-use bait to be used in tamper-resistant bait stations - [Covered and protected baiting points] |
| Application rate(s) and frequency | Application rate: Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters Dilution (%): 0 Number and timing of application: Bait products: - Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters |
| Category(ies) of users | trained professional |
| Pack sizes and packaging material | The product is wrapped in individual PE sachet (25, 50, 100g) or in loose. Sachets are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) Loose grains are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) |

| | |
|--|--|
| | - in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg |
|--|--|

(In France only : minimum pack size of 5 kg)

4.1.1. Use-specific instructions

- Remove the remaining product at the end of treatment period.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use the product in pulsed baiting treatments.

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.2. Use description

Table 2. Mice and/or rats – trained professionals – outdoor around buildings

| | |
|--------------|--------------------|
| Product type | PT14: Rodenticides |
|--------------|--------------------|

| | |
|--|---|
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Outdoor around buildings |
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mice Development stage: Juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: black or roof rat Development stage: Juveniles and adults |
| Field(s) of use | outdoor use Outdoor around buildings |
| Application method(s) | Method: bait application Detailed description: Bait formulations:- Ready-to-use bait to be used in tamper-resistant bait stations.- [Covered and protected baiting points] |
| Application rate(s) and frequency | Application rate: Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters Dilution (%): 0 Number and timing of application: Bait products: - Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters |
| Category(ies) of users | trained professional |
| Pack sizes and packaging material | The product is wrapped in individual PE sachet (25, 50, 100g) or in loose. Sachets are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) Loose grains are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg <i>(In France only : minimum pack size of 5 kg)</i> |

| | |
|--|--|
| | |
|--|--|

4.2.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.3. Use description

Table 3. Mice and/or Rats – trained professionals – Outdoor open areas & waste dumps

| | |
|--------------|--------------------|
| Product type | PT14: Rodenticides |
|--------------|--------------------|

| | |
|--|---|
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Outdoor open areas Outdoor waste dumps |
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mice Development stage: Juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: black or roof rat Development stage: Juveniles and adults |
| Field(s) of use | outdoor use Autre Outdoor open areas Outdoor waste dumps |
| Application method(s) | Method: bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations.- [Covered and protected baiting points] |
| Application rate(s) and frequency | Application rate: Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters Dilution (%): 0 Number and timing of application: Bait products: - Rats : 100 g of bait per baiting point every 5 to 10 meters - Mice: 25-30 g of bait per baiting point every 3 to 5 meters |
| Category(ies) of users | trained professional ; professional |
| Pack sizes and packaging material | The product is wrapped in individual PE sachet (25, 50, 100g) or in loose. Sachets are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) Loose grains are packed - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg <i>(In France only : minimum pack size of 5 kg)</i> |

| | |
|--|--|
| | |
|--|--|

4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period [Not applicable where explicitly authorised according to addenda 4].
- [When available] Follow any additional instructions provided by the relevant code of best practice.
- [For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species].

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.4. Use description

Table 4. House mice – general public – indoor

| | |
|--|---|
| Product type | PT14: Rodenticides |
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Indoor |

| | |
|--|---|
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mice Development stage: Juveniles and adults |
| Field(s) of use | indoor use Indoor |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait [in sachets for loose bait] to be used in tamper-resistant bait stations |
| Application rate(s) and frequency | Application rate: 25-30 g of bait per bait station Dilution (%): 0 Number and timing of application: Bait products: - 25-30 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 3 to 5 meters. |
| Category(ies) of users | general public (non-professional) |
| Pack sizes and packaging material | Maximum pack size: 150g The product is wrapped in individual sachet (25, 50, 100g). Sachets are packed - in pre-filled bait stations (PVC, PP, PS)(25, 50, 100g) - In carton box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In PP or PE box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In metal box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - PE/MET PET/PET pouch, PET/PE pouch, PE pouch, OPP pouch or PET/PE pouch with zip: 50, 75, 90, 100, 110, 120, 125, 130, 140, 150 g |

4.4.1. Use-specific instructions

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.4.2. Use-specific risk mitigation measures

4.4.3. Use-specific risk mitigation measures

-

4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.5. Use description

Table 5. Rats – general public – indoor

| | |
|--|--|
| Product type | PT14: Rodenticides |
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Indoor |
| Target organism(s) (including development stage) | Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> (in france only) Common name: roof rat Development stage: juveniles and adults |
| Field(s) of use | indoor use Indoor |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait [in sachets for loose bait] to be used in tamper-resistant bait stations |
| Application rate(s) and frequency | Application rate: 100 g of bait per bait station Dilution (%): 0 Number and timing of application: Bait products: - 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 to 10 meters. |
| Category(ies) of users | general public (non-professional) |

| | |
|-----------------------------------|--|
| Pack sizes and packaging material | <p>Maximum pack size: 150g The product is wrapped in individual sachet (25, 50, 100g).</p> <p>Sachets are packed</p> <ul style="list-style-type: none"> - in pre-filled bait stations (PVC, PP, PS)(25, 50, 100g) - In carton box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In PP or PE box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In metal box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - PE/MET PET/PET pouch, PET/PE pouch, PE pouch, OPP pouch or PET/PE pouch with zip: 50, 75, 90, 100, 110, 120, 125, 130, 140, 150 g |
|-----------------------------------|--|

4.5.1. Use-specific instructions

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.5.2. Use-specific risk mitigation measures

4.5.3. Use-specific risk mitigation measures

-

4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.6. Use description

Table 6. Rats – general public – outdoor around buildings

| | |
|--------------|--------------------|
| Product type | PT14: Rodenticides |
|--------------|--------------------|

| | |
|--|---|
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Outdoor around buildings |
| Target organism(s) (including development stage) | Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Juveniles and adults Scientific name: <i>Rattus rattus</i> (in France only) Common name: black or roof rat Development stage: Juveniles and adults |
| Field(s) of use | outdoor use Outdoor around buildings |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait [in sachets for loose bait] to be used in tamper-resistant bait stations |
| Application rate(s) and frequency | Application rate: 100 g of bait per bait station Dilution (%): 0 Number and timing of application: Bait products: - 100 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 to 10 meters. |
| Category(ies) of users | general public (non-professional) |
| Pack sizes and packaging material | Maximum pack size: 150g The product is wrapped in individual sachet (25, 50, 100g). Sachets are packed - in pre-filled bait stations (PVC, PP, PS)(25, 50, 100g) - In carton box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In PP or PE box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - In metal box (50, 75, 90, 100, 110, 120, 125, 130, 140, 150g) - PE/MET PET/PET pouch, PET/PE pouch, PE pouch, OPP pouch or PET/PE pouch with zip: 50, 75, 90, 100, 110, 120, 125, 130, 140, 150 g |

4.6.1. Use-specific instructions

- Place the bait stations in areas not liable to flooding.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

-

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.7. Use description

Table 7. (not relevant in France)– House mice – professionals – indoor

| | |
|--|---|
| Product type | PT14: Rodenticides |
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Indoor |
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles and adults |
| Field(s) of use | indoor use Indoor |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations [1] |
| Application rate(s) and frequency | Application rate: Mice: 25-30 g of bait per baiting point every 3 to 5 meters Dilution (%): 0 Number and timing of application: |

| | |
|-----------------------------------|--|
| | Bait products: - Mice: 25-30 g of bait per baiting point every 3 to 5 meters |
| Category(ies) of users | professional |
| Pack sizes and packaging material | <p>The product is wrapped in individual PE sachet (25, 50, 100g) or in loose.</p> <p>Sachets are packed</p> <ul style="list-style-type: none"> • in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) • in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) <p>Loose grains are packed</p> <ul style="list-style-type: none"> • in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) • in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg <p><i>(In France only : minimum pack size of 5 kg)</i></p> |

4.7.1. Use-specific instructions

- The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice.

4.7.2. Use-specific risk mitigation measures

4.7.3. Use-specific risk mitigation measures

-

4.7.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.7.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.7.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.8. Use description

Table 8. (not relevant in France)– Rats – professionals – indoor

| | |
|--|--|
| Product type | PT14: Rodenticides |
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides indoor |
| Target organism(s) (including development stage) | Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: juveniles and adults |
| Field(s) of use | indoor use indoor |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations |
| Application rate(s) and frequency | Application rate: Rats : 100 g of bait per baiting point every 5 to 10 meters Dilution (%): 0 Number and timing of application: Bait products: • Rats : 100 g of bait per baiting point every 5 to 10 meters |
| Category(ies) of users | professional |
| Pack sizes and packaging material | The product is wrapped in individual PE sachet (25, 50, 100g) or in loose. Sachets are packed • in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) |

| | |
|--|--|
| | <ul style="list-style-type: none"> • in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) <p>Loose grains are packed</p> <ul style="list-style-type: none"> • in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) • in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg <p><i>(In France only : minimum pack size of 5 kg)</i></p> |
|--|--|

4.8.1. Use-specific instructions

- The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice.

4.8.2. Use-specific risk mitigation measures

4.8.3. Use-specific risk mitigation measures

-

4.8.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.8.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.8.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.9. Use description

Table 9. (not relevant in France)– House mice and/or rats – professionals – outdoor around buildings

| | |
|--------------|--------------------|
| Product type | PT14: Rodenticides |
|--------------|--------------------|

| | |
|--|---|
| Where relevant, an exact description of the authorised use | Not relevant for rodenticides Outdoor around buildings |
| Target organism(s) (including development stage) | Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: juveniles and adults |
| Field(s) of use | outdoor use Outdoor around buildings |
| Application method(s) | Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations |
| Application rate(s) and frequency | Application rate: Rats : 100 g of bait per baiting point every 5 to 10 meters Mice: 25-30 g of bait per baiting point every 3 to 5 meters Dilution (%): 0 Number and timing of application: Bait products: <ul style="list-style-type: none"> • Rats : 100 g of bait per baiting point every 5 to 10 meters • Mice: 25-30 g of bait per baiting point every 3 to 5 meters |
| Category(ies) of users | professional |
| Pack sizes and packaging material | The product is wrapped in individual PE sachet (25, 50, 100g) or in loose. Sachets are packed <ul style="list-style-type: none"> - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in carton box (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) Loose grains are packed <ul style="list-style-type: none"> - in PP or PE bucket (1, 1.5, 2, 2.5, 3, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10kg) - in Cardboard box with inner liner in PE / paper craft bag with inner liner in PE : 5 and 10 kg <i>(In France only : minimum pack size of 5 kg)</i> |

| | |
|--|--|
| | |
|--|--|

4.9.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [*for mice* - at least every 2 to 3 days at] [*for rats* - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [*When available*] Follow any additional instructions provided by the relevant code of best practice.

4.9.2. Use-specific risk mitigation measures

4.9.3. Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.9.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.9.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.9.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
 - Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
 - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
 - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
 - The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).
- *[If national policy or legislation requires it]* When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
 - Bait should be secured so that it cannot be dragged away from the bait station.
 - Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
 - Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

FOR TRAINED PROFESSIONAL ONLY - *The* frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

FOR PROFESSIONALS ONLY Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

FOR PROFESSIONALS ONLY Remove the remaining bait or the bait stations at the end of the treatment period.

- Do not open the sachets containing the bait.
- *Loose grains:* Decanting is to be avoided. In case decanting cannot be avoided, an RPE of APF 10 has to be used.

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

-
- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
 - Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
 - Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
 - Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - Where possible, bait stations must be fixed to the ground or other structures.
 - *[Do not open the sachets containing the bait - where relevant for the bait formulation in the product].*
 - Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
 - Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
 - Do not place bait stations near water drainage systems where they can come into contact with water.
 - When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
 - Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign *[in accordance with the applicable code of good practice, if any]*.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- **FOR TRAINED PROFESSIONAL ONLY** Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- **FOR TRAINED PROFESSIONAL ONLY** Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements *[The method of disposal shall be described specifically in the national SPC and be reflected on the product label]*.
- **FOR PROFESSIONAL ONLY** To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week). *[Where relevant, specify if more frequent or daily inspection is required]*.
- **FOR PROFESSIONAL ONLY** Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- **FOR PROFESSIONAL ONLY.** The product information (i.e. label and/or leaflet) shall clearly show that:
 - the product shall not be supplied to the general public (e.g. "for professionals only").
 - the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").

- **FOR PROFESSIONAL ONLY** Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).

- The product information (i.e. label and/or leaflet) shall clearly show that:

the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").

users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").

- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.

- Dispose dead rodents in accordance with local requirements [*The method of disposal shall be described specifically in the national SPC and be reflected on the product label*].

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

- Dermal exposure, wash skin with water and then with water and soap.

- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case of ingestion by a pet [insert country specific information]

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number]"

- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements [*The method of disposal shall be described specifically in the national SPC and be reflected on the product label*].

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

- Store in places prevented from the access of children, birds, pets and farm animals.

- Shelf life: 2 years

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.
- The authorisation holder has to monitor the resistance phenomenon of rodent populations toward the active substance difenacoum. Results of the resistance monitoring must be submitted at the renewal of the product.)Provide a field test on *R. rattus* within 1 year in post authorisation.