

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

Escatop Pasta Extreme

Product type(s)

PT14: Rodenticides

Authorisation number: IT/2018/00470/AUT

R4BP asset number: IT-0018676-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	Escatop Pasta Extreme Murin Pasta Plus
---------------	---

1.2. Authorisation holder

Name and address of the authorisation holder	Name	Vebi Istituto Biochimico s.r.l.
	Address	via Desman, 43 35010 Borgoricco Italy
Authorisation number	IT/2018/00470/AUT	
<i>R4BP asset number</i>	IT-0018676-0000	
Date of the authorisation	29/04/2015	
Expiry date of the authorisation	31/12/2026	

1.3. Manufacturer(s) of the product

Name of manufacturer	Vebi Istituto Biochimico s.r.l.
Address of manufacturer	via Desman, 43 35010 Borgoricco (PD) Italy
Location of manufacturing sites	via Desman, 43 35010 Borgoricco (PD) Italy

1.4. Manufacturer(s) of the active substance(s)

Active substance	Brodifacoum
Name of manufacturer	Dr Tezza s.r.l.
Address of manufacturer	via Tre Ponti, 22 37050 Santa Maria di Zevio (VR) Italy
Location of manufacturing sites	via Tre Ponti, 22 37050 Santa Maria di Zevio (VR) Italy

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	active substance	56073-10-0	259-980-5	0,0025

2.2. Type(s) of formulation

PA Paste

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H373: May cause damage to organs {0:or state all organs affected, if known:} through prolonged or repeated exposure { 1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard:}.
Precautionary statements	P103: Read carefully and follow all instructions. P270: Do not eat, drink or smoke when using this product. P273: Avoid release to the environment. P301+P310: IF SWALLOWED: Immediately call a doctor. P501: Dispose of contents to in accordance to national law. P102: Keep out of reach of children.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. General public

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: adults Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juveniles Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: adults Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: juveniles Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: adults Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles
Field(s) of use	indoor use outdoor use Mice: indoor Rats indoor and outdoor around buildings
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: rat 150 g, mice 50 g Dilution (%): 0 Number and timing of application: Mice Bait products: - 50 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. In case of high infestation level, applications sites are located 2 meters apart Mice - The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

	<p>Rats: Bait products: - 150 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 10 meters. In case of high infestation level, applications sites are located 5 meters apart</p> <p>Rats - The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p>
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	<p>MURIN PASTA Extreme is furnished in two different type:</p> <ol style="list-style-type: none"> 1. Bait of pasta each containing from 10 to 15g, wrapped into paper food,- 2. Plastic tray each containing 25, 30, 40, 50, 75, 80, 100g <p>Pack size: <i>from 50 to 150 g</i></p> <p>Packaging material:</p> <ul style="list-style-type: none"> - envelope, Plastic: coextruded/ HDPE/polethylene - Can /Tin, Paper, cardboard , - Box, Paper, cardboard with envelope inside coextruded/ HDPE/ polethylene

4.1.1. Use-specific instructions

- Place the bait stations in areas not liable to flooding.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Do not open the sachets containing the bait
- When the product is used in single-dose plastic trays, once positioned cut the upper protective film.
- Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.
- Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Do not place bait stations near water drainage systems where they can come into contact with water.

-
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
 - Remove the remaining bait or the bait stations at the end of the treatment period.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Do not use anticoagulant rodenticides as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).
- The product information (i.e. label and/or leaflet) shall clearly show that: the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only"). Users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. "label bait stations according to the product recommendations").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
- Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.
- Dispose dead rodents in accordance with local requirements

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "Murin pasta defend or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".
- Hazardous to wildlife.

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements
- Use of gloves is recommended.

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight .
 - Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life 2 years

4.2. Use description

Table 2. Professional user

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	<p>Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: adults</p> <p>Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juveniles</p> <p>Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: adults</p> <p>Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: juveniles</p> <p>Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: adults</p> <p>Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles</p>
Field(s) of use	<p>indoor use outdoor use</p> <p>indoor and outdoor around buildings</p>
Application method(s)	<p>Method: bait application</p> <p>Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p>
Application rate(s) and frequency	<p>Application rate: rat 200 g, mice 50 g</p> <p>Dilution (%): 0</p> <p>Number and timing of application:</p>

	<p>Mice Bait products: - 50 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. In case of high infestation level, applications sites are located 2 meters apart</p> <p>Mice - The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p> <p>Rats: Bait products: - 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 10 meters. In case of high infestation level, applications sites are located 5 meters apart</p> <p>Rats - The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p>
Category(ies) of users	industrial ; professional
Pack sizes and packaging material	<p>MURIN PASTA Extreme is furnished in two different type:</p> <ol style="list-style-type: none"> 1. Bait of pasta each containing from 10 to 15g, wrapped into paper food,- 2. Plastic tray each containing 25, 30, 40, 50, 75, 80, 100g <p>Pack size: <i>from 1,5 kg to 25 kg</i></p> <p><i>In case of packaging higher than 10 kg Package is restricted to separately packed bags with a maximum bag size of 5 or 10 kg</i></p> <p>Packaging material:</p> <ul style="list-style-type: none"> - envelope, Plastic: coextruded/ HDPE/polyethylene - Can /Tin, Paper, cardboard , - Box, Paper, cardboard with envelope inside coextruded/ HDPE/ polyethylene - bucket: HDPE

4.2.1. Use-specific instructions

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

- Where possible, bait stations must be fixed to the ground or other structures.

- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).

- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

- Bait should be secured so that it cannot be dragged away from the bait station.

- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

- *Wear protective chemical resistant gloves during product handling phase*

- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait stations to further places and the possibility to change to another bait formulation.

- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

- Remove the remaining bait or the bait stations at the end of the treatment period.

- *Do not open the sachets containing the bait*

- *When the product is used in single-dose plastic trays, once positioned cut the upper protective film.*

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign

- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week). [

- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.

- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

- The product information (i.e. label and/or leaflet) shall clearly show that: the product shall not be supplied to the general public (e.g. "for professionals only").

-
- the product shall be used in adequate tamper resistant bait stations (e.g. "use in tamper resistant bait stations only").
 - users shall properly label bait stations with the information referred to in section 5.3 of the SPC (e.g. label bait stations according to the product recommendations").
 - Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.
 - Do not wash the bait stations with water between applications.
 - Dispose dead rodents in accordance with local requirements
- Do not apply directly in the burrows

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "Murin pasta defend or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".
- Hazardous to wildlife.

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements
- Use of gloves is recommended.

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life 2 years

4.3. Use description

Table 3. Trained Professional

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	<p>Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: adults</p> <p>Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juveniles</p> <p>Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: adults</p> <p>Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: juveniles</p> <p>Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: adults</p> <p>Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles</p>
Field(s) of use	<p>indoor use outdoor use</p> <p>Mice and rats: indoor and outdoor around buildings Rats: Outdoor open areas Rats: Outdoor waste dumps</p>
Application method(s)	<p>Method: bait application</p> <p>Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations Covered and protected baiting points as long as they provide the same level of protection for non target species and human as temper resistant bait stations Direct applications of ready to use bait into the burrows</p>
Application rate(s) and frequency	<p>Application rate: rat 200 g, mice 50 g</p> <p>Dilution (%): 0</p> <p>Number and timing of application:</p> <p>Mice Bait products: - 50 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. In case of high infestation level, applications sites are located 2 meters apart</p> <p>Mice - The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait</p>

	<p>stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p> <p>Rats: Bait products: - 200 g of bait per bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 10 meters. In case of high infestation level, applications sites are located 5 meters apart</p> <p>Rats - The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</p>
Category(ies) of users	trained professional
Pack sizes and packaging material	<p>MURIN PASTA Extreme is furnished in two different type:</p> <ol style="list-style-type: none"> 1. Bait of pasta each containing from 10 to 15g, wrapped into paper food,- 2. Plastic tray each containing 25, 30, 40, 50, 75, 80, 100g <p>Pack size: <i>from 1,5 kg to 25 kg</i></p> <p><i>In case of packaging higher than 10 kg Package is restricted to separately packed bags with a maximum bag size of 5 or 10 kg</i></p> <p>Packaging material:</p> <ul style="list-style-type: none"> - envelope, Plastic: coextruded/ HDPE/polethylene - Can /Tin, Paper, cardboard , - Box, Paper, cardboard with envelope inside coextruded/ HDPE/ polethylene - bucket: HDPE

4.3.1. Use-specific instructions

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

-
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
 - The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
 - Where possible, bait stations must be fixed to the ground or other structures.

- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened

- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

- *Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.*

- Bait should be secured so that it cannot be dragged away from the bait station.

- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

- Wear protective chemical resistant gloves during product handling phase.

- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.

- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

- *When using direct application in burrows:*

- Baits must be placed to minimise the exposure to non-target species and children.

- Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled

- *Remove the remaining product at the end of treatment period*

- Bait should be secured so that it cannot be dragged away from the bait station.

- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

-In case of use in a pulsed baiting system

Replace eaten bait only after 3 days and then at maximum 7 day intervals. Collect any spilled material and dead rodents

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet

-
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "Murin facoum esca grassa or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".
 - Hazardous to wildlife.

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements
- Use of gloves is recommended.

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life 2 years

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

please refer to specific authorised uses

5.2. Risk mitigation measures

please refer to specific authorised uses

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

- Dermal exposure, wash skin with water and then with water and soap.

- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "Murin pasta defend or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".

- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements

- Use of gloves is recommended.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

- Store in places prevented from the access of children, birds, pets and farm animals.

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

- Shelf life 2 years

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.