

# Summary of product characteristics for a biocidal product

**Product name:** Solo Blox

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** TP14-0126

**R4BP 3 asset reference number:** GR-0017673-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
5. General directions for use	14
5.1. Instructions for use	14
5.2. Risk mitigation measures	15
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	16
5.4. Instructions for safe disposal of the product and its packaging	16
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	17
6. Other information	17

## Administrative information

### 1.1. Trade names of the product

Solo Blox
JAGUAR BLOX
Solo Super Size Blox
Jaguar Super Size Blox
Final Blox
Final Super Size Blox

### 1.2. Authorisation holder

<b>Name and address of the authorisation holder</b>	Name	Bell Laboratories Netherlands B.V.
	Address	De Cuserstraat 93 1081 CN Amsterdam Netherlands
<b>Authorisation number</b>	TP14-0126	
<b>R4BP 3 asset reference number</b>	GR-0017673-0000	
<b>Date of the authorisation</b>	25/07/2017	
<b>Expiry date of the authorisation</b>	14/07/2024	

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	Bell Laboratories Inc
<b>Address of the manufacturer</b>	6551 N Towne Rd WI 53598 Windsor United States
<b>Location of manufacturing sites</b>	6551 N Towne Rd WI 53598 Windsor United States

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	11 - Brodifacoum
<b>Name of the manufacturer</b>	Activa
<b>Address of the manufacturer</b>	Via Feltre, 32 20132 Milano Italy
<b>Location of manufacturing sites</b>	Tezza s.r.l. Via Tre Ponti 37050 S. Maria di Zevio Italy 37050 Maria di Zevio Italy

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	Active Substance	56073-10-0	259-980-5	0,005

### 2.2. Type of formulation

RB - Bait (ready for use)

## 3. Hazard and precautionary statements

<b>Hazard statements</b>	May damage the unborn child. May cause damage to organs blood through prolonged or repeated exposure .
<b>Precautionary statements</b>	Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves. IF exposed or concerned: Get medical attention. Store locked up.

Dispose of contents to according to local regulations.

Get medical advice if you feel unwell.

## 4. Authorised use(s)

### 4.1 Use description

#### Use 1 - Use #1 Mice and rats - professional users - indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults  Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults  Scientific name: Rattus rattus Common name: Roof rat Development stage: Juveniles  Scientific name: Mus musculus Common name: House mouse Development stage: Adults  Scientific name: Mus musculus Common name: House mouse Development stage: Juveniles
<b>Field(s) of use</b>	Indoor  Indoor
<b>Application method(s)</b>	Method: Bait application Detailed description:  Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations
<b>Application rate(s) and frequencies</b>	Application Rate: Rats High infestation – up to 60g bait in baiting points every 5 metres. Low infestation – up to 60g bait in baiting points every 10 metres. Mice High infestation – up to 20g bait in baiting points every 2 metres. Low infestation – up to 20g bait in baiting points every 5 metres Dilution (%):

	<p>Number and timing of application:</p> <p>Rats</p> <p>High infestation – up to 60g bait in baiting points every 5 metres</p> <p>Low infestation – up to 60g bait in baiting points every 10 metres</p> <p>Mice</p> <p>High infestation – up to 20g bait in baiting points every 2 metres</p> <p>Low infestation - – up to 20g bait in baiting points every 5 metres</p>
<b>Category(ies) of users</b>	Professional
<b>Pack sizes and packaging material</b>	<p>Minimum pack size of 3 kg.</p> <p>Minimum pack size 3kg: Loose 5g, 10g, 15g, 20g, 28g, 50g, 100g, 150g and 200g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box, chipboard box, or polyethylene plastic pouch with Ziploc seal - 3 to 25kg.</p>

#### 4.1.1 Use-specific instructions for use

<ul style="list-style-type: none"> <li>- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</li> <li>- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.</li> <li>- <i>[When available]</i> Follow any additional instructions provided by the relevant code of best practice.</li> </ul>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

#### 4.1.2 Use-specific risk mitigation measures

<ul style="list-style-type: none"> <li>- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.</li> <li>- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).</li> <li>- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.</li> <li>- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.</li> <li>- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").</li> <li>- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.</li> <li>- Do not wash the bait stations with water between applications.</li> <li>- Dispose dead rodents in accordance with local requirements.</li> </ul>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

**4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

**4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
  
- Shelf life: 24 months

**4.2 Use description**

**Use 2 - Use #2 Mice and rats - trained professional - indoor**

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults  Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles  Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults  Scientific name: Rattus rattus Common name: Roof rat Development stage: Juveniles  Scientific name: Mus musculus Common name: House mouse Development stage: Adults  Scientific name: Mus musculus Common name: House mouse Development stage: Juveniles

<b>Field(s) of use</b>	Indoor  Indoor
<b>Application method(s)</b>	Method: Bait application Detailed description: - Ready-to-use bait to be used in tamper-resistant bait stations - Covered and protected baiting points
<b>Application rate(s) and frequencies</b>	Application Rate: Rats High infestation – up to 60g bait in baiting points every 5 metres. Low infestation – up to 60g bait in baiting points every 10 metres. Mice High infestation – up to 20g bait in baiting points every 2 metres. Low infestation – up to 20g bait in baiting points every 5 metres Dilution (%): Number and timing of application:  Rats  High infestation – up to 60g bait in baiting points every 5 metres  Low infestation – up to 60g bait in baiting points every 10 metres  Mice  High infestation – up to 20g bait in baiting points every 2 metres  Low infestation - – up to 20g bait in baiting points every 5 metres
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	Minimum pack size of 3 kg.  Minimum pack size 3kg: Loose 5g, 10g, 15g, 20g, 28g, 50g, 100g, 150g and 200g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box, chipboard box, or polyethylene plastic pouch with Ziploc seal -  3 to 25kg

#### 4.2.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.



## 4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the applicable code of good practice.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.

## 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

## 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.
- Use of gloves is recommended.

**UK only** : For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

## 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight .
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

## 4.3 Use description

### Use 3 - Use #3 Mice and rats - professional users - outdoor around buildings

#### Product type

PT14 - Rodenticides (Pest control)

#### Where relevant, an exact description of the authorised use

Not relevant for rodenticides

#### Target organism(s) (including development stage)

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Adults

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: Juveniles

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: Adults

Scientific name: Rattus rattus  
Common name: Roof rat  
Development stage: Juveniles

Scientific name: Mus musculus  
Common name: House mouse  
Development stage: Adults

Scientific name: Mus musculus  
Common name: House mouse  
Development stage: Juveniles

**Field(s) of use**

Outdoor  
  
Outdoor

**Application method(s)**

Method: Bait application  
Detailed description:  
  
Bait formulations:  
Ready-to-use bait to be used in tamper-resistant bait stations

**Application rate(s) and frequencies**

Application Rate: Rats High infestation – up to 60g bait in baiting points every 5 metres. Low infestation – up to 60g bait in baiting points every 10 metres. Mice High infestation – up to 20g bait in baiting points every 2 metres. Low infestation – up to 20g bait in baiting points every 5 metres  
Dilution (%):  
Number and timing of application:  
  
Rats  
  
High infestation – up to 60g bait in baiting points every 5 metres  
  
Low infestation – up to 60g bait in baiting points every 10 metres  
  
Mice  
  
High infestation – up to 20g bait in baiting points every 2 metres  
  
Low infestation - – up to 20g bait in baiting points every 5 metres

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

Minimum pack size of 3 kg.  
  
Minimum pack size 3kg: Loose 5g, 10g, 15g, 20g, 28g, 50g, 100g, 150g and 200g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box, chipboard box, or polyethylene plastic pouch with Ziploc seal - 3 to 25kg



#### 4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas non-labile to flooding.
- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- *[When available]* Follow any additional instructions provided by the relevant code of best practice.

#### 4.3.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

#### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface water (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

#### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months.

#### 4.4 Use description

Use 4 - Use #4 Mice and rats - trained professional - outdoor around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults</p> <p>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: Adults</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: Juveniles</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: Adults</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: Juveniles</p>
<b>Field(s) of use</b>	<p>Outdoor</p> <p>Outdoor</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description:</p> <ul style="list-style-type: none"> <li>- Ready-to-use bait to be used in tamper-resistant bait stations.</li> <li>- Covered and protected baiting points</li> </ul>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: Rats High infestation – up to 60g bait in baiting points every 5 metres. Low infestation – up to 60g bait in baiting points every 10 metres. Mice High infestation – up to 20g bait in baiting points every 2 metres. Low infestation – up to 20g bait in baiting points every 5 metres Dilution (%): Number and timing of application:</p> <p>Rats</p> <p>High infestation – up to 60g bait in baiting points every 5 metres</p> <p>Low infestation – up to 60g bait in baiting points every 10 metres</p> <p>Mice</p> <p>High infestation – up to 20g bait in baiting points every 2 metres</p>

	Low infestation - – up to 20g bait in baiting points every 5 metres
<b>Category(ies) of users</b>	Trained professional
<b>Pack sizes and packaging material</b>	<p>Minimum pack size of 3 kg</p> <p>Minimum pack size 3kg: Loose 5g, 10g, 15g, 20g, 28g, 50g, 100g, 150g and 200g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box, chipboard box, or polyethylene plastic pouch with Ziploc seal -</p> <p>3 to 25kg</p>

#### 4.4.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period.
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.

#### 4.4.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

#### 4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.  
- Use of gloves is recommended.  
**UK only** : For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

#### 4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight .  
- Store in places prevented from the access of children, birds, pets and farm animals.  
- Shelf life: 24 months

### 4.5 Use description

#### Use 5 - Use #5 Rats - trained professional - sewers

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Adults  Scientific name: Rattus norvegicus Common name: Brown rat Development stage: Juveniles
<b>Field(s) of use</b>	Other  Sewers
<b>Application method(s)</b>	Method: Bait application Detailed description:  Bait formulations: - Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. - Covered and protected baiting points
<b>Application rate(s) and frequencies</b>	Application Rate: Up to 200g bait per manhole. Dilution (%):

	<p>Number and timing of application:</p> <p>Up to 200g bait per manhole.</p>
<b>Category(ies) of users</b>	<p>Trained professional</p>
<b>Pack sizes and packaging material</b>	<p>Minimum pack size of 2.5 kg</p> <p>Minimum pack size 3kg: Loose 5g, 10g, 15g, 20g, 28g, 50g, 100g, 150g and 200g blocks in High Density Poly Ethylene (HDPE) pail, cardboard box, chipboard box, or polyethylene plastic pouch with Ziploc seal -</p> <p>3 to 25kg</p>

#### 4.5.1 Use-specific instructions for use

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

#### 4.5.2 Use-specific risk mitigation measures

- Do not use this product in pulsed baiting treatments.

#### 4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

- Antidote: Vitamin K1 administered by medical/veterinary personnel only.

- In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Medical professionals for information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-809 2166), retain the label for reference.

Contact a veterinary surgeon in case of ingestion by a pet.

- Hazardous to wildlife.

#### **4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.
- Use of gloves is recommended.

#### **4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight .
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

## **5. General directions for use**

### **5.1. Instructions for use**



### **Professional users**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- *[If national policy or legislation require it]* When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

### **Trained professionals**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

## **5.2. Risk mitigation measures**

#### **Professional users**

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

#### **Professional users with demonstrated competence (equivalent to trained professionals)**

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements

### **5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

#### **Professional users**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
  - Dermal exposure, wash skin with water and then with water and soap.
  - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
  - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.
- Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of accident, call a poison center".
- Hazardous to wildlife.

#### **Professional users with demonstrated competence (equivalent to trained professionals)**

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
  - Dermal exposure, wash skin with water and then with water and soap.
  - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
  - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

### **5.4. Instructions for safe disposal of the product and its packaging**

**Professional users**

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.

**Trained professionals**

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.  
- Use of gloves is recommended.

**5.5. Conditions of storage and shelf-life of the product under normal conditions of storage**

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.  
- Store in places prevented from the access of children, birds, pets and farm animals.  
- Shelf life: 24 months.

**6. Other information**

- Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

- This product contains a bittering agent and a dye.

- Use Biocides Safely and Sustainably

- It is illegal to use this product for uses or in a manner other than that prescribed on this label.

- Poison Information: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.

- Please note IRELAND applies a minimum Professional pack size of 2.5 Kg instead of 3.0 kg.