

Substance Name: Phenanthrene

EC Number: 201-581-5

CAS Number: 85-01-8

MEMBER STATE COMMITTEE

SUPPORT DOCUMENT FOR IDENTIFICATION OF

PHENANTHRENE

**AS A SUBSTANCE OF VERY HIGH CONCERN
BECAUSE OF ITS vPvB¹ (ARTICLE 57E)
PROPERTIES**

Adopted on 12 December 2018

¹ vPvB means very persistent and very bioaccumulative

CONTENT

| | |
|--|-----------|
| FOREWORD | 3 |
| IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57 | 4 |
| JUSTIFICATION | 5 |
| 1. IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES | 5 |
| 1.1 Name and other identifiers of the substance | 5 |
| 1.2 Composition of the substance | 5 |
| 1.3 Physicochemical properties | 6 |
| 2. HARMONISED CLASSIFICATION AND LABELLING | 6 |
| 3. ENVIRONMENTAL FATE PROPERTIES | 6 |
| 3.1 Degradation..... | 6 |
| 3.1.1 Abiotic degradation..... | 6 |
| 3.1.2 Biodegradation..... | 8 |
| 3.1.3 Summary and discussion on degradation..... | 14 |
| 3.2 Environmental distribution..... | 15 |
| 3.2.1 Adsorption/desorption..... | 15 |
| 3.2.2 Volatilisation | 15 |
| 3.2.3 Distribution modelling | 15 |
| 3.2.4 Summary and discussion of environmental distribution | 16 |
| 3.3 Bioaccumulation..... | 16 |
| 3.3.1 Bioaccumulation in aquatic organisms (pelagic & sediment organisms)..... | 16 |
| 3.3.2 Bioaccumulation in terrestrial organisms (soil dwelling organisms, vertebrates)..... | 17 |
| 3.3.3 Field data | 17 |
| 3.3.4 Summary and discussion of bioaccumulation..... | 18 |
| 4. HUMAN HEALTH HAZARD ASSESSMENT | 18 |
| 5. ENVIRONMENTAL HAZARD ASSESSMENT | 18 |
| 6. CONCLUSIONS ON THE SVHC PROPERTIES | 18 |
| 6.1 CMR assessment | 18 |
| 6.2 PBT and vPvB assessment | 19 |
| 6.2.1 Assessment of PBT/vPvB properties | 19 |
| 6.2.2 Summary and overall conclusions on the PBT and vPvB properties..... | 20 |
| 6.3 Assessment under Article 57(f) | 21 |
| REFERENCES | 22 |
| ANNEX I – ENVIRONMENTAL HAZARD ASSESSMENT | 29 |
| 1.1 Aquatic compartment (including sediment)..... | 29 |
| 1.1.1 Fish | 29 |
| 1.1.2 Aquatic invertebrates..... | 30 |
| 1.1.3 Algae and aquatic plants | 30 |
| 1.1.4 Sediment organisms | 31 |
| 1.2 Terrestrial compartment..... | 31 |
| 1.3 Summary and discussion of the environmental hazard assessment..... | 31 |

TABLES

| | |
|---|----|
| Table 1: Substance identity..... | 5 |
| Table 2: Overview of physicochemical properties | 6 |
| Table 3: Fugacity Model calculation (EPI Suite, version 4.11) of phenanthrene | 16 |

FOREWORD

Phenanthrene belongs to the substance group of Polycyclic Aromatic Hydrocarbons (PAHs) of which many are well-known to be hazardous for human health and the environment. Phenanthrene does not possess a harmonised classification according to the CLP Regulation (EC 1272/2008).

Until now, several Annex XV dossiers for the identification of substances of very high concern (SVHC) were explicitly based on the properties of PAHs as constituents of concern in the identified substances, such as Anthracene, Anthracene Oils, Coal Tar Pitch High Temperature (CTPHT), Benzo[a]pyrene, Benz[a]anthracene and Chrysene.

Phenanthrene is constituent, inter alia, in CTPHT. In the Support Document of CTPHT, it has been concluded by the Member State Committee (MSC) that phenanthrene fulfils the vPvB criteria of Annex XIII to the REACH Regulation (ECHA, 2009). However, phenanthrene and further PAHs whose properties have already been agreed on by the MSC in the CTPHT SVHC identification process have not yet been proposed for formal SVHC identification and inclusion in the Candidate List.

Phenanthrene was assessed with respect to vPvB properties based on the MSC Support Document for identification of CTPHT as SVHC (ECHA, 2009). For the purpose of the present SVHC proposal for phenanthrene, a supplementary literature search was made. The search identified only few studies not included in the EU Risk Assessment Report on CTPHT (European Commission, 2008) and the Annex XV Transitional Dossier on CTPHT (The Netherlands, 2008). Thus, the assessment of the vPvB properties in the present dossier and the conclusion that phenanthrene fulfils the criteria in Article 57 (e) was based mainly on the information in the MSC Support Document on CTPHT (ECHA, 2009) and supplemented with information from newer studies that are presented as further evidence as they do not trigger a need to modify the conclusions taken by authorities earlier on.

IDENTIFICATION OF A SUBSTANCE OF VERY HIGH CONCERN ON THE BASIS OF THE CRITERIA SET OUT IN REACH ARTICLE 57

Substance Name: Phenanthrene

EC Number: 201-581-5

CAS number: 85-01-8

- The substance is identified as very persistent and very bioaccumulative (vPvB) according to Article 57 (e) of Regulation (EC) No 1907/2006 (REACH).

Summary of how the substance meets the criteria set out in Article 57 of the REACH Regulation

The assessment of the vPvB properties in the present dossier and the conclusion that phenanthrene fulfils the criteria in Article 57 (e) were based mainly on the information in the MSC Support Document on CTPHT (ECHA, 2009) and supplemented with information from newer studies. All available information (such as the results of standard tests, modelling and (Q)SAR results) were considered together in a weight-of-evidence approach.

Persistence

The available experimental information show that phenanthrene degrades very slowly in sediment with half-life greater than 180 d.

In soil simulation test performed in laboratory condition, phenanthrene shows a fast degradation. However, the available field studies show that phenanthrene could degrade slower depending on e.g. the local conditions, matrix, methodological conditions. The current information does not allow for a final conclusion on persistency of phenanthrene in soil.

Thus, the P and the vP criteria of REACH Annex XIII are fulfilled by phenanthrene for sediment.

Bioaccumulation

Phenanthrene fulfils the B criterion when the bioconcentration factor in aquatic species is greater than 2 000, and the vB criterion when the bioconcentration factor in aquatic species is greater than 5 000. The bioaccumulation of phenanthrene was measured in three studies with fish (BCFs ranging from 2 229 to 6 118 L/kg), two studies with crustacean (BCFs ranging from 5 513 to 28 145 L/kg), two studies with copepod (BCFs ranging from 5 252 to 71 077 L/kg) and one study with an oligochaete species (BCF = 5 222 L/kg). Thus, BCFs greater than 2 000 and 5 000 were obtained.

Thus, the B and the vB criteria of REACH Annex XIII are fulfilled by phenanthrene.

Overall conclusion:

In conclusion, phenanthrene meets the criteria for a vPvB substance according to Article 57 (e) of REACH Regulation by comparing all relevant and available information according to the criteria set out in the Annex XIII of REACH in a weight-of-evidence determination.

Registration dossiers submitted for the substance: No

Justification


1. Identity of the substance and physical and chemical properties

1.1 Name and other identifiers of the substance

Table 1: Substance identity

| | |
|---|---|
| EC number: | 201-581-5 |
| EC name: | Phenanthrene |
| CAS number (in the EC inventory): | 85-01-8 |
| CAS number: Deleted CAS numbers: | - |
| CAS name: | Phenanthrene |
| IUPAC name: | Phenanthrene |
| Index number in Annex VI of the CLP Regulation | No harmonised classification |
| Molecular formula: | C ₁₄ H ₁₀ |
| Molecular weight range: | 178.229 g.mol ⁻¹ |
| Synonyms: | <i>o</i> -diphenyleneethylene, phenanthren, phenanthrin |

Structural formula:



1.2 Composition of the substance

Name: Phenanthrene

Description: Phenanthrene belongs to the group of Polycyclic Aromatic Hydrocarbons (PAHs). Phenanthrene is not produced as such. However, it may occur as a UVCB² constituent in UVCB-substances that are derived from coal or in petroleum streams. The dossier addresses the substance phenanthrene as a substance itself.

Substance type: mono-constituent

² Substances of Unknown or Variable composition, Complex reaction products or Biological materials

1.3 Physicochemical properties

Table 2: Overview of physicochemical properties

| Property | Value [Unit] | Reference/source of information |
|--|-------------------------------------|---------------------------------|
| Physical state at 20°C and 101.3 kPa | solid | INERIS, 2010 |
| Melting/freezing point | 100.5 °C | ECHA, 2009 |
| Boiling point | 340°C | ECHA, 2009 |
| Vapour pressure | 2.6 x 10 ⁻² Pa at 25°C | ECHA, 2009 |
| Density | 0.980 Kg.L ⁻¹ at 20°C | ECHA, 2009 |
| Water solubility | 0.95 mg.L ⁻¹ at 24°C | ECHA, 2009 |
| Partition coefficient n-octanol/water (Log value) | 4.57 at 25°C | ECHA, 2009 |
| Henry's constant | 3.7 Pa m ³ /mol at 25 °C | ECHA, 2009 |

2. Harmonised classification and labelling

No harmonised classification for phenanthrene.

3. Environmental fate properties

3.1 Degradation

The data provided on the degradation of phenanthrene in the Support Document for identification of CTPHT as SVHC (ECHA, 2009) are not assessed or discussed again in this dossier but included in order to have access to the full dataset for assessment (flagged by *italic print*). Additional information is available in the EU risk assessment report on CTPHT (EC, 2008) and the Annex XV Transitional Dossier for CTPHT (The Netherlands, 2008). When more recent data were identified, they have been included in the document and cited accordingly.

3.1.1 Abiotic degradation

3.1.1.1 Hydrolysis

As assessed before in the Support Document for identification of CTPHT as SVHC (ECHA, 2009), phenanthrene does not present functional groups that result in hydrolysis and is therefore expected to be *hydrolytically stable in aquatic systems*. The Support Document furthermore states as a result that *hydrolysis is not expected to contribute to the degradation of PAHs under environmental conditions*.

3.1.1.2 Oxidation

The oxidation of PAHs was assessed in the Annex XV Transitional Dossier for CTPHT (the

Netherlands, 2008), and summarised and discussed in the Support Document for identification of CTPHT as SVHC (ECHA, 2009) as follows:

In the atmosphere, the PAHs are either gas phase or particle-associated. It has been shown that the 2-4 ring PAHs with vapour pressure higher than or equal to 10^{-4} Pa are mostly gas phase-related and PAHs of 4 rings or more with vapour pressure below 10^{-4} Pa are particle-associated. In the gas phase PAHs are oxidised by atmospheric hydroxyl (OH) and nitrate radicals and ozone, whereas the particle-associated PAHs are expected to be degraded by direct photolysis and by reaction with ozone (The Netherlands, 2008). Phenanthrene has 3 aromatic rings and a reported vapour pressure of 2.6×10^{-2} Pa at 25°C, which is higher than 10^{-4} Pa. Therefore, it is assumed that phenanthrene is mainly in the gas phase and is degraded mainly by oxidisation and, to a lesser extent, by photolysis. In the gas phase, phenanthrene can undergo oxidation in the presence of OH radicals exhibiting short lifetimes between a few hours to less than two days (ECHA, 2009).

In the atmosphere, phenanthrene can be partitioned in the gas phase and also adsorbed to the particle phase. Among the semi-volatile PAHs, phenanthrene is mainly in the gas phase but a small fraction (up to 12.4%) can be found adsorbed onto particles which reduces the degradation rate due to stabilisation of PAHs (ECHA, 2009). In the atmosphere, phenanthrene reacts with OH radicals and undergoes “alkene-like” reactions with NO_3 or O_3 radicals (Atkinson, 1994). Nevertheless, “alkene-like” reactions are highly variable because NO_3 radical concentrations are very fluctuant due to the fact that their formation requires the presence of both O_3 and NO_2 . Furthermore, the NO_3 radical reaction is only a night-time loss process because the NO_3 radical rapidly photolyses. The “alkene-like” reaction that phenanthrene can undergo also contributes to its oxidation in the atmosphere.

3.1.1.3 Phototransformation/photolysis

3.1.1.3.1 Phototransformation in air

Photolysis of PAHs in the atmosphere was already assessed in the EU risk assessment report (2008) and summarised in the Support Document for identification of CTPHT as SVHC (ECHA, 2009) and in the Annex XV Transitional Dossier for CTPHT (The Netherlands, 2008) as follows:

Photolysis in the troposphere results in the formation of reactive hydroxyl (OH) and nitrate (NO_3) radicals and ozone (O_3), which reacts as oxidizing agent with organic compounds, like PAHs. These radical and ozone reactions comprise mainly degradation of gas-phase PAHs (Calvert et al., 2002). Particle-associated PAHs are expected to degrade in air predominantly via direct photolysis by light with a wavelength < 290 nm (Kamens et al., 1988), although reaction with ozone will also occur (Peters and Seifert, 1980; Grosjean et al., 1983; Pitts et al., 1986; Coutant et al., 1988).

As stated earlier, phenanthrene in the air is mainly in the gas phase and undergoes oxidisation. The small amount of phenanthrene that is particle-associated undergoes photolysis, with a reaction rate dependent on the type of particle.

3.1.1.3.2 Photo-transformation in water

As assessed before in the Support Document for identification of CTPHT as SVHC (ECHA, 2009), *photo-degradation in natural waters takes normally place only in the upper few centimetres of the water-column and is therefore not considered to have significant impact on the overall persistency of PAHs in the aquatic environment.*

Due to the number of factors that affect photo-degradation rates, this process is not generally considered in the persistence assessment for substances registered under

REACH. Further discussion on photo-degradation is provided in Chapter R.7b of the Guidance on IR&CSA.

According to Castro-Jiménez and de Meent (2011), light absorption in natural water is significantly slower than measured in laboratory water with photo degradation occurring around 30 times more slowly for typical fresh water, 400 times more slowly for typical coastal sea water, and 500 times more slowly for ocean water. These authors also conclude that the "contribution of photo-degradation in water to overall degradation is significant only for substances that reside in water to a considerable extent". They highlight that many substances reside in sediment and soil, rather than in water.

Therefore, aquatic photo-degradation is not considered to have a significant impact on the overall persistency of phenanthrene in the environment.

3.1.1.3.3 Photo-transformation in soil

As assessed before in the Support Document for identification of CTPHT as SVHC (ECHA, 2009), *as exposure to light is even more limited in soils, photo-degradation is as well not considered a relevant degradation process in terrestrial environments.* Thus, photo-degradation is not considered a relevant degradation process in soils.

3.1.1.4 Summary on abiotic degradation

It is concluded that in the atmosphere, free PAHs degrade within a range of minutes to days by direct photolysis. The action of an oxidant on phenanthrene is also an important path for its degradation. A very small part of phenanthrene (12.4%) may be particle-associated and when adsorbed onto fine particles, phenanthrene may be more stable in the atmosphere. In water, phenanthrene is not hydrolysed but can be photo-degraded. However, this only appears at the upper few centimetres of a water-column and is therefore not considered having a significant impact on the overall persistence of phenanthrene in the aquatic environment. In soil, exposure to light is even more limited. Thus, photo-degradation is not considered as relevant degradation process in water and terrestrial environments. Phenanthrene is hydrolytically stable under environmental conditions.

This conclusion was already drawn in the Support Document for identification of CTPHT as SVHC (ECHA, 2009).

3.1.2 Biodegradation

3.1.2.1 Biodegradation in water and sediments

Regarding the biodegradation in water, as assessed in the Support Document for identification of CTPHT as SVHC (ECHA, 2009):

Experimental information for biodegradation in water has demonstrated that PAH substances with up to four aromatic rings are biodegradable under aerobic conditions, but that biodegradation rates of PAHs with more than four aromatic rings, are very low (The Netherlands, 2008).

In general, the biodegradation rates decrease with increasing number of aromatic rings. This correlation has been attributed to factors like the bacterial uptake rate and the bioavailability. The bacterial uptake rate has been shown to be lower for the higher molecular weight PAHs as compared to the PAHs of lower molecular weight. This may be due to the size of high molecular weight members, which limits their ability to cross cellular membranes. In addition, bioavailability is lower for higher molecular PAHs due to adsorption to organic matter in water and sediment. It has further been shown that half-

lives of PAHs in estuarine sediment are proportionally related to the octanol-water partition coefficient (Kow) (Durant et al, (1995) cited in The Netherlands, 2008). [...]

The results from standard tests for biodegradation in water show that PAHs with up to four aromatic rings are biodegradable under aerobic conditions but that the biodegradation rate of PAH with more aromatic rings is very low (ECHA, 2009). Regarding biodegradation on sediments, although there is evidence for anaerobic transformation, PAHs are usually considered to be persistent under anaerobic conditions (Neff, 1979; Volkering & Breure, 2003, cited in The Netherlands, 2008), thus a low biodegradation of phenanthrene is expected in sediments.

Although the biodegradation pathway of the different PAHs is very similar, their biodegradation rates differ considerably. In general, the biodegradation rates decreases with increasing number of aromatic rings. Biodegradation rates also are extremely dependent on the (a)biotic conditions, both in the lab and in the field. Important influencing factors are (1) the substrate concentration; with low PAH concentrations leading to longer half-lives; (2) temperature, which reversely relates to the half-live and (3) the presence or absence of a lag phase (De Maagd, 1996). In addition, the desorption rate of PAHs appears to decrease with increase of the residence time of PAHs due to slow sorption onto micropores and organic matter, and polymerisation or covalent binding to the organic fraction. The consequence of this aging process is a decreased biodegradability and a decreased toxicity (Volkering and Breure, 2003).

As assessed in the Support Document for identification of CTPHT as SVHC (ECHA, 2009) *Mackay et al. (1992) estimated half-lives in the different environmental compartments based on model calculations and literature research. The calculated half-lives of phenanthrene in water and sediments are in the range of 12 to 42 days and longer than 420 to 1250 days respectively.*

In a 28 day ready biodegradability test (MITI I, OECD 301C) using 100 mg/L PAHs and 30 mg/L sludge, phenanthrene did not fulfil the criteria to be considered as readily biodegradable (54% degradation after 4 weeks based on BOD measurement), similarly to fluorene, carbazole, acenaphthene and dibenzofuran. According to the MITI test, which is suitable for substances with low water solubility, these PAHs are not considered as readily biodegradable (CITI, 1992, INERIS, 2010).

Contrarily to this result, in a ready biodegradability test performed according to the OECD 301C (MITI) guideline methodology, using 100 mg/L PAHs and 30 mg/L sludge, phenanthrene achieved 67.2% mineralisation (BOD/ThOD) over 28 days (Junker et al., 2016).

As stated in the R.11 ECHA guidance, "Available data consisting solely of screening information can be employed to derive a conclusion mainly for "not P and not vP" or "may fulfil the P or vP criteria". After the latter conclusion on screening, higher tier information generally needs to be made available. Appropriate data need to be available to conclude the P/vP-assessment with a conclusion "not P/vP" on all three compartments (or five, with marine compartments): water (marine water), sediment (marine sediment) and soil. Either the available data, including in normal case simulation test data from one or two compartments, can be interpreted so that a conclusion can be derived on the remaining compartment(s) for which no higher tier data are available, or data need to be available directly on all compartments, or there is another justification for why a conclusion does not need to be drawn for all three (five) compartments. Moreover, "the conclusion should be based on a Weight-of-Evidence consideration by expert judgement where all relevant and available data for all endpoints are considered in conjunction". As data were available for other compartments than water, these data need to be taken into account in a Weight-of-Evidence consideration for the persistency assessment of phenanthrene.

During the public consultation, references of seven studies related to the degradation of phenanthrene and diesel in water were provided (Birch et al., 2018, Brakstad et al., 2018a, 2018b, Loftus et al., 2018, Ribicic et al., 2018, Prince et al., 2008, Prince et al., 2013).

The studies have been assessed for their reliability and relevance according to OECD and ECHA guidelines (R11 and R7b). Their suitability for the P assessment is limited as they present methodological limitations, as for example the addition of mineral media in the experiments, the origin of the samples that could be considered as pre-adapted to PAHs, the absence of abiotic control and reference substance, the use of dispersant in the experiments, the absence of information about the dissolved phenanthrene in the experiments, or the use of inappropriate internal markers for the half-life determination.

During the public consultation, a poster³ was provided presenting half-lives in surface water measured at temperatures ranging from 5 – 22°C. Although the authors seems to suggest that the half-life is not depending on temperature, such conclusion cannot be drawn without further information of the test conditions at which the half-lives are measured to allow an appropriate comparison. Therefore no conclusion on the temperature dependency of the degradation of phenanthrene from this poster can be drawn (Concawe, 2018).

In addition, the results of one poster⁴ (Hammershoj et al., 2018) related to the degradation of mixtures was transmitted. Phenanthrene was not part of the mixture, the information have not been further considered in the current assessment.

In a microcosm study (Bahr et al., 2015), the biodegradation of four PAHs (naphthalene, fluorene, phenanthrene, and acenaphthene added as ¹³C-labelled substrates) was investigated as single substances in groundwater (pH around 7) from an aquifer located at the site of a former gas plant with oxic conditions. ¹³CO₂-values of controls amended with non-labelled phenanthrene shifted from -20‰ to -27‰, while the sterile controls remained stable. There is no information on the effect of pre-adaptation on biodegradation in this study. ¹³C-enrichment of the produced CO₂ revealed mineralisation of phenanthrene comprised between 14.2% and 33.1% over a period of 62 days of incubation in the dark at 14°C, in order to approximate field conditions. This study used a BACTRAP® system (composed of activated carbon pellets) that allowed trapping a bacterial community already present in the aquifer by the system for 100 days. In this experiment, the percentage of dissolved oxygen was always recorded to ensure a minimum of 1% content in order to maintain an oxic condition representative of the *in-situ* conditions. The use of an adapted bacterial community may lead to an overestimation of the mineralisation rate in comparison to the situation arising in other aquifer. This study highlighted that phenanthrene degradation was slow in oxic aquifer, indicating possible persistence. Biodegradation was observed in the presence of an adapted bacterial community (100 days of incubation plus 62 days of exposure to reach 33% of mineralisation).

An extended summary of a water-sediment simulation OECD 308 study was provided during the public consultation focusing on phenanthrene (Meisterjahn et al. 2018a). The tests were conducted in closed biometer systems due to significant observed loss of test material in flow-through setups during pre-tests. All other parameters were followed as expressed in the guideline. The reported half-lives for the total system estimated by CAKE calculation ranged between 114 to 150 days depending on the considered sediment and statistical assessment (one sediment had fine texture and high organic carbon content; the other had a coarse texture and low organic carbon content). The DegT₅₀ for the first sediment range from 114 to 130 days and from 116 to 150 days for the second sediment. These data are below the cut-off value for P criteria. Nevertheless, the ModelMaker calculation for compartment specific (Water and sediment phase) provide half-lives (DT₅₀) for the first sediment of 56 days for water and 305 days for sediment. For the second sediment, the ModelMaker calculation provides DT₅₀ of 172 days for water and 116 days for sediment. These model calculations indicate, at the opposite of the other DegT₅₀

³ See it ("*Persistence assessment of phenanthrene: a case study*") in the Response-to-comment document (RCOM) on the SVHC proposal for phenanthrene, in the embedded attachment of comment 5257 submitted by Concawe.

⁴ See it ("*Mixture effects on biodegradation kinetics of petroleum hydrocarbons in surface water*") in the Response-to-comment document (RCOM) on the SVHC proposal for phenanthrene, in the embedded attachment of comment 5257 submitted by Concawe.

determine by the CAKE calculation, that the phenanthrene is vP for sediment. Moreover, the experiments were performed at 20°C. However, for a regulatory purpose, the recommended temperature is 12°C, as recommended in the R7-Guidance on information requirements and chemical safety assessment. When recalculations were done with the simplified Arrhenius equation recommended by ECHA, the DegT50 were all above the cut-off value of 180 days, comprised between 216 to 247 days for the first sediment and comprised between 220 to 285 days for the second sediment. If the methodology used the non-simplified Arrhenius equation, the DegT50 were between 242 to 276 days for the first sediment and between 246 to 319 days for the second sediment. Thus, data indicate that a low biodegradation of phenanthrene is observed in sediments, based on which phenanthrene meets the vP criteria for sediment.

3.1.2.2 Biodegradation in soil

Biodegradation in soil was assessed in the Support Document for identification of CTPHT as SVHC (ECHA, 2009) as follows:

Biodegradation rates of PAHs in soil depend on several factors related to the soil type, including pH, moisture content, nutrients, oxygen, and the diversity of the soil microbial population. Various species (bacteria, fungi, yeasts and algae) are known to degrade PAHs in soil (The Netherlands, 2008). It has been shown that the number of PAH-degrading microorganisms and the degradation capacity is higher in PAH-contaminated soils than in pristine soils, something explained by the development of an adapted soil microbial community. Several studies have also been demonstrated enhanced PAH degradation rates when the soil had been enriched with isolated PAH-degrading microorganisms ([...] The Netherlands, 2008). On the basis of a comparison between two studies (Wild et al. 1991 and Wild and Jones, 1993) it was illustrated that the half-lives observed under laboratory conditions can be much shorter than those obtained from long-term field studies. This was attributed by the authors to the more optimal conditions (temperature, moisture content, nutrient and oxygen supply) applied in the laboratory tests.

Furthermore the Support Document for identification of CTPHT as SVHC (ECHA, 2009) discusses "aging" for PAHs as follows:

"Aging" is a phenomenon associated with increased residence time of PAHs in soil, which can further decrease the bioavailability of PAHs in the terrestrial environment. Freshly spiked PAHs are more readily desorbed and thus more bioavailable than PAHs that have been in soil or sediment for a longer period of time (The Netherlands, 2008). This means that studies involving artificially added PAHs (e.g. ¹⁴C-labelled) often result in biodegradation rates much higher than rates observed for the same substances present in soil as part of a contamination by coal tar."

Regarding the information provided in the studies by Wild and Jones (1993) and Wild et al. (1991), it should be noted that the MSC has already considered their use in the assessment of CTPHT as SVHC, and hence also for phenanthrene. In agreement with MSC conclusions made within the CTPHT assessment, the SVHC dossier submitter evaluated the study Wild et al. (1991) as a most reliable evidence of persistency of CTPHT (thus also of phenanthrene), which is suitable for PBT assessment of pristine environment.

Wild and Jones (1993) and Wild et al. (1991) studied the biodegradation of PAHs in soil amended with sewage sludge under laboratory and field conditions, respectively. They determined dissipation half-lives for phenanthrene of 83 to 193 days (corresponding to a half-life up to 643 days when converted to 12°C by using the Arrhenius equation) in laboratory soil microcosms (three soil types: sandy loams, forest soil, and roadside soil conducted at a range of temperature between 20 and 30°C) and, under field conditions, a half-life of 5.7 years. When phenanthrene was spiked to soil, the dissipation half-life was lower, i.e. 16 days. However, as the abiotic loss, due to e.g. volatilization, was high (65%), the reliability of this as a degradation half-life is questionable.

Mackay et al. (1992) as summarised by The Netherlands (2008) suggested a half-life for

phenanthrene in the range of 125 to 420 days in soil. These predictions are in line with half-lives observed by the field studies mentioned above.

Mineralisation of ^{14}C -phenanthrene was investigated with the indigenous microbial communities present in five pristine soils of Livingstone Island, one of the South Shetland Islands separated from the Antarctica Peninsula by the Bransfield Strait. The temperature rarely exceeded 3°C in summer and decreased to -11°C in winter. Mineralisation of ^{14}C -phenanthrene was investigated using respirometric assay at different temperature (4° , 12° , 22°C) and under slurry condition (22°C + Mineral Basal Salt (MBS) medium) in the dark for 35 days. Briefly, 10g of soils were rehydrated to 40-60% water holding capacity and spiked with unlabelled and labelled ^{14}C -phenanthrene. The latter condition (slurry) was not considered relevant for the present assessment due to the adjunction of MBS who improved the bacterial growth, dispersion, accessibility and enhanced phenanthrene bioavailability. The maximum extent of mineralisation of ^{14}C -phenanthrene was 1.14% at 4°C for 35 days, indicating that, in natural temperature conditions of Livingstone Island, phenanthrene mineralisation is extremely slow compared to higher temperatures. At 12°C , maximum mineralisation extent was 35.15 % and 39.09% at 22°C indicating that phenanthrene degradation was limited, which was consistent with the plateau in degradation reached in two of the studied soils (soil 4 and 5). At 4 and 12°C , degradation did not start for 35 days of incubation which was reflected by lag phases (time to reach 5% mineralisation). 5% mineralisation was not reached in the 5 soils at 4°C and 3 soils at 12°C and even 2 soils at 22°C (Okere *et al.*, 2012).

In 2017, the same research group evaluated the biodegradation of phenanthrene in the same Antarctic soils over 150 days and at various temperatures. Briefly, 50g of soils were spiked with ^{12}C -phenanthrene and then mixed with 200g of non-spiked soils to avoid adverse effects to bacterial community. Then soils were left incubated in the dark at 4° , 12° and 22°C for 1, 30, 60 and 150 days. These soils were used to determine the catabolic activity of indigenous bacterial community exposed to ^{14}C -phenanthrene using respirometric assay at different temperature (4° , 12° , 22°C) and under slurry condition (22°C + Mineral Basal Salt (MBS) medium). As detailed before, the latter condition was not considered. In the set-up, 10g of soils were rehydrated to 40-60% water holding capacity and spiked with ^{12}C -phenanthrene (>99.6%) and ^{14}C -phenanthrene. The respirometers were stored in the dark, incubated at temperatures set for pre-exposition of the soils for 21 days. The lag phase for bacterial growth (time to reach 5% mineralisation) was never reach for the soils incubated at 4°C , was reach in 4/20 (2 for 60 days of contact and 2 for 150 days of contact) at 12°C and in 6/20 at 22°C (3 for 60 days of contact and 3 for 150 days of contact). At 4°C , the maximum extent of mineralisation of ^{14}C -phenanthrene was 0.46% for 1 day contact soil. At 12°C , the maximum extent of mineralisation of ^{14}C -phenanthrene was 12.21% for 60 day contact soil and was greater than 1% in only 5/20 conditions (2 for 60 days of contact and 3 for 150 days of contact). At 22°C , the maximum extent of mineralisation of ^{14}C -phenanthrene was 24.82% and was greater than 5% in only 6/20 conditions (3 for 60 days of contact and 3 for 150 days of contact). This study results highlighted the fact that in soil with poor organic matter content, adsorption of phenanthrene will occur on mineral surfaces, reducing its bioavailability. When increasing temperature and exposure duration, the mineralisation of phenanthrene increases. Nevertheless, it can be observed from these results that in natural conditions that can be found on Livingstone Island, no biodegradation of phenanthrene was observed, indicating that phenanthrene will remain over a long period of time in the environment. Moreover, when the contact time between phenanthrene and soils was increased, the maximum amounts of phenanthrene available for biodegradation was decreased. This was reflected by the plateau of 7% of mineralisation reached at 12°C after 150 days of pre-exposure indicating that soil could accumulate phenanthrene due to ageing (increase in contact time) (Okere *et al.*, 2017).

A microcosm study evaluated a phenanthrene-contaminated Patagonian soil (Pico Truncado, Patagonia, Argentina), maintained under arid conditions and incubated at $20 \pm 2^\circ\text{C}$ for 250 days. There was no abiotic control for this experiment. No biodegradation of

phenanthrene measured as primary degradation was observed for 150 even with the adjunction of fertilisers at day 86 of treatment (microcosms fertilised with 7.45 g kg⁻¹ dry soil of commercial fertiliser Nitrofoska® (BASF, Research Triangle Park, NC), taking the relation C/N/P to 100:5:2). The lack of biodegradation was observed with and without bioaugmentation (adjunction of *Sphingobium sp.* strain 22B, a PAH-degrading strain), indicating that phenanthrene is recalcitrant to biodegradation under arid conditions. After day 150, the water content of soil was increased from 10 % to 15%, drastically increasing the biodegradation of phenanthrene, from no degradation of phenanthrene (day 150) to 98% degradation of the added phenanthrene at the end of the experiment (day 210) (Madueño, Alvarez, and Morelli, 2015).

Bioavailability of the substance to microorganisms constitutes a crucial factor in biodegradation. However, the main criterion to decide on the persistence character of a substance in soil/sediment is the half-life in relevant conditions.

During public consultation, an extended summary on a soil simulation test according to OECD 307 was provided (Meisterjahn et al. 2018b). The study met all the quality criteria except a slight over-recovery of radioactivity in one case (mainly 111.4 % normally limited to 110 %). In the four soils studied, the resulted degradation half-lives ranged from 6.8 to 17.3 days (geometric mean: 11.1 days) when considering only the extractable fraction. However, for a regulatory purpose, the recommended temperature is 12°C, as recommended in the R7- Guidance on information requirements and chemical safety assessment. The half-lives then ranged from 13 to 33 days. The mass balance resulted in a high percentage of NER (comprised between 42- 51%) for the samples tested, whereas under sterile conditions, the formation of NER was much lower, which suggest most of the NER cannot be considered as parent substance. This study indicates that degradation of phenanthrene can occur in soil.

During the public consultation, a reference (Sigmund et al, 2018) regarding explicitly the biodegradation of phenanthrene on spiked soils was transmitted. This study has been assessed for its reliability and relevance according to OECD and ECHA guidelines (R11 and R7b). It is noted that the study Sigmund et al, 2018 is not comparable to the study of Wild and Jones, 1993 due to methodological differences.

The experiment objective was to investigate the effects of compost amendment, both with and without biochar on PAHs and NSO-PAH degradation in soils. The soil samples were spiked with 25 mg of phenanthrene/kg of soil, and additionally spiked with 100 mg/kg of Zn and 10 mg/kg of Cd, heavy metals. This may cause changes in bacterial community and their capacity to biodegrade PAHs. The study was conducted for 120 days with non-radiolabelled phenanthrene, no information about the recovery of the substance (formation of NER) is available in the publication. Furthermore, the study did not perform an abiotic control. The data presented in the publication for the half-life is limited to a graphical representation making the determination of the real half-life value difficult. It seems that the half-life presented in this graphical representation is comparable to the half-life determined in the OECD 307 test. Due to methodological limitations, the reliability of this half-life cannot be assessed.

Harmsen and Rietra, 2018, reference provided during the public consultation, performed a long-term study on soil and sediment focusing on biodegradation of PAHs and total petroleum hydrocarbons (TPH), which has been monitored on seven experimental fields during periods up to 25 years. The study took place on experimental fields at Kreekraksluizen, situated in the Netherlands. Landfarms were initiated on semi-field scale in 1990. About 50 cm of dredged sediment was applied and the layer thickness of the dewatered sediments was about 30 cm. The sediments were intensively treated to stimulate biodegradation during the first years. Treatments used were cultivation, no cultivation, adding wood chips and adding sludge from a sewage system adapted to PAHs. In 1993 the sediment from the different experimental fields were combined into two new experimental fields containing sediments that originated from two harbors in a thicker layer. These sediments were further treated in a passive way (passive landfarming), only

allowing vegetation to grow to create aerobic conditions and were followed for 20 years. The bioavailable fraction was measured using Tenax extraction. The study detailed the remaining concentration of every PAHs measured and also phenanthrene after 7.2 years and 22 years. The initial concentration of phenanthrene was 168 mg/kg, after 7.2 years, the remaining amount was estimated being 3 mg/kg and after 22 years, estimated of being 1.3 mg/kg. In other sediment from harbour in Wemeldinge were initial concentration of phenanthrene was 9 mg/kg, after 7.2 years it remains 2.5 mg/kg and after 22 years 0.8 mg/kg. It is observed that the degradation rate of phenanthrene may depend on the initial concentration and that a certain fraction can remain for a long period of time in the environment (Harmsen and Rietra, 2018).

3.1.2.3 Summary and discussion on biodegradation

Regarding water and sediment, Mackay et al. (1992) indicated that predicted phenanthrene elimination half-lives ranged between 13 and 42 days and that the substance persisted in sediment with half-lives between 420 to 1250 days. Regarding the available information from the ready biodegradation tests showing not ready biodegradation for one result (Junker et al., 2016), and ready biodegradation from the other (INERIS, 2010), biodegradation of phenanthrene may occur in water. An extended summary of a water-sediment simulation OECD 308 study was provided during the public consultation (Meisterjahn et al., 2018a). The DegT50 for the first sediment ranged from 114 to 130 days and from 116 to 150 days for the second sediment. These data are below the cut-off value for P criteria. When recalculations were done at 12°C DegT50 were all above the cut-off value of 180 days, comprised between 216 to 319 days. Thus, data indicate that a low biodegradation of phenanthrene is observed in sediments, based on which phenanthrene meets the vP criteria for sediment.

For soil, Mackay et al. (1992) indicated that phenanthrene persisted with half-lives between 125 to 420 days. These half-lives are comparable to those observed in field conditions when phenanthrene is introduced via sludge, ranging between 83 to 193 days (corresponding to a half-life up to 643 days when converted to 12°C by using the Arrhenius equation) (Wild and Jones, 1993). It was demonstrated that under even more harsh conditions biodegradation of phenanthrene can go even slower. In contrast to the high half-life observed under field conditions, in a standard soil simulation test according to OECD 307, a much lower degradation half-life of phenanthrene were observed, ranging from 6.8 to 17.3 days at 20°C, (corresponding to 13 to 33 days at 12°C). This study indicates that degradation of phenanthrene can occur in soil.

A study of Harmsen and Rietra, 2018 suggests that the degradation rate of phenanthrene may depend on the initial concentration and that a certain fraction can remain for a long period of time in the environment.

In a soil simulation test performed in laboratory conditions, phenanthrene shows a fast degradation. However, the available field studies show that phenanthrene could degrade slower depending on e.g. the local conditions, matrix, methodological conditions. The current information does not allow for a final conclusion on persistency of phenanthrene in soil.

3.1.3 Summary and discussion on degradation

In the atmosphere, phenanthrene can be partitioned in the gas phase and, to a fewer extent, also be adsorbed to particulates. Degradation in the gas phase by oxidation in the presence of OH radicals takes place between a few hours to less than two days.

In the water and soil compartments, photolysis is only relevant in the upper few centimeters of the water column and the upper few millimeters of the soil. Thus, photodegradation is not considered as a relevant degradation process in water and terrestrial environments.

In general, PAHs have no functional groups that result in hydrolysis in the water and soil compartments. Therefore, phenanthrene is considered as hydrolytically stable.

The predicted half-lives range between 13 to 42 days for degradation in water and between 420 to 1250 days for sediment. In the sediment compartment, a low biodegradation of phenanthrene is observed. The low degradation rate is confirmed in a sediment simulation study according to OECD TG 308, provided during the public consultation (Meisterjahn et al. 2018a), based on which phenanthrene meets the vP criteria for sediment. In a standard soil simulation test according to OECD 307, much lower degradation half-lives of phenanthrene were observed, ranging from 6.8 to 17.3 days at 20°C, (corresponding to 13 to 33 days at 12°C). This study indicates that degradation of phenanthrene can occur in soil. Wild and Jones (1993) reported dissipation half-lives for phenanthrene of 83 to 193 days (corresponding to a half-life up to 643 days when converted to 12°C by using the Arrhenius equation) in a laboratory soil microcosm study when applied via sewage sludge. A study of Harmsen and Rietra, 2018, suggeststhat the degradation rate of phenanthrene may depend on the initial concentration and that a certain fraction can remain for a long period of time in the environment.

Under field conditions, Wild *et al.* (1991) demonstrated a half-life of more than 5.7 years in soil for phenanthrene.

In a soil simulation test performed in laboratory condition, phenanthrene shows a fast degradation. However, the available field studies show that phenanthrene could degrade slower depending on e.g. the local conditions, matrix, methodological conditions. The current information does not allow for a final conclusion on persistency of phenanthrene in soil.

3.2 Environmental distribution

3.2.1 Adsorption/desorption

In the Support Document for identification of CTPHT as SVHC (ECHA, 2009) the adsorption properties of PAHs are described as follows: *“A linear relationship between K_{ow} and the organic carbon-water partitioning coefficient K_{oc} has been demonstrated for PAHs in sediments and soil. The Log K_{ow} values from 4.6 to 6.6 can be translated as a high potential for partitioning to soils and sediments. Partitioning processes like adsorption to airborne particulate matter, as well as accumulation in sludge during wastewater treatment, have been demonstrated especially for high molecular weight PAHs (The Netherlands, 2008).”*

Based on the Log K_{ow} of 4.57 for phenanthrene, reported in the Annex XV Transitional Dossier on CTPHT (The Netherlands, 2008), the coefficient of partitioning between organic carbon and water, Log K_{oc} , has been estimated at 4.36 (The Netherlands, 2008) and it is concluded that phenanthrene has a high potential to adsorb to particles in the environment.

3.2.2 Volatilisation

Phenanthrene has a reported vapour pressure of 2.6×10^{-2} Pa at 25°C and a Henry’s law constant of 3.7 Pa m³/mol at 25°C (Mackay et al., 2006, cited by The Netherlands, 2008). Thus, some volatilisation is expected from water or soil surfaces.

3.2.3 Distribution modelling

Mackay Level III fugacity modelling was done using EPI Suite (version 4.11) with default values of environmental emission rates (it is assumed that phenanthrene is released at equal rates to air, water, and soil) (computations done in April 2018). The calculations

revealed a distribution of phenanthrene almost exclusively to soil and a small part to sediment and water phases (Table 3).

Table 3: Fugacity Model calculation (EPI Suite, version 4.11) of phenanthrene

| Distribution to: | Mass amount (percent) |
|------------------|-----------------------|
| Air | 0.484 |
| Water | 9.75 |
| Soil | 77.3 |
| Sediment | 12.4 |

3.2.4 Summary and discussion of environmental distribution

Phenanthrene exhibits a high potential to adsorb to organic matter and some volatilisation from soil and water is expected. Furthermore, according to the fugacity model, phenanthrene is expected to be mainly distributed in the soil compartment followed by the sediment and water compartment. The air compartment is not expected to be a relevant route of distribution.

3.3 Bioaccumulation

3.3.1 Bioaccumulation in aquatic organisms (pelagic & sediment organisms)

Experimental studies were available to determine the BCF of phenanthrene in different organisms. They were considered in the Support Document for the identification of CTPHT as SVHC (ECHA, 2009) and reassessed later by the RIVM (Bleeker and Verbruggen, 2009) and completed with a literature search (April 2018). When bioaccumulation data have been reassessed by the RIVM, these are used in preference to the CTPHT data. When more recent supporting data was identified, they were included in the current document and cited accordingly.

BCF values were reported above 2 000 L/kg for sheepshead minnows (*Cyprinodon variegatus* exposed to phenanthrene during 36 days in continuous flow system followed by 8 days of depuration providing a BCF of 2 229 L/kg) and fathead minnows (*Pimephales promelas*; BCF_K: 3 611 L/kg) (Carlson et al., 1979, Jonsson et al., 2004). The BCF value obtained with fathead minnows (*Pimephales promelas*; BCF_K: 3 611 L/kg) were reassessed and normalised with a 5% lipid content, providing a BCF value of 1149 L/kg for sheepshead minnows and 4 751 L/kg for fathead minnows close to the 5 000 L/kg threshold for vB criteria. For fathead minnows, the study realised by De Maagd (De Maagd, 1996) was used in the European Union Risk Assessment Report on CTPHT (EC, 2008) and provided a BCF value of 6 760 L/kg. However, after reassessing this study, it was concluded that this study cannot any longer be rated as reliability index of 2 but 3 (namely, unreliable) due to uncertainties in the determination method. It was stated by the RIVM that the results from De Maagd (1996) should be considered with care, but are still important as circumstantial evidence in a weight of evidence approach to decide on the bioaccumulation potential of the studies PAHs (Verbruggen and Herwijnen, 2011).

During public consultation, more recent studies (8 studies: Niimi et al., 1986, Cheikyula et al., 2008, Lo et al., 2016, Kobayashi et al., 2013, Ke et al., 2007, Xia et al., 2015, Wang et al., 2018, Li et al., 2018) conducted with fish were transmitted. Reported BCF values vary between the fish species, ranging from 76 L/kg for Japanese flounder (Cheikyula et al., 2008) to 1954 L/Kg for the benthic fish *Pseudopleuronectes yokohamae* (Kobayashi et al, 2013). It is noted that the reliability of these studies has not been evaluated in detail by the dossier submitter. Nevertheless, the study by Cheikyula et al., 2008, Niimi et al.,

1986 were rated as R 3 and 4 by Verbruggen and Herwijnen, 2011. Overall, these data do not undermine the data already available for fish.

The study of Carlson et al., 1979, providing a kinetic BCF of 4751 L/kg after lipid normalisation also provides a final equilibrium static BCF value of 5 100 L/kg considered as reliable with restriction (Bleeker and Verbruggen, 2009). Moreover, when lipid normalization was applied to the different static BCFs available in this study, it provides a maximum BCF value of 6118 considered as reliable with restriction (The Netherlands, 2018). In this study, *Fathead minnows* were exposed to a series of PAHs present in lake water, via flow through conditions for 28 days followed by 5 days of depuration. In molluscs phenanthrene accumulates, but BCF levels remain below 2 000 L/kg (maximum: 1 280 L/kg). In crustaceans, two studies provided very high BCF values. Landrum et al., 2003, conducted a static renewal experiment of toxicity and bioaccumulation with the amphipod *Diporeia spp.* exposed to a range of concentrations of ¹⁴C PAHs (from 57.1 to 637.8 µg/L of phenanthrene) for 28 days. The authors determined BCF values for phenanthrene by kinetic approach ranging from 5 513 to 11 440 L/kg (8 889 L/kg) for the tested concentrations. The second one was determined in flow-through systems by the kinetic method with the amphipod *Pontoporeia hoyi* exposed to selected ¹⁴C radiolabelled PAHs during 6 hours followed by 14 days of depuration phase. This study provided a BCF value of 28 145 L/kg. The two studies were considered as reliable with restriction and reliable without restriction respectively.

In oligochaete *Stylodrilus heringianus*, a flow-through experiment using the kinetic method rated as reliable with restriction provided a BCF value of 5 222 L/kg (Frank et al., 1986, as cited in Bleeker and Verbruggen, 2009).

Regarding *copepods*, a laboratory experiment in *Calanus finmarchicus* conducted a BCF determination in a 192 h semi-static daily renewal exposure with ¹⁴C-labeled phenanthrene followed by a 96 h depuration time. Phenanthrene accumulated rapidly, reaching steady state within 96 h and providing, at the end of depuration step, a lipid normalised BCF (with the kinetic method) of 5 252 L/kg (Jensen et al., 2012). In a more recent study, BCF determination with the kinetic methodology on arctic copepod species *Calanus hyperboreus* for 4 days of uptake and 3 days depuration provided BCF values ranging from 40 330 to 71 077 L/kg depending on the copepod developmental stage. (Agersted et al., 2018).

For insects (*Hexagenia limbata*), BCFs clearly depend on lipid content of the organism, and may reach values as high as 5 697 L/kg.

3.3.2 Bioaccumulation in terrestrial organisms (soil dwelling organisms, vertebrates)

The European Union Risk Assessment Report on CTPHT (EC, 2008) estimated BCF values on earthworm for several PAHs according to the equilibrium partitioning approach (EP). For phenanthrene, it was estimated a BCF earthworm of 450 L/kg. This value represents a reasonable worst case (EC, 2008).

3.3.3 Field data

Trophic magnification studies are available for phenanthrene. In the study of Wan et al., 2007, not mentioned in the CTPHT Support Document, PAHs concentrations in phytoplankton/seston, zooplankton, invertebrates, fish and one seabird species collected from Bohai Bay on the north of China were analysed. The trophic magnification factor (TMF) calculated for phenanthrene was 0.43. Another study of Nfon et al., 2008, calculated a TMF of 0.82 (0.73–0.92) for phenanthrene, referring to a study conducted in a benthic and pelagic food chain from the Baltic Sea. The species of the food chain analysed included pelagic species of phytoplankton, zooplankton and several benthic species. A study

investigated the bioaccumulation and biomagnification of PAHs by analyzing 11 finfish species and the blue crab, *Callinectes sapidus*, collected from the fresh-brackish portion of the Passaic River. For phenanthrene, a TMF of 0.34 was calculated (Khairy, Weinstein, and R. Lohmann *et al.*, 2014).

The study conducted by Wang (Wang *et al.*, 2012) reported estimated TMF values for several PAHs, based on the analysis of several fish species with different feeding behaviours (herbivorous, omnivorous and carnivorous) collected on Taihu Lake in China. A TMF of 1.27 was calculated for phenanthrene, suggesting biomagnification of the chemical through the food chain.

It is mentioned in the chapter R.11 of the ECHA guidance on PBT or vPvB that the results from field studies should be considered as part of the overall evaluation of the bioaccumulation properties of a substance. However, currently there is no consensus about standard methodologies and guidelines for the interpretation of such results, generating uncertainties for the interpretation of those results. In particular, temporal and spatial variability or the inclusion/exclusion of a few or even a single species could affect the outcome of the TMFs. Thus, the data treatment could have a deep impact on the TMF values that were calculated. Ultimately, chapter R.11 of the ECHA guidance on PBT or vPvB indicates that the absence of a biomagnification potential cannot be used on its own to conclude that the B or vB criteria are not fulfilled.

3.3.4 Summary and discussion of bioaccumulation

The bioaccumulation potential of phenanthrene differs between the organisms due to their capacity to metabolise (biotransform) PAHs. It is likely that phenanthrene is transformed in fish using enzymes belonging to the Cytochrome P450 enzymes (Cyt P450) and other mechanisms, resulting in low BCF values. However, invertebrate species may have a lower metabolic capacity than fish species, e.g. as is the case for polycyclic aromatic hydrocarbons (Bleeker and Verbruggen, 2009). Bioaccumulation in these invertebrates may therefore be higher than in fish under the same exposure conditions and this situation should be considered in a Weight-of-Evidence approach.

The bioaccumulation of phenanthrene was measured in fish (BCFs of 2 229 to 6 118 L/kg), crustacean (BCFs of 5 513 to 28145 L/kg), copepod (BCFs of 5 252 to 71 077 L/kg) and oligochaete (BCF of 5 222 L/kg). Thus, the BCFs values were higher than 2000 and 5000.

4. Human health hazard assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 point (e) of REACH.

5. Environmental hazard assessment

Not relevant for the identification of the substance as SVHC in accordance with Article 57 point (e) of REACH. Information related to the T criterion of Article 57 (d) of REACH is presented in Annex I as additional information.

6. Conclusions on the SVHC Properties

6.1 CMR assessment

This section is not relevant for the identification of the substance phenanthrene as SVHC in accordance with Article 57 (e).

6.2 PBT and vPvB assessment

6.2.1 Assessment of PBT/vPvB properties

An assessment of the vPvB properties of phenanthrene has already been carried out by the MSC in the context of the identification of CTPHT as SVHC, as documented in the MSC Support Document for identification of CTPHT as SVHC (ECHA, 2009). Additional information was assessed earlier in the EU Risk Assessment Report on CTPHT (European Commission, 2008) and the Annex XV Transitional Dossier on CTPHT (The Netherlands, 2008) and was further supplemented with information from newer studies.

6.2.1.1 Persistence

Phenanthrene has low water solubility and shows a high tendency to adsorb to particles and organic matter in the environment. The resulting low bioavailability is one of the limiting factors of its biodegradation.

Regarding water and sediment, Mackay *et al.* (1992) indicated that predicted phenanthrene elimination half-lives range between 13 and 42 days and it persists in sediment with half-lives comprised between 420 to 1250 days. Regarding the available information from the ready biodegradation test showing ready biodegradation for one result (INERIS, 2010) and no ready biodegradation from a test system based on the OECD 301 C (MITI) (Junker *et al.*, 2016), biodegradation of phenanthrene may occur in water. An extended summary of a water-sediment simulation OECD 308 study was provided during the public consultation (Meisterjahn *et al.* 2018a). The DegT50 for the first sediment range from 114 to 130 days and from 116 to 150 days for the second sediment. These data are below the cut-off value for P criteria. However, for a regulatory purpose, the recommended temperature is 12°C. When recalculations were done at 12°C with the simplified Arrhenius equation recommended by ECHA, the DegT50 were all above the cut-off value of 180 days, comprised between 216 to 247 days for the first sediment and comprised between 220 to 285 days for the second sediment. If the methodology used the non-simplified Arrhenius equation, the DegT50 were between 242 to 276 days for the first sediment and between 246 to 319 days for the second sediment. Thus, data indicate that a low biodegradation of phenanthrene is observed in sediments, based on which phenanthrene meets the vP criteria for sediment.

For soil, Mackay *et al.* (1992) indicated that phenanthrene persisted with half-lives between 125 to 420 days. These half-lives are comparable to those observed in field conditions when phenanthrene is introduced via sludge, ranging between 83 to 193 days (corresponding to a half-life up to 643 days when converted to 12°C by using the Arrhenius equation) (Wild and Jones, 1993). It was demonstrated that under even more harsh conditions biodegradation of phenanthrene can go even slower. In contrast to the high half-life observed under field conditions, in a standard soil simulation test according to OECD 307, a much lower degradation half-life of phenanthrene were observed, ranging from 6.8 to 17.3 days at 20°C, (corresponding to 13 to 33 days at 12°C). This study indicate that degradation of phenanthrene can occur in soil.

A study of Harmsen and Rietra 2018, suggests that the degradation rate of phenanthrene may depend on the initial concentration and that a certain fraction can remain for a long period of time in the environment. Finally, biodegradation studies on soil done by Wild *et al.* (1991) demonstrated a phenanthrene half-life of more than 5.7 years under field conditions.

In soil simulation test performed in laboratory condition, phenanthrene show a fast degradation. However, the available field studies shows that phenanthrene could degrade slower depending on e.g. the local conditions, matrix, methodological conditions. The current information does not allow for a final conclusion on persistency of phenanthrene in soil. The data from the OECD 307 cannot be used, based on our current knowledge, to

discredit the available data from field studies.

Overall it is concluded that phenanthrene fulfils the P and vP criteria for sediment according to REACH Annex XIII.

6.2.1.2 Bioaccumulation

Experimentally obtained BCF values > 2 000 and 5 000 are reported with phenanthrene in fish (2 229 to 6 118 L/kg), crustacean (5 513 to 28 145 L/kg), copepod (5 252 to 71 077 L/kg) and oligochaete (5 222 L/kg). In accordance to REACH Annex XIII, phenanthrene fulfils the B and vB criteria. This conclusion was already drawn by the MSC in the context of the identification of CTPHT as SVHC (ECHA, 2009). New data retrieved in the literature search done in April 2018 have been included in the current report and did not challenge the previous conclusion.

6.2.2 Summary and overall conclusions on the PBT and vPvB properties

The assessment of the PvB properties in the present dossier and the conclusion that phenanthrene fulfils the criteria in Article 57 (e) were based mainly on the information in the MSC Support Document on CTPHT (ECHA, 2009) and supplemented with information from newer studies. All available information (such as the results of standard tests, modelling and (Q)SAR results) were considered together in a weight-of-evidence approach.

Persistence

The available experimental information show that phenanthrene degrades very slowly in sediment with half-life greater than 180 days.

In soil simulation test performed in laboratory condition, phenanthrene shows a fast degradation. However, the available field studies show that phenanthrene could degrade slower depending on e.g. the local conditions, matrix, methodological conditions. The current information does not allow for a final conclusion on persistency of phenanthrene in soil.

Thus, the P and the vP criteria of REACH Annex XIII are fulfilled by phenanthrene for sediments.

Bioaccumulation

Phenanthrene fulfils the B criterion when the bioconcentration factor in aquatic species is greater than 2 000, and the vB criterion when the bioconcentration factor in aquatic species is greater than 5 000. The bioaccumulation of phenanthrene was measured in three studies with fish (BCFs ranging from 2 229 to 6 118 L/kg), two studies with crustacean (BCFs ranging from 5 513 to 28 145 L/kg), two studies with copepod (BCFs ranging from 5 252 to 71 077 L/kg) and one study with an oligochaete species (BCF = 5 222 L/kg). Thus, BCFs greater than 2 000 and 5 000 were obtained.

Thus, the B and the vB criteria of REACH Annex XIII are fulfilled by phenanthrene.

Overall conclusion:

In conclusion, phenanthrene meets the criteria for a vPvB substance according to Article 57 (e) of REACH Regulation by comparing all relevant and available information according to the criteria set out in the Annex XIII of REACH in a weight-of-evidence determination.

6.3 Assessment under Article 57(f)

This section is not relevant for the identification of phenanthrene as SVHC in accordance with Article 57 (e) of REACH.

References

- Afnor. 2009. "Bio indicateur de la toxicité potentielle de milieu aqueux." XP-T90-382. Afnor, La Plaine-Saint-Denis, France, pp 19.
- Agersted, M. D., E. F. Møller, and K. Gustavson. 2018. "Bioaccumulation of oil compounds in the high-Arctic copepod *Calanus hyperboreus*." *Aquatic Toxicology* 195:8-14. doi: 10.1016/j.aquatox.2017.12.001.
- Alves, R. N., C. F. Mariz, Jr., D. V. Paulo, and P. S. Carvalho. 2017. "Toxicity of effluents from gasoline stations oil-water separators to early life stages of zebrafish *Danio rerio*." *Chemosphere* 178:224-230. doi: 10.1016/j.chemosphere.2017.03.023.
- Amorim, M. J., E. Oliveira, A. S. Teixeira, C. S. Gravato, S. Loureiro, L. C. Guilhermino, C. A. Van Gestel, and A. M. Soares. 2011. "Toxicity and bioaccumulation of phenanthrene in *Enchytraeus albidus* (Oligochaeta: Enchytraeidae)." *Environmental Toxicology and Chemistry* 30 (4):967-972. doi: 10.1002/etc.464.
- Atkinson, R. 1994. "Gas-phase tropospheric chemistry of organic compounds." *Journal of Physical and Chemical Reference Data*, Monograph, 2, 1-216
- Bahr, A., A. Fischer, C. Vogt, and P. Bombach. 2015. "Evidence of polycyclic aromatic hydrocarbon biodegradation in a contaminated aquifer by combined application of in situ and laboratory microcosms using ¹³C-labelled target compounds." *Water Research* 69:100-109. doi: 10.1016/j.watres.2014.10.045.
- BAUA. (2017) Analysis of the most appropriate risk management option (RMOA). Fluoranthene, Pyrene, Benz(a)anthracene, Chrysene, Benzo(k)fluoranthene, Benzo(ghi)perylene, Phenanthrene. Authority: Germany. Date: June 2017
- Behymer, T. D., and R. A. Hites. 1988. "Photolysis of polycyclic aromatic hydrocarbons adsorbed on fly ash." *Environmental Science & Technology* 22 (11):1311-1319. doi: 10.1021/es00176a011.
- Birch, H., Hammershøj, R., Mayer, P., 2018. Determining Biodegradation Kinetics of Hydrocarbons at Low Concentrations: Covering 5 and 9 Orders of Magnitude of K_{ow} and K_{aw}. *Environmental Science & Technology* 52, 2143-2151.
- Bisson, M., Dujardin R., Flammarion P., Garric J., Babut M., Lamy M.-H., Porcher J.-M., Thybaud É., and Vindimian É., 2000. "Complément au SEQ-Eau: méthode de détermination des seuils de qualité pour les substances génotoxiques." Verneuil-en-Halatte, France: Institut National de l'Environnement Industriel et des Risques (INERIS), Agence de l'eau Rhin-Meuse.
- Black, J. A., W. J. Birge, A. G. Westerman, and P. C. Francis. 1983. "Comparative aquatic toxicology of aromatic hydrocarbons." *Fundamental and Applied Toxicology* 3 (5):353-358. doi: [https://doi.org/10.1016/S0272-0590\(83\)80004-9](https://doi.org/10.1016/S0272-0590(83)80004-9).
- Bleeker E.A.J., Wiegman S., Droge S.T.J., Kraak M.H.S., and Van Gestel C.A.M. 2003. "Towards an improvement of the risk assessment of polycyclic (hetero)aromatic hydrocarbons." Amsterdam, The Netherlands: Aquatic Ecology and Ecotoxicology, Faculty of Science, University of Amsterdam and Animal Ecology, Institute of Ecological Science, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam. Report 2003-01 (UvA)/Report 2003-04 (VU).
- Bleeker E.A.J., and Verbruggen E.M.J. 2009. "Bioaccumulation of polycyclic aromatic hydrocarbons in aquatic organisms." Bilthoven: RIVM. Report no. 601779002.
- Boese, B. L., J. O. Lamberson, R. C. Swartz, R. Ozretich, and F. Cole. 1998. "Photoinduced toxicity of PAHs and alkylated PAHs to a marine infaunal amphipod." *Arch Environ Contam Toxicol* 34 (3):235-40.
- Brakstad, O.G., Nordtug, T., Throne-Holst, M., 2015. "Biodegradation of dispersed Macondo oil in seawater at low temperature and different oil droplet sizes." *Marine Pollution Bulletin* 93, 144-152.

- Brakstad, O.G., Ribicic, D., Winkler, A., Netzera, R., 2018a. "Biodegradation of dispersed oil in seawater is not inhibited by a commercial oil spill dispersant." *Marine Pollution Bulletin* 129, 555-561.
- Brakstad et al., 2018b. "Dispersibility and biotransformation of oils with different properties in seawater." *Chemosphere* 191, 44-53.
- Bruner, K. A., S. W. Fisher, and P. F. Landrum. 1994. "The Role of the Zebra Mussel, *Dreissena polymorpha*, in Contaminant Cycling: I. The Effect of Body Size and Lipid Content on the Bioconcentration of PCBs and PAHs." *Journal of Great Lakes Research* 20 (4):725-734. doi: [https://doi.org/10.1016/S0380-1330\(94\)71190-4](https://doi.org/10.1016/S0380-1330(94)71190-4).
- Butler, J. D., T. F. Parkerton, D. J. Letinski, G. E. Bragin, M. A. Lampi, and K. R. Cooper. 2013. "A novel passive dosing system for determining the toxicity of phenanthrene to early life stages of zebrafish." *Science of the Total Environment* 463-464:952-958. doi: 10.1016/j.scitotenv.2013.06.079.
- Carlson R.M., Oyler A.R., Gerhart E.H., Caple R., Welch K.J., Kopperman H.L., Bodenner D. and Swanson D.1979. "Implications to the aquatic environment of polynuclear aromatic hydrocarbons liberated from Northern Great Plains coal." US EPA report EPA-600/3-79-093, U.S. Environmental Protection Agency (US EPA), Duluth, MN.
- Calvert, J.G., R. Atkinson, K.H. Becker, R.M. Kamens, J.H. Seinfeld, T.H. Wallington, and G. Yarwood. 2002. *The Mechanisms of Atmospheric Oxidation of the Aromatic Hydrocarbons*: Oxford University Press.
- Cheikyula, J.O., J. Koyama, and S. Uno, 2008. "Comparative study of bioconcentration and EROD activity induction in the Japanese flounder, red sea bream, and Java medaka exposed to polycyclic aromatic hydrocarbons." *Environmental Toxicology: An International Journal*. 23(3): p. 354-362.
- CITI-Japan, 1992, Biodegradation and bioaccumulation data of existing chemicals based on CSCL Japan. Chemicals Inspection and Testing Institute
- Concawe, 2018. "Persistence assessment of phenanthrene: a case study." Poster. Concawe Ecology Group,
- Coutant, R. W., L. Brown, J. C. Chuang, R. M. Riggin, and R. G. Lewis. 1988. "Phase distribution and artifact formation in ambient air sampling for polynuclear aromatic hydrocarbons." *Atmospheric Environment (1967)* 22 (2):403-409. doi: [https://doi.org/10.1016/0004-6981\(88\)90046-7](https://doi.org/10.1016/0004-6981(88)90046-7).
- De Maagd PG-J. 1996. "Polycyclic aromatic hydrocarbons: fate and effects in aquatic environment." Utrecht, Utrecht University.
- Droge, S. T.J., M. L. Paumen, E. A.J. Bleeker, M. H.S. Kraak, and C. A.M. van Gestel. 2006. "Chronic toxicity of polycyclic aromatic compounds to the springtail *Folsomia candida* and the enchytraeid *Enchytraeus crypticus*." *Environmental Toxicology and Chemistry* 25 (9):2423-2431. doi: doi:10.1897/05-628R.1.
- Durant, N. D., L. P. Wilson, and E. J. Bouwer. 1995. "Microcosm studies of subsurface PAH-degrading bacteria from a former manufactured gas plant." *Journal of Contaminant Hydrology* 17 (3):213-237. doi: [https://doi.org/10.1016/0169-7722\(94\)00034-F](https://doi.org/10.1016/0169-7722(94)00034-F).
- EC. 2008. "European Union Risk Assessment Report, Coal Tar Pitch High Temperature, CAS No: 65996-93-2, EINECS No: 266-028-2."
- ECHA. 2009. "Support Document for identification of Coal Tar Pitch, High Temperature as a SVHC because of its PBT and CMR properties." <http://echa.europa.eu/documents/10162/73d246d4-8c2a-4150-b656-c15948bf0e77>
- Evans, A. D., and M. Nipper. 2007. "Toxicity of phenanthrene and lindane mixtures to marine invertebrates." *Environ Toxicol* 22 (5):495-501. doi: 10.1002/tox.20279.

- Finch, B. E., S. Marzocchi, D. M. Di Toro, and W. A. Stubblefield. 2017. "Evaluation of the phototoxicity of unsubstituted and alkylated polycyclic aromatic hydrocarbons to mysid shrimp (*Americamysis bahia*): Validation of predictive models." *Environmental Toxicology and Chemistry* 36 (8):2043-2049. doi: doi:10.1002/etc.3733.
- Finnish Environment Institute 2018. Open information. Ympäristötiedon hallintajärjestelmä Hertta. Kertymärekiesteri (KERTY). Available at <https://wwwp2.ymparisto.fi/scripts/kirjaudu.asp> (registration required) (in Finnish)
- Frank, A. P., P. F. Landrum, and B. J. Eadie. 1986. "Polycyclic aromatic hydrocarbon rates of uptake, depuration, and biotransformation by Lake Michigan *Stylodrilus heringianus*." *Chemosphere* 15 (3):317-330. doi: [https://doi.org/10.1016/0045-6535\(86\)90025-1](https://doi.org/10.1016/0045-6535(86)90025-1).
- Grosjean, D., K. Fung, and J. Harrison. 1983. "Interactions of polycyclic aromatic hydrocarbons with atmospheric pollutants." *Environmental Science & Technology* 17 (11):673-679. doi: 10.1021/es00117a010.
- Halling-Sørensen, B., N. Nyholm, Kresten O. Kusk, and E. Jacobsson. 2000. "Influence of Nitrogen Status on the Bioconcentration of Hydrophobic Organic Compounds to *Selenastrum capricornutum*." *Ecotoxicology and Environmental Safety* 45 (1):33-42. doi: <https://doi.org/10.1006/eesa.1999.1818>.
- Hammershøj, R., H. Birch, P. Mayer. 2017. "Mixture effects on biodegradation kinetics of petroleum hydrocarbons in surface water." Poster Presentation. SETAC Europe 27th Annual Meeting, Brussels. Society of Environmental Toxicology and Chemistry.
- Harmesen, J. & R.P.J.J. Rietra, 2018. "25 years monitoring of PAHs and petroleum hydrocarbons biodegradation in soil." *Chemosphere* 207, 229–238.
- Hodson, P. V. 2017. "The Toxicity to Fish Embryos of PAH in Crude and Refined Oils." *Archives of Environmental Contamination and Toxicology* 73 (1):12-18. doi: 10.1007/s00244-016-0357-6.
- Holst L., and Giesy J. P. 1989. "Chronic effects of the photoenhanced toxicity of anthracene on *Daphnia magna* reproduction." *Environmental Toxicology and Chemistry* 8 (10):933-942. doi: doi:10.1002/etc.5620081012.
- INERIS .2010. "Fiche de données toxicologiques et environnementales des substances chimiques – PHENANTHRENE. "
- Jensen, L. K., J. O. Honkanen, Jæ, I. ger, and J. Carroll. 2012. "Bioaccumulation of phenanthrene and benzo[a]pyrene in *Calanus finmarchicus*." *Ecotoxicology and Environmental Safety* 78:225-231. doi: 10.1016/j.ecoenv.2011.11.029.
- Jonsson, G., R. K. Bechmann, S. D. Bamber, and T. Baussant. 2004. "Bioconcentration, biotransformation, and elimination of polycyclic aromatic hydrocarbons in sheepshead minnows (*Cyprinodon variegatus*) exposed to contaminated seawater." *Environ Toxicol Chem* 23 (6):1538-48.
- Junker T., A. Coors, G.Schüürmann, 2016. "Development and application of screening tools for biodegradation in water–sediment systems and soil." *Science of The Total Environment* 544, 1020-1030.
- Kamens, R. M., Z. Guo, J. N. Fulcher, and D. A. Bell. 1988. "The influence of humidity, sunlight, and temperature on the daytime decay of polyaromatic hydrocarbons on atmospheric soot particles." *Environmental Science & Technology* 22 (1):103-108. doi: 10.1021/es00166a012.
- Ke, R., J. Li, M. Qiao, Y. Xu and Z. Wang, 2007. "Using semipermeable membrane devices, bioassays, and chemical analysis for evaluation of bioavailable polycyclic aromatic hydrocarbons in water." *Archives of environmental contamination and toxicology*. 53(3): p. 313-320.
- KEMI .2003. "KEMI Report No 5/03, HA oils in automotive tyres – prospects of a national ban." Report on a government commission, Stockholm, EN, https://www.kemi.se/Documents/Publikationer/Trycksaker/Rapporter/Rapport5_03.pdf

- Khairy, M. A., M. P. Weinstein, and R. Lohmann. 2014. "Trophodynamic Behavior of Hydrophobic Organic Contaminants in the Aquatic Food Web of a Tidal River." *Environmental Science & Technology* 48 (21):12533-12542. doi: 10.1021/es502886n.
- Kobayashi, J., et al., 2013. "Respiratory uptake kinetics of neutral hydrophobic organic chemicals in a marine benthic fish, *Pseudopleuronectes yokohamae*." *Chemosphere*. 93(8): p. 1479-1486.
- Lampi, M. A., J. Gurska, K. I. McDonald, F. Xie, X. D. Huang, D. G. Dixon, and B. M. Greenberg. 2006. "Photoinduced toxicity of polycyclic aromatic hydrocarbons to *Daphnia magna*: ultraviolet-mediated effects and the toxicity of polycyclic aromatic hydrocarbon photoproducts." *Environ Toxicol Chem* 25 (4):1079-87.
- Landrum, P. F., G. R. Lotufo, D. C. Gossiaux, M. L. Gedeon, and J. H. Lee. 2003. "Bioaccumulation and critical body residue of PAHs in the amphipod, *Diporeia* spp: additional evidence to support toxicity additivity for PAH mixtures." *Chemosphere* 51 (6):481-9. doi: 10.1016/s0045-6535(02)00863-9.
- Landrum, P.F. 1988. "Toxicokinetics of organic xenobiotics in the amphipod, *Pontoporeia hoyi*: Role of physiological and environmental variables." *Aquatic Toxicology* 12 (3):245-271.
- Li, Y., et al., 2018. "Dissolved organic matter affects both bioconcentration kinetics and steady-state concentrations of polycyclic aromatic hydrocarbons in zebrafish (*Danio rerio*)." *Science of The Total Environment*. 639: p. 648-656.
- Lo, J.C., et al., 2016. "In Vivo Biotransformation Rates of Organic Chemicals in Fish: Relationship with Bioconcentration and Biomagnification Factors." *Environmental Science & Technology*. 50(24): p. 13299-13308.
- Lofthus et al., 2018. "Biodegradation in seawater of PAH and alkylphenols from produced water of a North Sea platform." *Chemosphere* 206, 465-473.
- Lu, J., S. Tian, X. Lv, Z. Chen, B. Chen, X. Zhu, and Z. Cai. 2018. "TiO₂ nanoparticles in the marine environment: Impact on the toxicity of phenanthrene and Cd²⁺ to marine zooplankton *Artemia salina*." *Science of the Total Environment* 615:375-380. doi: 10.1016/j.scitotenv.2017.09.292.
- Ma, Y., A. Huang, S. Cao, F. Sun, L. Wang, H. Guo, and R. Ji. 2016. "Effects of nanoplastics and microplastics on toxicity, bioaccumulation, and environmental fate of phenanthrene in fresh water." *Environ Pollut* 219:166-173. doi: 10.1016/j.envpol.2016.10.061.
- Mackay, D., W.-Y. Shiu, K.-C. Ma, and S. C. Lee. 2006. *Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals; Volume 2-Halogenated Hydrocarbons*.
- Mackay, D., W. Y. Shiu, and K. C. Ma. 1992. "Illustrated handbook of physical-chemical properties and environmental fate for organic chemicals. Volume II: polynuclear aromatic hydrocarbons, polychlorinated dioxins, and dibenzofurans." *Lewis Publishers, Boca Raton, FL*. 1992. 597.
- Madueño, L., H. M. Alvarez, and I. S. Morelli. 2015. "Autochthonous bioaugmentation to enhance phenanthrene degradation in soil microcosms under arid conditions." *International Journal of Environmental Science and Technology* 12 (7):2317-2326. doi: 10.1007/s13762-014-0637-5.
- McLeese, D. W., and L. E. Burrige. 1987. "Comparative accumulation of PAHs in four marine invertebrates."
- Meisterjahn et al., 2018a (Report in preparation). Degradation of ¹⁴C labelled hydrocarbons in Soil using standard and optimized OECD test guidelines - 307. Germany: Fraunhofer IME-AE.
- Meisterjahn et al., 2018b (Report in preparation). Degradation of ¹⁴C labelled hydrocarbons in Water-Sediment using standard and optimized OECD test guidelines - 308. Germany: Fraunhofer IME-AE.

- Mhadhbi, L., M. Boumaiza, and R. Beiras. 2010. "A standard ecotoxicological bioassay using early life stages of the marine fish *Psetta maxima*." *Aquatic Living Resources* 23 (2):209-216. doi: 10.1051/alr/2010014.
- Mu, J., J. Wang, F. Jin, X. Wang, and H. Hong. 2014. "Comparative embryotoxicity of phenanthrene and alkyl-phenanthrene to marine medaka (*Oryzias melastigma*)." *Marine Pollution Bulletin* 85 (2):505-515. doi: 10.1016/j.marpolbul.2014.01.040.
- Neff, J.M. 1979. "Polycyclic aromatic hydrocarbons in the aquatic environment. Sources, fate and biological effects." Applied Science Publishers, London, 262 pp.
- Niimi, A. and V. Palazzo, 1986. "Biological half-lives of eight polycyclic aromatic hydrocarbons (PAHs) in rainbow trout (*Salmo gairdneri*)." *Water Research*. 20(4): p. 503-507.
- Nfon, E., I. T. Cousins, and D. Broman. 2008. "Biomagnification of organic pollutants in benthic and pelagic marine food chains from the Baltic Sea." *Sci Total Environ* 397 (1-3):190-204. doi: 10.1016/j.scitotenv.2008.02.029.
- Okay, O. S., and B. Karacik. 2007. "Photoinduced toxicity of selected PAHs to the marine microalga *Phaeodactylum tricornutum*." *Journal of Environmental Science and Health, Part A* 42 (6):707-714. doi: 10.1080/10934520701304344.
- Okere, U. V., A. Cabrerizo, J. Dachs, K. C. Jones, and K. T. Semple. 2012. "Biodegradation of phenanthrene by indigenous microorganisms in soils from Livingstone Island, Antarctica." *FEMS Microbiol Lett* 329 (1):69-77. doi: 10.1111/j.1574-6968.2012.02501.x.
- Okere, U. V., A. Cabrerizo, J. Dachs, U. O. Ogbonnaya, K. C. Jones, and K. T. Semple. 2017. "Effects of pre-exposure on the indigenous biodegradation of 14C-phenanthrene in Antarctic soils." *International Biodeterioration and Biodegradation* 125:189-199. doi: 10.1016/j.ibiod.2017.09.013.
- Owojori, O. J., K. Waszak, and J. Roembke. 2014. "Avoidance and reproduction tests with the predatory mite *Hypoaspis aculeifer*: Effects of different chemical substances." *Environmental Toxicology and Chemistry* 33 (1):230-237. doi: 10.1002/etc.2421.
- Peters, J., and B. Seifert. 1980. "Losses of benzo(a)pyrene under the conditions of high-volume sampling." *Atmospheric Environment* (1967) 14 (1):117-119. doi: [https://doi.org/10.1016/0004-6981\(80\)90114-6](https://doi.org/10.1016/0004-6981(80)90114-6).
- Pitts, J. N., H.-R. Paur, B. Zielinska, J. Arey, A. M. Winer, T. Ramdahl, and V. Mejia. 1986. "Factors influencing the reactivity of polycyclic aromatic hydrocarbons adsorbed on filters and ambient POM with ozone." *Chemosphere* 15 (6):675-685. doi: [https://doi.org/10.1016/0045-6535\(86\)90033-0](https://doi.org/10.1016/0045-6535(86)90033-0).
- Ribicic et al., 2018. "Oil type and temperature dependent biodegradation dynamics - Combining chemical and microbial community data through multivariate analysis." *BMC Microbiology* 18:83.
- S.R., Wild, and Jones K.C. 1993. "Biological and abiotic losses of polynuclear aromatic hydrocarbons (PAHs) from soils freshly amended with sewage sludge." *Environmental Toxicology and Chemistry* 12 (1):5-12. doi: doi:10.1002/etc.5620120103.
- Seiler, T. B., N. Best, M. M. Fernqvist, H. Hercht, K. E. C. Smith, T. Braunbeck, P. Mayer, and H. Hollert. 2014. "PAH toxicity at aqueous solubility in the fish embryo test with *Danio rerio* using passive dosing." *Chemosphere* 112:77-84. doi: 10.1016/j.chemosphere.2014.02.064.
- Sigmund G., C. Poyntner, G. Piñar, M. Kah and T. Hofmann. 2018. "Influence of compost and biochar on microbial communities and the sorption/degradation of PAHs and NSO-substituted PAHs in contaminated soils." *Journal of Hazardous Materials* 345, 107–113.
- Smith, K. E., N. Dom, R. Blust, and P. Mayer. 2010. "Controlling and maintaining exposure of hydrophobic organic compounds in aquatic toxicity tests by passive dosing." *Aquat Toxicol* 98 (1):15-24. doi: 10.1016/j.aquatox.2010.01.007.

- Sverdrup, L. E., J. Jensen, P. H. Krogh, and J. Stenersen. 2002. "Studies on the effect of soil aging on the toxicity of pyrene and phenanthrene to a soil-dwelling springtail." *Environmental Toxicology and Chemistry* 21 (3):489-492. doi: doi:10.1002/etc.5620210303.
- Sverdrup, L. E., A. E. Kelley, P. H. Krogh, T. Nielsen, J. Jensen, J. J. Scott-Fordsmand, and J. Stenersen. 2001. "Effects of eight polycyclic aromatic compounds on the survival and reproduction of the springtail *Folsomia fimetaria* L. (Collembola, isotomidae)." *Environ Toxicol Chem* 20 (6):1332-8.
- Sverdrup, L. E., P. H. Krogh, T. Nielsen, and J. Stenersen. 2002. "Relative sensitivity of three terrestrial invertebrate tests to polycyclic aromatic compounds." *Environ Toxicol Chem* 21 (9):1927-33.
- Swartz, R. C., S. P. Ferraro, J. O. Lamberson, F. A. Cole, R. J. Ozretich, B. L. Boese, D. W. Schults, M. Behrenfeld, and G. T. Ankley. 1997. "Photoactivation and toxicity of mixtures of polycyclic aromatic hydrocarbon compounds in marine sediment." *Environmental Toxicology and Chemistry* 16 (10):2151-2157. doi: doi:10.1002/etc.5620161023.
- The Netherlands. 2008. Annex XV Transitional Dossier for substance Coal Tar Pitch, High Temperature, CAS Number: 65996-93-2.
- The Netherlands. 2018. PBT factsheet on three ring PAHs, PBT-EG, october 2018.
- Turcotte, D., P. Akhtar, M. Bowerman, Y. Kiparissis, R. S. Brown, and P. V. Hodson. 2011. "Measuring the toxicity of alkyl-phenanthrenes to early life stages of medaka (*Oryzias latipes*) using partition-controlled delivery." *Environmental Toxicology and Chemistry* 30 (2):487-495. doi: doi:10.1002/etc.404.
- Turja, R., Soirinsuo, A., Budzinski, H., Devier, M. H., Lehtonen, K. K. 2013. Biomarker responses and accumulation of hazardous substances in mussels (*Mytilus trossulus*) transplanted along a pollution gradient close to an oil terminal in the Gulf of Finland (Baltic Sea). *Comparative Biochemistry and Physiology, Part C* 157 (2013) 80-92. <http://dx.doi.org/10.1016/j.cbpc.2012.09.006>
- Turja, R., Höher, N., Snoeijs, P., Baršienė, J., Butrimavičienė, L., Kuznetsova, T., Kholodkevich, S. V., Devier, M.-H., Budzinski, H., Lehtonen, K.K. 2014. A multibiomarker approach to the assessment of pollution impacts in two Baltic Sea coastal areas in Sweden using caged mussels (*Mytilus trossulus*). *Science of the Total Environment* 473-474, 398-409. <https://doi.org/10.1016/j.scitotenv.2013.12.038>
- Verbruggen E.M.J., and van Herwijnen R. 2011. "Environmental risk limits for phenanthrene." RIVM Letter report 601357007/2011.
- Verbruggen, EMJ. 2012. Environmental risk limits for polycyclic aromatic hydrocarbons (PAHs): For direct aquatic, benthic, and terrestrial toxicity. Rijksinstituut voor Volksgezondheid en Milieu RIVM.
- Verrhiest, G., B. Clément, and G. Blake. 2001. "Single and Combined Effects of Sediment-Associated PAHs on Three Species of Freshwater Macroinvertebrates." *Ecotoxicology* 10 (6):363-372. doi: 10.1023/a:1012223014534.
- Vieira, L. R., and L. Guilhermino. 2012. "Multiple stress effects on marine planktonic organisms: Influence of temperature on the toxicity of polycyclic aromatic hydrocarbons to *Tetraselmis chuii*." *Journal of Sea Research* 72:94-98. doi: <https://doi.org/10.1016/j.seares.2012.02.004>.
- Volkering F., and Breure A.M. 2003. "Biodegradation and general aspects of bioavailability." In Douben (ed.) *PAHs: An Ecotoxicological Perspective.*, Wiley, 81- 98.
- Wan, Y., X. Jin, J. Hu, and F. Jin. 2007. "Trophic dilution of polycyclic aromatic hydrocarbons (PAHs) in a marine food web from Bohai Bay, north China." *Environ Sci Technol* 41 (9):3109-14.
- Wang, J., X. Zhang, and G. H. Li. 2012. "Compositions and diagnostic ratios of heavily degraded crude oil residues in contaminated soil in oilfields." *Huanjing Kexue/Environmental Science* 33 (4):1352-1360.

- Wang, H., et al., 2018. "Relationship between metabolic enzyme activities and bioaccumulation kinetics of PAHs in zebrafish (*Danio rerio*)." *Journal of Environmental Sciences*. 65: p. 43-52.
- Wild, S. R., M. L. Berrow, and K. C. Jones. 1991. "The persistence of polynuclear aromatic hydrocarbons (PAHs) in sewage sludge amended agricultural soils." *Environmental Pollution* 72 (2):141-157. doi: [https://doi.org/10.1016/0269-7491\(91\)90064-4](https://doi.org/10.1016/0269-7491(91)90064-4).
- Willis, A. M., and J. T. Oris. 2014. "Acute photo-induced toxicity and toxicokinetics of single compounds and mixtures of polycyclic aromatic hydrocarbons in zebrafish." *Environmental Toxicology and Chemistry* 33 (9):2028-2037. doi: 10.1002/etc.2648.
- Wu, S., H. Zhang, S. Zhao, J. Wang, H. Li, and J. Chen. 2012. "Biomarker responses of earthworms (*Eisenia fetida*) exposed to phenanthrene and pyrene both singly and combined in microcosms." *Chemosphere* 87 (4):285-293. doi: 10.1016/j.chemosphere.2011.11.055.
- Xia, X., et al., 2015. "How does predation affect the bioaccumulation of hydrophobic organic compounds in aquatic organisms?" *Environmental science & technology*. 49(8): p. 4911-4920.
- Zhang, X., X. Xia, J. Dong, Y. Bao, and H. Li. 2014. "Enhancement of toxic effects of phenanthrene to *Daphnia magna* due to the presence of suspended sediment." *Chemosphere* 104:162-9. doi: 10.1016/j.chemosphere.2013.10.086.
- Zhao, Y., X. Wang, X. Lin, S. Zhao, and J. Lin. 2017. "Comparative developmental toxicity of eight typical organic pollutants to red sea bream (*Pagrosomus major*) embryos and larvae." *Environmental Science and Pollution Research* 24 (10):9067-9078. doi: 10.1007/s11356-016-6282-4.
- Zindler, F., B. Glomstad, D. Altin, J. Liu, B. M. Jenssen, and A. M. Booth. 2016. "Phenanthrene Bioavailability and Toxicity to *Daphnia magna* in the Presence of Carbon Nanotubes with Different Physicochemical Properties." *Environ Sci Technol* 50 (22):12446-12454. doi: 10.1021/acs.est.6b03228.

Annex I – Environmental Hazard Assessment

In the current Annex XV report, Phenanthrene is assessed with respect to its vPvB properties based on the MSC Support Document for identification of CTPHT as a SVHC according to the criteria set out in the Annex XIII (ECHA, 2009). As this dossier only focuses on the identification of phenanthrene as a SVHC substance according to article 57 (e), the toxicity data presented here are only informative.

Thus, several environmental toxicity studies have been assessed in the Annex XV Transitional Dossier for CTPHT (The Netherlands, 2008) and summarised in the Support Document for identification of CTPHT as SVHC (ECHA, 2009). As the data presented in the following sections is based on these documents, they will not be assessed and discussed again within this dossier. Additional relevant studies were retrieved in a bibliographic search in April 2018 and have been included in the analysis hereafter.

1.1 Aquatic compartment (including sediment)

The Support Document for the identification of CTPHT as SVHC summarises the following on environmental hazard assessment for the aquatic compartment. *PAHs can be toxic via different modes of action, such as non-polar narcosis and phototoxicity. Phototoxicity is caused by the ability of PAHs to absorb UVA radiation, UVB radiation, and in some instances, visible light. It may occur as the result of the production of singlet oxygen, which is highly damaging to biological material, or as result of the formation of new, more toxic compounds from the photomodification (usually oxidation) of PAHs (Lampi et al., 2006). Phototoxic effects can be observed after a short period of exposure, which explains why for PAHs like anthracene, fluoranthene and pyrene, where phototoxicity is most evident, the acute toxicity values under simulated solar radiation may be lower than the chronic toxicity values determined under less harsh radiation.*

The phototoxicity of PAHs is relevant where the PAHs are exposed to light and UV radiation, and considered to be most important for upper layers of aquatic and terrestrial environments. Although UV penetration depths may vary among PAH-contaminated sites, it is not unlikely that significant portions of the aquatic community may be exposed to UV levels sufficient to induce phototoxicity, as UV levels occurring under normal sun light conditions have been shown to elicit these effects. There is growing evidence which suggests that phototoxic PAHs may be degrading aquatic habitats, particularly those in highly contaminated areas with shallow or clear water. Photo-induced chronic effects have been reported for anthracene at UV intensities occurring at depths of 10-12 m in Lake Michigan (Holst & Giesy, 1989). Phototoxicity of PAHs may also be initiated in aquatic organisms which have accumulated PAHs from the sediment and subsequently are exposed to sun light closer to the surface (The Netherlands, 2008). Phototoxic effects of PAHs are therefore considered relevant in this hazard, respectively T-assessment.

For phenanthrene acute as well as chronic toxicity data are available for fresh water algae, crustaceans including daphnids, fish, and insects. In addition, chronic toxicity data are also available for protozoans, cyanobacteria, aquatic plants, and insects.

1.1.1 Fish

1.1.1.1 Short-term toxicity to fish

A test performed with *Danio rerio* according to the OECD 236 FET test, provided an 96h-LC₅₀ of 11 µg/L (Alves et al., 2017). The 96h UV-enhanced LC₅₀ for zebrafish (*Danio rerio*) for phenanthrene was 271 µg/L (Willis and Oris 2014). The available data in the literature provides values ranging from up to 10 µg/L to phenanthrene solubility value and even higher (Black et al., 1983; Seiler et al., 2014; Turcotte et al., 2010; Hodson, 2017;

Verbruggen, 2012).

For marine water, data were available with red sea bream (*Pagrosomus major*) providing values of toxicity ranging from 0.15 mg/L for larvae NOEC to 48h-LC₅₀ of 1.97 mg/L for embryos (Zhao *et al.*, 2017). In turbot embryos (*Psetta maxima*), tested according to the early life stage OECD guideline, the values for hatching success 48h-EC₁₀ were of 102.5 and 42.9 µg/L under light and black condition respectively. The embryos survival 96h-EC₁₀ values were 4.5 and 6.66 µg/L under light and black condition respectively (Mhadhbi, Boumaiza and Beiras, 2010). Moreover, the FET with *Danio rerio* provides an effluent soluble fraction of 96h-LC₅₀ of 11 µg/L and a 96h-LC₅₀ of 4.9 µg/L for 50 % frequency of fish without an inflated swim bladder (Alves *et al.*, 2017).

1.1.1.2 Long-term toxicity to fish

Regarding long-term toxicity, data for fishes were assessed by Verbruggen and van Herwijnen in 2011 for the RIVM (Verbruggen and van Herwijnen, 2011) and updated by Verbruggen in 2012 (Verbruggen, 2012). In these reports, NOEC or EC₁₀ value ranging from 11 µg/L and 93 µg/L for mortality in an ELS test with the largemouth bass *Micropterus salmoides* and *Oryzias latipes* (based on the most sensitive parameters, malformations) in freshwater. The chronic toxicity data assessed by the RIVM for fish ranged from 11 µg/L to up to 560 µg/L. An assay with the zebrafish embryos larval survival assay provides a 30d-LC₁₀ value of 44 µg/L (Butler *et al.*, 2013) and was highlighted to be above 756 µg/L in 25d-LC₅₀ medaka embryos assay (Mu *et al.*, 2014). In marine water, NOEC and EC₁₀ values ranged from 68 to 168 µg/L (Verbruggen, 2012).

1.1.2 Aquatic invertebrates

Toxicity data of phenanthrene on aquatic invertebrates provides quite similar values when tested with daphnia (*Daphnia magna*). The available data, when performing acute tests, provide values ranging from 48h-EC₅₀ of 0.34 mg/L (OECD 202, Zindler *et al.*, 2016), 48h-EC₅₀ of 0.48 mg/L (Smith *et al.*, 2010), 48h-EC₅₀ of 0.55 mg/L (Zhang *et al.*, 2014) to 48h-EC₅₀ of 0.59 mg/L (Ma *et al.*, 2016). Verbruggen in 2012 report values ranging from 100 µg/L to 1200 µg/L in daphnia magna 48h-LC₅₀.

In marine environment, the model marine zooplankton *Artemia salina* provide a 48h-LC₅₀ of 0.49 mg/L (Lu *et al.*, 2018). A study with the mollusc *Mytilus galloprovincialis*, performed in the dark, gives a NOEC/EC₁₀ value of 29 µg/L and the crustacean *Acartia tonsa* a value of phenanthrene toxicity of 69 µg/L (Verbruggen, 2012). An acute 24h-EC₅₀ embryo-larval development assessment of phenanthrene on the pacific oyster *Crassostrea gigas* provide a value of 1.91µg/L and was performed according to the standard procedure AFNOR XP T90-382 (Afnor, 2009). Photo-enhanced toxicity of phenanthrene was assessed by exposing 3-days-old mysid shrimp (*Americamysis bahia*) under artificial ultraviolet light or in the dark during a 48-h acute toxicity test. The 96h-LC₅₀ was 22.8 µg/L and expressed no phototoxicity (Finch *et al.*, 2017).

Despite the recent literature search (April 2018), no new long term data were highlighted to be used to provide a more restrictive classification regarding the toxicity of phenanthrene. The data already assessed and described in the CTPHT and in the report of the RIVM are the most sensitive for phenanthrene on aquatic invertebrates. These reports provide the lowest EC₁₀ value for reproduction of *Ceriodaphnia dubia* in a 7d toxicity test. The value of this EC₁₀ is 13 µg/L and is based on measured concentrations.

1.1.3 Algae and aquatic plants

A 48h-EC₅₀ assay performed with *Pseudokirchneriella subcapitata* provides a toxicity value of 438.3µg/L. An OECD 201 algae assay performed with *T. chuii* at 20°C and 25°C provide value of 96h-IC₅₀ of 1.316 and 0.262 mg/L, respectively (Vieira and Guilhermino 2012).

For the marine microalgae *Phaeodactylum tricornutum*, the effect of phenanthrene was determined at 22 ± 2 °C and result in a 96 h IC₅₀ of 0.347 mg L⁻¹ (Okay and Karacik, 2007). For chronic toxicity data, the lowest 48h-EC₁₀ value available in the literature is 10 µg/L and was obtained for growth rate of the algae *P. subcapitata* (Halling-Sørensen *et al.*, 1996). The values of chronic toxicity to algae and macrophytes were ranging from EC₁₀ of 10 to 4910 µg/L (Verbruggen, 2012).

1.1.4 Sediment organisms

Phenanthrene toxicity to sediment was evidenced with different species (Evans and Nipper, 2007, Verbruggen and van Herwijnen 2011, ECHA, 2009). It is stated in CTPHT that “For sediment data are available for both fresh water sediment and marine sediment. The data for fresh water sediment include chronic tests with annelids, crustaceans and insects. The lowest NOEC is 50 mg/kg dw, recalculated to sediment with 10% organic carbon, for mortality and growth of both *Hyaella azteca* exposed for 14 days and *Chironomus riparius*, exposed for 10 days (Verrhiest *et al.*, 2001). Effect concentrations are based on measured concentrations. In another 28-d study with *Chironomus riparius*, emergence appeared to be somewhat less sensitive (Bleeker *et al.*, 2003).

For marine sediment toxicity data are available for two species of crustaceans. Effect concentrations for the Amphipod *Rhepoxynius abronius* (Swartz *et al.*, 1997; Boese *et al.*, 1998) appeared to be all above 200 mg/kg dw, recalculated to sediment with 10% organic carbon.” The most sensitive assay was performed with *Schizopera knabeni* and provides a NOEC/EC₁₀ value of 7.8 mg/kg dw based on the most sensitive parameter, the reproduction (Verbruggen, 2012).

1.2 Terrestrial compartment

The environmental hazard assessment for phenanthrene was previously done for the identification of CTPHT as SVHC (ECHA, 2009). It was reported that “chronic toxicity data for phenanthrene in soil are available for annelids, collembola, plants, crustaceans, and microbial processes. Again for phenanthrene, the EC₁₀ for reproduction of *Folsomia fimetaria* was the lowest EC₁₀ or NOEC (Sverdrup *et al.*, 2001, 2002, 2002c). Four EC₁₀ for *Folsomia* in sandy loam soil were available. The difference between the four values was the ageing of phenanthrene in the soil. Soils were spiked and toxicity testing started after 0, 10, 40, or 120 days after spiking. The EC_{10s} for these cases were 29, 18, 18, and 12 mg/kg dw, recalculated to a soil with 2% organic carbon. The EC_{10s} were based on measured concentrations. The geometric mean of these EC_{10s} is 18 mg/kg dw. This value was based on measured concentrations”.

This data is the lowest available for phenanthrene toxicity on soil organisms. Indeed, recent literature search (done in April 2018) highlight other phenanthrene toxicity data. Nevertheless, those data, described later, provide higher effect concentrations on different organisms. In a study with the predatory mite *Hypoaspis aculeifer*, performed according to the OECD guideline 226, the 14d-LC₅₀ was 684 mg/kg. When focusing on the reproduction, the 14d-EC₅₀ was 49 mg/kg and the avoidance 48h-EC₅₀ was 26 mg/kg (Owojori, Waszak, and Roembke, 2014). In an OECD 220 assay, the 21d-EC₁₀ was 37 mg/kg for *Enchytraeus crypticus* (Roelofs *et al.*, 2016). For *E. albidus*, the 3w-LC₅₀ was 135 mg/kg dry soil (Amorin *et al.*, 2011). According to an experiment realised in microcosm according to the ISO 11268 with *Eisenia fetida*, the 4w-EC₅₀ was 40.67 mg/kg (Wu *et al.*, 2012). One other study with *E. fetida* gives a NOEC/EC₁₀ value of 36 mg/kg (Verbruggen, 2012). A study with *Folsomia candida*, performed according to the ISO 11267, gives an EC₁₀ value of 24.95 mg/kg (Droge *et al.*, 2006).

1.3 Summary and discussion of the environmental hazard assessment

In the Support Document for identification of CTPHT as SVHC (ECHA, 2009), the issue has already been summarised and discussed as follows:

The experimental data indicate a high chronic and acute toxicity of the PAH constituents of CTPHT for aquatic organisms.

The reassessment of toxicity data by the RIVM (Verbruggen, 2012) and the recent literature search performed in April 2018 and included in the current report did not challenge this conclusion for phenanthrene.