

# Summary of product characteristics for a biocidal product

**Product name:** RATAACUM PELLETT

**Product type(s):** PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

**Authorisation number:** IT/2012/00062/AUT

**R4BP 3 asset reference number:** IT-0001128-0000

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
5. General directions for use	17
5.1. Instructions for use	17
5.2. Risk mitigation measures	18
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	19
5.4. Instructions for safe disposal of the product and its packaging	19
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	19
6. Other information	19

## Administrative information

### 1.1. Trade names of the product

RATACUM PELLETT
-----------------

### 1.2. Authorisation holder

<b>Name and address of the authorisation holder</b>	Name	Copyr S.p.A.
	Address	Via G. Stephenson 73 20157 Milano Italy
<b>Authorisation number</b>	IT/2012/00062/AUT	
<b>R4BP 3 asset reference number</b>	IT-0001128-0000	
<b>Date of the authorisation</b>	09/08/2012	
<b>Expiry date of the authorisation</b>	01/07/2024	

### 1.3. Manufacturer(s) of the biocidal products

<b>Name of the manufacturer</b>	Copyr S.p.A.
<b>Address of the manufacturer</b>	Via G. Stephenson, 73 20157 Milano Italy
<b>Location of manufacturing sites</b>	Indupharma S.r.l. - Via Sorgaglia, 40 35020 Arre (PD) Italy

### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	26 - Difenacoum
<b>Name of the manufacturer</b>	ACTIVA SRL
<b>Address of the manufacturer</b>	Via Feltre 32 20132 Milan Italy
<b>Location of manufacturing sites</b>	Tezza S.r.l.-via Tre Ponti 22 37050 S. Maria di Zevio Italy

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Difenacoum	3-(3-biphenyl-4-yl-1,2,3,4- tetrahydro-1-naphthyl)-4-hydroxycoumarin	Active Substance	56073-07-5	259-978-4	0,005
Vanilla aroma		Non-active substance			0,3
Green Pigment		Non-active substance			0,08

### 2.2. Type of formulation

RB - Bait (ready for use)
---------------------------

## 3. Hazard and precautionary statements

<b>Hazard statements</b>	<p>May damage the unborn child.</p> <p>May cause damage to organs to BLOOD through prolonged or repeated exposure .</p>
<b>Precautionary statements</b>	<p>Keep out of reach of children.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Wear protective gloves.</p> <p>IF exposed or concerned: Get medical advice.</p> <p>Dispose of contents to in accordance with national and international regulations.</p> <p>Do not breathe dust.</p>

## 4. Authorised use(s)

### 4.1 Use description

#### Use 1 - House mice - professionals - indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: juveniles to adult
<b>Field(s) of use</b>	Indoor  Indoor
<b>Application method(s)</b>	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
<b>Application rate(s) and frequencies</b>	Application Rate: 40g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): 0 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
<b>Category(ies) of users</b>	Industrial  Professional
<b>Pack sizes and packaging material</b>	<b>Type of packaging</b> <b>Size/volume of the packaging</b> <b>Material of the packaging</b> <b>Type and material of closure(s)</b> <b>Intended user (e.g. professional, non-professional)</b> <b>Compatibility of the product with the proposed packaging materials (Yes/No)</b> Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled or Printed Bag/Sack - with Inner plastic liner

from 1.5 Kg - to 10 kg - max size inner liner 10 Kg  
Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes  
Labelled Plastic Bucket  
from 1.5 Kg -to 25 kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes  
Labelled Plastic Bucket- with inner plastic liner  
from 1.5 - to 10 kg - max size inner liner 10 Kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes  
Labelled or Printed Carton sachet with plastic inner liner  
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes  
Labelled or Printed Cardboard with inner plastic liner  
1,5-10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated cardboard  
Professional, trained professional  
Yes

#### 4.1.1 Use-specific instructions for use

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

#### 4.1.2 Use-specific risk mitigation measures

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

#### 4.2 Use description

##### Use 2 - Mice and rats - trained professionals - outdoor around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	Scientific name: Rattus norvegicus Common name: Brown rat Development stage: juvenile to adult  Scientific name: Rattus rattus Common name: Roof rat Development stage: other: juvenile to adult  Scientific name: Mus musculus Common name: House mouse Development stage: other: juvenile to adult
<b>Field(s) of use</b>	Outdoor  outdoor around buildings
<b>Application method(s)</b>	Method: Bait application Detailed description: Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations
<b>Application rate(s) and frequencies</b>	Application Rate: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point. Dilution (%): 0 Number and timing of application: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.
<b>Category(ies) of users</b>	

**Pack sizes and packaging material**

Trained professional
<p><b>Type of packaging</b> <b>Size/volume of the packaging</b> <b>Material of the packaging</b> <b>Type and material of closure(s)</b> <b>Intended user (e.g. professional, non-professional)</b> <b>Compatibility of the product with the proposed packaging materials (Yes/No)</b></p> <p>Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled Plastic Bucket from 1.5 Kg -to 25 kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes</p> <p>Labelled Plastic Bucket- with inner plastic liner from 1.5 - to 10 kg - max size inner liner 10 Kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes</p> <p>Labelled or Printed Carton sachet with plastic inner liner from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Cardboard with inner plastic liner 1,5-10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated cardboard Professional, trained professional Yes</p>

**4.2.1 Use-specific instructions for use**

Remove the remaining product at the end of treatment period.  
For permanent baiting :  
- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.  
- Follow any additional instructions provided by the relevant code of best practice.

For application in covered and protected bait points:



- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Follow any additional instructions provided by the relevant code of best practice.

#### 4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product in pulsed baiting treatments.
- do not apply directly in burrows

In case of permanent baiting:

- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

### 4.3 Use description

#### Use 3 - Rats - professionals - indoor

**Product type**

PT14 - Rodenticides (Pest control)

**Where relevant, an exact description of the authorised use**

Not relevant for rodenticides

**Target organism(s) (including development stage)**

Scientific name: Rattus norvegicus  
Common name: Brown rat  
Development stage: juvenile to adult

Scientific name: Rattus rattus  
Common name: Roof rat  
Development stage: other: juvenile to adult

**Field(s) of use**

Indoor

Indoor

**Application method(s)**

Method: Bait application  
Detailed description:  
Ready-to-use bait to be used in tamper-resistant bait stations

**Application rate(s) and frequencies**

Application Rate: 60-100g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.  
Dilution (%): 0  
Number and timing of application:  
If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.

**Category(ies) of users**

Industrial

Professional

**Pack sizes and packaging material**

**Type of packaging**  
**Size/volume of the packaging**  
**Material of the packaging**  
**Type and material of closure(s)**  
**Intended user (e.g. professional, non-professional)**  
**Compatibility of the product with the proposed packaging materials (Yes/No)**

Labelled or Printed Bottle  
from 1.5 Kg- to 5 kg  
Plastic composite (HDPE)  
Bottle sealed  
Professional, trained professional  
Yes

Labelled or Printed Bag/Sack  
from 1.5 Kg - to 25 kg  
Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes

Labelled or Printed Bag/Sack - with Inner plastic liner  
from 1.5 Kg - to 10 kg - max size inner liner 10 Kg  
Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes

Labelled Plastic Bucket  
from 1.5 Kg -to 25 kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes

Labelled Plastic Bucket- with inner plastic liner  
from 1.5 - to 10 kg - max size inner liner 10 Kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed

Professional, trained professional  
Yes  
Labelled or Printed Carton sachet with plastic inner liner  
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes  
Labelled or Printed Cardboard with inner plastic liner  
1,5-10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated cardboard  
Professional, trained professional  
Yes

#### 4.3.1 Use-specific instructions for use

The bait stations should be visited at least every 5 to 7 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

#### 4.3.2 Use-specific risk mitigation measures

#### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

#### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

#### 4.4 Use description

Use 4 - House mice and rats - professionals -outdoor around buildings

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: juvenile to adult</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: other: juvenile to adult</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: other: juvenile to adult</p>
<b>Field(s) of use</b>	<p>Outdoor</p> <p>outdoor around buildings</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: House mouse: 40g of bait per bait point. Rattus: 60-100g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): 0 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.</p>
<b>Category(ies) of users</b>	<p>Industrial</p> <p>Professional</p>
<b>Pack sizes and packaging material</b>	<p>M</p> <p><b>Type of packaging</b> <b>Size/volume of the packaging</b> <b>Material of the packaging</b> <b>Type and material of closure(s)</b> <b>Intended user (e.g. professional, non-professional)</b> <b>Compatibility of the product with the proposed packaging materials (Yes/No)</b></p> <p>Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded</p>

Professional, trained professional  
Yes  
Labelled Plastic Bucket  
from 1.5 Kg -to 25 kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes  
Labelled Plastic Bucket- with inner plastic liner  
from 1.5 - to 10 kg - max size inner liner 10 Kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes  
Labelled or Printed Carton sachet with plastic inner liner  
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes  
Labelled or Printed Cardboard with inner plastic liner  
1,5-10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated cardboard  
Professional, trained professional  
Yes

#### 4.4.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.  
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.  
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

#### 4.4.2 Use-specific risk mitigation measures

Do not apply this product directly in the burrows.

#### 4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### 4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

#### 4.5 Use description

##### Use 5 - House mice and rats - trained professionals - indoor

<b>Product type</b>	PT14 - Rodenticides (Pest control)
<b>Where relevant, an exact description of the authorised use</b>	Not relevant for rodenticides
<b>Target organism(s) (including development stage)</b>	<p>Scientific name: Rattus norvegicus Common name: Brown rat Development stage: juvenile to adult</p> <p>Scientific name: Rattus rattus Common name: Roof rat Development stage: other: juvenile to adult</p> <p>Scientific name: Mus musculus Common name: House mouse Development stage: other: juvenile to adult</p>
<b>Field(s) of use</b>	<p>Indoor</p> <p>indoor</p>
<b>Application method(s)</b>	<p>Method: Bait application Detailed description: Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations.</p>
<b>Application rate(s) and frequencies</b>	<p>Application Rate: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point. Dilution (%): 0 Number and timing of application: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.</p>
<b>Category(ies) of users</b>	Trained professional

## Pack sizes and packaging material

**Type of packaging**  
**Size/volume of the packaging**  
**Material of the packaging**  
**Type and material of closure(s)**  
**Intended user (e.g. professional, non-professional)**  
**Compatibility of the product with the proposed packaging materials (Yes/No)**

Labelled or Printed Bottle  
from 1.5 Kg- to 5 kg  
Plastic composite (HDPE)  
Bottle sealed  
Professional, trained professional  
Yes

Labelled or Printed Bag/Sack  
from 1.5 Kg - to 25 kg  
Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes

Labelled or Printed Bag/Sack - with Inner plastic liner  
from 1.5 Kg - to 10 kg - max size inner liner 10 Kg  
Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes

Labelled Plastic Bucket  
from 1.5 Kg -to 25 kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes

Labelled Plastic Bucket- with inner plastic liner  
from 1.5 - to 10 kg - max size inner liner 10 Kg  
Labelled plastic (HDPE) bucket  
Rectangular or truncated cone bucket sealed  
Professional, trained professional  
Yes

Labelled or Printed Carton sachet with plastic inner liner  
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated bags or serial production bags , both thermal welded  
Professional, trained professional  
Yes

Labelled or Printed Cardboard with inner plastic liner  
1,5-10 kg single inner plastic bags of up to 10 Kg each  
Carton + Plastic composite  
Prefabricated cardboard  
Professional, trained professional  
Yes

### 4.5.1 Use-specific instructions for use

Remove the remaining product at the end of treatment period.  
For permanent baiting :  
- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.  
- Follow any additional instructions provided by the relevant code of best practice.

### 4.5.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product in pulsed baiting treatments.

In case of permanent baiting:

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

#### 4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage system, ensure that bait contact with water is avoided.

#### 4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

#### 4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

#### 4.6 Use description

##### Use 6 - Rats - trained professionals - Outdoor open areas

**Product type**

PT14 - Rodenticides (Pest control)

**Where relevant, an exact description of the authorised use**

Not relevant for rodenticides

**Target organism(s) (including development stage)**

Scientific name: *Rattus norvegicus*  
Common name: Brown rat  
Development stage: juvenile to adult

Scientific name: *Rattus rattus*  
Common name: Roof rat  
Development stage: other: juvenile to adult

**Field(s) of use**



<p><b>Application method(s)</b></p>	<p>Outdoor</p> <p>outdoor open areas</p> <hr/> <p>Method: Bait application  Detailed description:  Bait formulations:  - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations</p>
<p><b>Application rate(s) and frequencies</b></p>	<p>Application Rate: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.  Dilution (%): 0  Number and timing of application:  Bait products  Rats  - High infestation: 100 g of bait per baiting point.  - Low infestation: 60 g of bait per baiting point.</p> <p>Permanent baiting 60-100 gr of bait per bait point.</p>
<p><b>Category(ies) of users</b></p>	<p>Trained professional</p>
<p><b>Pack sizes and packaging material</b></p>	<p><b>Type of packaging</b>  <b>Size/volume of the packaging</b>  <b>Material of the packaging</b>  <b>Type and material of closure(s)</b>  <b>Intended user (e.g. professional, non-professional)</b>  <b>Compatibility of the product with the proposed packaging materials (Yes/No)</b></p> <p>Labelled or Printed Bottle  from 1.5 Kg- to 5 kg  Plastic composite (HDPE)  Bottle sealed  Professional, trained professional  Yes</p> <p>Labelled or Printed Bag/Sack  from 1.5 Kg - to 25 kg  Plastic composite  Prefabricated bags or serial production bags , both thermal welded  Professional, trained professional  Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner  from 1.5 Kg - to 10 kg - max size inner liner 10 Kg  Plastic composite  Prefabricated bags or serial production bags , both thermal welded  Professional, trained professional  Yes</p> <p>Labelled Plastic Bucket  from 1.5 Kg -to 25 kg  Labelled plastic (HDPE) bucket  Rectangular or truncated cone bucket sealed  Professional, trained professional  Yes</p> <p>Labelled Plastic Bucket- with inner plastic liner  from 1.5 - to 10 kg - max size inner liner 10 Kg  Labelled plastic (HDPE) bucket  Rectangular or truncated cone bucket sealed</p>

Professional, trained professional  
 Yes  
 Labelled or Printed Carton sachet with plastic inner liner  
 from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each  
 Carton + Plastic composite  
 Prefabricated bags or serial production bags , both thermal welded  
 Professional, trained professional  
 Yes  
 Labelled or Printed Cardboard with inner plastic liner  
 1,5-10 kg single inner plastic bags of up to 10 Kg each  
 Carton + Plastic composite  
 Prefabricated cardboard  
 Professional, trained professional  
 Yes

#### 4.6.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- Follow any additional instructions provided by the relevant code of best practice.

For permanent baiting :

- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.
- Follow any additional instructions provided by the relevant code of best practice.

For application in covered and protected bait points:

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Follow any additional instructions provided by the relevant code of best practice.

#### 4.6.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
  - Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
  - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
  - Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
  - Do not use the product in pulsed baiting treatments.
  - do not apply directly in burrows
- In case of permanent baiting:
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

#### 4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

#### **4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging**

#### **4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

## **5. General directions for use**

### **5.1. Instructions for use**

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves (i.e rubber) during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.- Instructions for use that are "bait-specific":-Loose baits: place the bait in the bait station by using a dosage device. Bait should not be decanted, avoid breath of dust

## 5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to general public (i.e. 2 for professionals only)
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment unless authorised for permanent baiting treatments.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements.

### 5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
  - Dermal exposure, wash skin with water and then with water and soap.
  - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
  - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre
- Hazardous to wildlife.

### 5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements

### 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

## 6. Other information

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.